

# The TP53 Arg72Pro and MDM2 309G > T polymorphisms are not associated with breast cancer risk in BRCA1 and BRCA2 mutation carriers

**OM Sinilnikova<sup>\*,1,2</sup>, AC Antoniou<sup>3</sup>, J Simard<sup>4</sup>, S Healey<sup>5</sup>, M Léoné<sup>1</sup>, D Sinnett<sup>6,7</sup>, AB Spurdle<sup>5</sup>, J Beesley<sup>5</sup>, X Chen<sup>5</sup>, kConFab<sup>8</sup>, MH Greene<sup>9</sup>, JT Loud<sup>9</sup>, F Lejbkowitz<sup>10</sup>, G Rennert<sup>10</sup>, S Dishon<sup>10</sup>, IL Andrusis<sup>11,12</sup>, OCGN<sup>11</sup>, SM Domchek<sup>13</sup>, KL Nathanson<sup>13</sup>, S Manoukian<sup>14</sup>, P Radice<sup>15,16</sup>, I Konstantopoulou<sup>17</sup>, I Blanco<sup>18</sup>, AL Laborde<sup>19</sup>, M Durán<sup>20</sup>, A Osorio<sup>21</sup>, J Benitez<sup>21</sup>, U Hamann<sup>22</sup>, FBL Hogervorst<sup>23</sup>, TAM van Os<sup>24</sup>, HJP Gille<sup>25</sup>, HEBON<sup>23</sup>, S Peock<sup>3</sup>, M Cook<sup>3</sup>, C Luccarini<sup>26</sup>, DG Evans<sup>27</sup>, F Laloo<sup>27</sup>, R Eeles<sup>28</sup>, G Pichert<sup>29</sup>, R Davidson<sup>30</sup>, T Cole<sup>31</sup>, J Cook<sup>32</sup>, J Paterson<sup>33</sup>, C Brewer<sup>34</sup>, EMBRACE<sup>3</sup>, DJ Hughes<sup>35</sup>, I Couplier<sup>36,37</sup>, S Giraud<sup>1</sup>, F Coulet<sup>38</sup>, C Colas<sup>38</sup>, F Soubrier<sup>38</sup>, E Rouleau<sup>39</sup>, I Bièche<sup>39</sup>, R Lidereau<sup>39</sup>, L Demange<sup>40</sup>, C Nogues<sup>40</sup>, HT Lynch<sup>41</sup>, GEMO<sup>1,2,42</sup>, RK Schmutzler<sup>43</sup>, B Versmold<sup>43</sup>, C Engel<sup>44</sup>, A Meindl<sup>45</sup>, N Arnold<sup>46</sup>, C Sutter<sup>47</sup>, H Deissler<sup>48</sup>, D Schaefer<sup>49</sup>, UG Froster<sup>50</sup>, GC-HBOC<sup>43,44,45,46,47,48,49,50</sup>, K Aittomäki<sup>51</sup>, H Nevanlinna<sup>52</sup>, L McGuffog<sup>3</sup>, DF Easton<sup>3</sup>, G Chenevix-Trench<sup>5</sup> and D Stoppa-Lyonnet<sup>42</sup> on behalf of the Consortium of Investigators of Modifiers of BRCA1/2**

<sup>1</sup>Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Hospices Civils de Lyon / Centre Léon Bérard, Lyon 69373, France; <sup>2</sup>Laboratoire de Génétique Moléculaire, Signalisation et Cancer, UMR5201 CNRS, Université de Lyon, Lyon 69373, France; <sup>3</sup>Cancer Research UK, Genetic Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge CB1 8RN, UK; <sup>4</sup>Canada Research Chair in Oncogenetics, Cancer Genomics Laboratory, Centre Hospitalier Universitaire de Quebec and Laval University, Quebec, Canada G1V 4G2; <sup>5</sup>Queensland Institute of Medical Research, Brisbane QLD 4029, Australia; <sup>6</sup>Division of Hematology-Oncology, Research Center, CHU Sainte-Justine, Montreal, Canada H3T 1C5; <sup>7</sup>Department of Pediatrics, University of Montreal, Montreal, Canada H3T 1C5; <sup>8</sup>Peter MacCallum Cancer Center, Melbourne, VIC 3002, Australia; <sup>9</sup>Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD 20852-7231, USA; <sup>10</sup>CHS National Cancer Control Center and Department of Community Medicine and Epidemiology, Carmel Medical Center and Technion Faculty of Medicine, Haifa 34362, Israel; <sup>11</sup>Ontario Cancer Genetics Network, Cancer Care Ontario, Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada M5G 1X5; <sup>12</sup>Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, ON, Canada M5G 1X5; <sup>13</sup>Department of Medicine, Abramson Cancer Center, University of Pennsylvania School of Medicine, Philadelphia, PA 19104, USA; <sup>14</sup>Unit of Medical Genetics, Fondazione IRCCS Istituto Nazionale Tumori, Milano, Italy; <sup>15</sup>Department of Experimental Oncology, Fondazione IRCCS Istituto Nazionale Tumori, Milano 20133, Italy; <sup>16</sup>IFOM, Fondazione Istituto FIRC di Oncologia Molecolare, Milano 20139, Italy; <sup>17</sup>Molecular Diagnostics Laboratory, IRRP, NCSR 'Demokritos', 15310 Ag. Paraskevi, Athens, Greece; <sup>18</sup>Cancer Genetic Counseling Program, Institut Català d'Oncologia, Barcelona 08907, Spain; <sup>19</sup>Servicio de Genética, Hospital de la Santa Creu i Sant Pau, Barcelona 08025, Spain; <sup>20</sup>Laboratorio de Genética del Cáncer, Instituto de Biología y Genética Molecular, Universidad de Valladolid, Valladolid 47002, Spain; <sup>21</sup>Human Cancer Genetics Programme, Spanish National Cancer Research Centre (CNIO), Madrid 28029, Spain; <sup>22</sup>Deutsches Krebsforschungszentrum (DKFZ), Molecular Genetics of Breast Cancer, Heidelberg 69120, Germany; <sup>23</sup>Department of Pathology and the Family Cancer Clinic, Netherlands Cancer Institute, Amsterdam 1066 CX, The Netherlands; <sup>24</sup>Department of Clinical Genetics, Amsterdam University Medical Centre, Amsterdam 1105 AZ, The Netherlands; <sup>25</sup>Department of Clinical Molecular Genetics, Free University Medical Centre, Amsterdam 1081 HV, The Netherlands; <sup>26</sup>Department of Oncology, University of Cambridge, Cambridge CB1 8RN, UK; <sup>27</sup>Manchester Academic Health Sciences Centre, Central Manchester University Hospitals NHS Foundation Trust, Manchester M13 0JH, UK; <sup>28</sup>Translational Cancer Genetics Team, The Institute of Cancer Research and Royal Marsden NHS Foundation Trust, Sutton, Surrey SM2 5NG, UK; <sup>29</sup>Clinical Genetics, Guy's Hospital, London SE1 9RT, UK; <sup>30</sup>Ferguson-Smith Centre for Clinical Genetics, Glasgow G3 8SJ, UK; <sup>31</sup>West Midlands Regional Genetics Service, Birmingham Women's Hospital Healthcare NHS Trust, Edgbaston, Birmingham B15 2TG, UK; <sup>32</sup>Sheffield Clinical Genetics Service, Sheffield Children's Hospital, Sheffield S10 2TH, UK; <sup>33</sup>Department of Clinical Genetics, East Anglian Regional Genetics Service, Addenbrookes Hospital, Cambridge CB2 8QE, UK; <sup>34</sup>Department of Clinical Genetics, Royal Devon & Exeter Hospital, Exeter EX2 5DW, UK; <sup>35</sup>Department of Clinical Medicine, Trinity College Centre for Health Sciences, Adelaide & Meath Hospital, Dublin 24, Ireland; <sup>36</sup>Service de Génétique Médicale, Unité d'Oncogénétique, CHU Anaud de Villeneuve, Montpellier 34295, France; <sup>37</sup>Unité d'Oncogénétique, CRLCC Val d'Aurelle, Montpellier 34295, France; <sup>38</sup>Groupe Hospitalier Pitié-Salpêtrière, Assistance Publique-Hopitaux de Paris, Laboratoire d'Oncogénétique et Angiogénétique Moléculaire, Université Pierre et Marie Curie, Paris 75651, France; <sup>39</sup>INSERM U735 / Oncogenetics, Centre René Huguenin, Saint-Cloud 92210, France; <sup>40</sup>Epidémiologie Clinique, Oncogénétique, Centre René Huguenin, Saint-Cloud 92210, France; <sup>41</sup>Department of Preventive Medicine, Creighton University, Omaha, NE 68178, USA; <sup>42</sup>INSERM U509, Service de Génétique Oncologique, Institut Curie, Université Paris-Descartes, Paris 75248, France; <sup>43</sup>Centre for hereditary Breast and Ovarian Cancer, Department of Obstetrics and Gynaecology, University of Cologne, Cologne 50924, Germany; <sup>44</sup>Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Leipzig 04103, Germany; <sup>45</sup>Department of Obstetrics and Gynaecology, Technical University Munich, Munich 80333, Germany; <sup>46</sup>Department of Obstetrics and Gynaecology, University of Schleswig-Holstein, Kiel 24105, Germany; <sup>47</sup>Institute of Human Genetics, University of Heidelberg, Heidelberg 69117, Germany; <sup>48</sup>Department of Obstetrics and Gynaecology, University of Ulm, Ulm 89069, Germany; <sup>49</sup>Institute of Human Genetics, University of Frankfurt, Frankfurt 60325, Germany; <sup>50</sup>Institute of Human Genetics, University of Leipzig, Leipzig 04103, Germany; <sup>51</sup>Department of Clinical Genetics, Helsinki University Central Hospital, Helsinki, 00029 HUS, Finland; <sup>52</sup>Department of Obstetrics and Gynecology, Helsinki University Central Hospital, Helsinki, 00029 HUS, Finland

\*Correspondence: Dr OM Sinilnikova; E-mail: sinilnik@lyon.fnclcc.fr

Received 17 April 2009; revised 15 July 2009; accepted 27 July 2009; published online 25 August 2009

**BACKGROUND:** The *TP53* pathway, in which *TP53* and its negative regulator *MDM2* are the central elements, has an important role in carcinogenesis, particularly in *BRCA1*- and *BRCA2*-mediated carcinogenesis. A single nucleotide polymorphism (SNP) in the promoter region of *MDM2* (309T>G, rs2279744) and a coding SNP of *TP53* (Arg72Pro, rs1042522) have been shown to be of functional significance.

**METHODS:** To investigate whether these SNPs modify breast cancer risk for *BRCA1* and *BRCA2* mutation carriers, we pooled genotype data on the *TP53* Arg72Pro SNP in 7011 mutation carriers and on the *MDM2* 309T>G SNP in 2222 mutation carriers from the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA). Data were analysed using a Cox proportional hazards model within a retrospective likelihood framework.

**RESULTS:** No association was found between these SNPs and breast cancer risk for *BRCA1* (*TP53*: per-allele hazard ratio (HR) = 1.01, 95% confidence interval (CI): 0.93–1.10,  $P_{\text{trend}} = 0.77$ ; *MDM2*: HR = 0.96, 95%CI: 0.84–1.09,  $P_{\text{trend}} = 0.54$ ) or for *BRCA2* mutation carriers (*TP53*: HR = 0.99, 95%CI: 0.87–1.12,  $P_{\text{trend}} = 0.83$ ; *MDM2*: HR = 0.98, 95%CI: 0.80–1.21,  $P_{\text{trend}} = 0.88$ ). We also evaluated the potential combined effects of both SNPs on breast cancer risk, however, none of their combined genotypes showed any evidence of association.

**CONCLUSION:** There was no evidence that *TP53* Arg72Pro or *MDM2* 309T>G, either singly or in combination, influence breast cancer risk in *BRCA1* or *BRCA2* mutation carriers.

*British Journal of Cancer* (2009) **101**, 1456–1460. doi:10.1038/sj.bjc.6605279 www.bjcancer.com

Published online 25 August 2009

© 2009 Cancer Research UK

**Keywords:** *TP53*; *MDM2*; *BRCA1/2*; breast cancer; polymorphism; risk

The *TP53* pathway is crucial for tumour suppression, acting through regulation of cell-cycle control, apoptosis, senescence and DNA repair. The *TP53* gene and its negative regulator *MDM2* are central to this pathway, promoting polyubiquitination and degradation of *TP53*, and also controlling the *TP53* synthesis (Toledo and Wahl, 2006; Candeias *et al*, 2008). Inactivation of the *TP53* pathway has an important role in *BRCA1*- and *BRCA2*-associated tumorigenesis. *BRCA1* and *BRCA2* mutations are associated with genomic instability caused by defective cell-cycle checkpoint and DNA damage repair (Deng, 2006). Mouse model studies have highlighted functional links between these genes. Biallelic inactivation of *BRCA1* and *BRCA2* in mice have shown that embryonic lethality because of growth retardation can be partially rescued in a *Trp53* null background (Evers and Jonkers, 2006). The development of mammary tumours in conditional *BRCA1* and *BRCA2* knockout mice was considerably accelerated in a *Trp53* knockout background (Evers and Jonkers, 2006). In addition, a high incidence of *TP53* mutations has been found in breast tumours of human *BRCA1* and *BRCA2* mutation carriers (Greenblatt *et al*, 2001; Manie *et al*, 2009). The observed interactions between *TP53* and *BRCA* pathways are integral to the progression of tumourigenesis in breast cancer.

A *TP53* polymorphism (rs1042522) has been found to be of functional significance, with the Pro72 allele being less efficient than Arg72 at inducing apoptosis, mainly due to weaker binding and ubiquitination by *MDM2* of the Pro72 variant protein (Dumont *et al*, 2003; Osorio *et al*, 2006). An SNP in the promoter region of *MDM2* (309T>G, rs2279744) has been shown to increase *MDM2* transcriptional activity, thus attenuating the *TP53* pathway (Bond *et al*, 2004). This latter SNP was associated with an earlier onset of breast cancer in Li-Fraumeni patients carrying *TP53* mutations (Bougeard *et al*, 2006; Ruijs *et al*, 2007). The effect on breast cancer risk of the *TP53* Arg72Pro and the *MDM2* 309T>G polymorphisms, separately and in combination, was investigated in a large case-control study by the Breast Cancer Association Consortium (BCAC), but no association was detected (Schmidt *et al*, 2007). However, several smaller studies examined these polymorphisms in *BRCA1* and *BRCA2* mutation carriers (Martin *et al*, 2003; Tommiska *et al*, 2005; Copson *et al*, 2006; Osorio *et al*, 2006; Wasielewski *et al*, 2007; Yarden *et al*, 2008), and some suggested an association between the *TP53* Pro72 and the *MDM2*

309G alleles with an earlier age at breast cancer diagnosis (Martin *et al*, 2003; Tommiska *et al*, 2005; Osorio *et al*, 2006; Yarden *et al*, 2008). We therefore investigated the associations between breast cancer risk and these *TP53* and *MDM2* polymorphisms in a large series of *BRCA1* and *BRCA2* mutation carriers from the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA) (Chenevix-Trench *et al*, 2007).

## MATERIALS AND METHODS

### Study sample

Eligibility was restricted to female carriers with pathogenic mutations in *BRCA1* or *BRCA2* who were  $\geq 18$  years. Data were obtained from 13 CIMBA studies (Table 1). The majority of carriers were recruited through cancer genetics clinics offering genetic testing, and enrolled into national or regional studies. Information collected included the year of birth; mutation description; age at last followup; ages at breast and ovarian cancer diagnosis; and age at bilateral prophylactic mastectomy. Information was also available on the country of residence, which was defined to be the country of the clinic at which the carrier family was recruited for the study. Related individuals were identified through a unique family identifier. Further details of the information collected on the *BRCA1* and *BRCA2* mutation carriers and other details of the CIMBA initiative can be found elsewhere. Additional specific acknowledgements to the CIMBA collaborating centres are included in the Supplementary Appendix. (<http://www.srl.cam.ac.uk/consortia/cimba/index.html>) (Chenevix-Trench *et al*, 2007). All carriers participated in clinical and research studies at the host institutions under IRB-approved protocols.

### Genotyping

We pooled genotype data from studies within CIMBA that had previously genotyped polymorphisms rs1042522 and rs2279744 (see Table 1). Deviation from Hardy-Weinberg equilibrium among unrelated subjects was evaluated separately for each SNP and study. There was evidence for deviation for only one study ( $P = 0.03$ ), but cluster plot examination did not show any unusual

**Table 1** Number of *BRCA1* and *BRCA2* mutation carriers by study and by single nucleotide polymorphism (SNP)

Study	Country	TP53 Arg72Pro (rs1042522)	MDM2 309T > G (rs2279744)	Genotyping platform
Spanish National Cancer Centre (CNIO)	Spain	788	0	Restriction enzyme digestion
Deutsches Krebsforschungszentrum (DKFZ)	Germany	170	0	PCR-based RFLP
Epidemiological study of <i>BRCA1</i> and <i>BRCA2</i> mutation carriers (EMBRACE)	U.K. and Eire	1131	0	iPLEX
Genetic Modifiers of cancer risk in <i>BRCA1/2</i> mutation carriers (GEMO)	France and U.S.A.	1405	1357	Taqman
German Consortium of Hereditary Breast and Ovarian Cancer (GC-HBOC)	Germany	815	0	Taqman
Helsinki Breast Cancer Study (HEBCS)	Finland	188	187	rs1042522: Amplifluor(tm) fluorescent genotyping (Kbiosciences); rs2279744: RFLP
HEreditary Breast and Ovarian study Netherlands (HEBON)	The Netherlands	438	432	Taqman
INterdisciplinary HEalth Research International Team BREast CANcer susceptibility (INHERIT BRCA5)	Quebec-Canada	146	155	Taqman
kConFab	Australia	790	0	iPLEX
National Cancer Institute (NCI)	USA	190	0	Taqman
National Israeli Cancer Control Center (NICCC)	Israel	470	0	Taqman
Ontario Cancer Genetics Network (OCGN)	Canada	84	91	Taqman
University of Pennsylvania (UPENN)	USA	396	0	iPLEX
Total		7011	2222	

pattern and the study was included in the analysis. Where available study specific genotyping quality control data were examined and data were included if the call rate was over 95% and the concordance among duplicates was over 98%.

**Statistical analysis**

Mutation carriers were classified according to their age at diagnosis of breast cancer or their age at last follow up. For this purpose, individuals were censored at the age of first breast cancer diagnosis, ovarian cancer diagnosis, bilateral prophylactic mastectomy or the age at last observation. Only individuals censored at breast cancer diagnosis were assumed to be affected (Table 2).

To correct for a potential bias related to the fact that *BRCA1* and *BRCA2* mutation carriers are not randomly sampled with respect to their disease status, the data were analysed within a survival analysis framework, by modelling the retrospective likelihood of the observed genotypes conditional on the disease phenotypes. A detailed description of the retrospective likelihood approach has been published (Antoniou et al, 2007). We used a Cox proportional hazards model, where the effect of each SNP was modelled either as a per-allele hazard ratio (HR) or using separate HRs for heterozygotes and homozygotes. To assess the combined effects of the SNPs, we fitted a model in which a separate HR parameter was estimated for each multilocus genotype. More details of the statistical analysis can be found elsewhere (Antoniou et al, 2008).

**RESULTS**

In total, 7011 *BRCA1* and *BRCA2* mutation carriers were genotyped for *TP53* Arg72Pro and 2222 mutation carriers were genotyped for *MDM2* 309T > G (Table 1). Table 2 shows summary statistics for the cohort of *BRCA1* and *BRCA2* mutation carriers with an observed genotype for either the *TP53* or *MDM2* polymorphism. There was no evidence of an association between either SNP and breast cancer risk in *BRCA1* or *BRCA2* mutation carriers combined or analysed separately (*TP53* Arg72Pro:  $P_{trend} = 0.89, 0.77$  and  $0.83$ , respectively; *MDM2* 309T > G:  $P_{trend} = 0.60, 0.54$  and  $0.88$ , respectively) (Table 3). There was no evidence for heterogeneity in the HRs between studies (*TP53* Arg72Pro:  $P = 0.22$  and  $0.93$ , *MDM2* 309T > G:  $P = 0.11$  and  $0.82$  for *BRCA1* or *BRCA2* mutation carriers

**Table 2** Summary characteristics for the 7109 eligible *BRCA1* and *BRCA2* carriers used in the analysis and typed for either single nucleotide polymorphism (SNP)

Characteristic	BRCA1		BRCA2	
	Unaffected	Breast cancer	Unaffected	Breast cancer
Number	2055	2567	1051	1436
Person-years follow-up	87571	104679	46315	63080
Median age at censure (IQR)	41 (33–51)	40 (34–46)	42 (34–52)	43 (37–50)
Age at censure (years), N (%)				
<30	327 (15.9)	225 (8.8)	139 (13.2)	78 (5.4)
30–39	584 (28.4)	1052 (41.0)	296 (28.1)	462 (32.2)
40–49	574 (27.9)	880 (34.3)	286 (27.2)	511 (35.6)
50–59	364 (17.7)	296 (11.5)	196 (18.7)	278 (19.4)
60–69	134 (6.5)	87 (3.4)	82 (7.8)	81 (5.6)
70+	72 (3.5)	27 (1.0)	52 (4.9)	26 (1.8)
Year of birth, N (%)				
<1920	18 (0.9)	32 (1.3)	12 (1.1)	10 (0.7)
1920–29	63 (3.1)	117 (4.6)	39 (3.7)	83 (5.8)
1930–39	171 (8.3)	267 (10.4)	96 (9.1)	196 (13.7)
1940–49	326 (15.9)	657 (25.6)	143 (13.6)	358 (24.9)
1950–59	481 (23.4)	820 (31.9)	241 (22.9)	459 (32.0)
1960+	996 (48.5)	674 (26.3)	520 (49.5)	330 (23.0)

Abbreviation: IQR = interquartile range.

respectively). The HRs for the 9 *TP53*–*MDM2* combined genotypes, estimated separately in *BRCA1* and *BRCA2* mutation carriers, ranged between 0.72 and 1.31, but none of them were significant.

**DISCUSSION**

To our knowledge, this is the largest study to investigate the hypothesis that *TP53* Arg72Pro and *MDM2* 309T > G influence breast cancer risk in *BRCA1* and *BRCA2* mutation carriers individually or in combination. Our findings of no association

**Table 3** Genotype frequencies by mutant gene and breast cancer status with hazard ratio (HR) estimates

	Unaffected (%)	Affected (%)	HR	95% CI	P-value
<i>TP53 Arg72Pro (rs1042522)</i>					
<i>BRCA1/2</i>					
GG	1660 (54.4)	2164 (54.7)	1.00		
GC	1178 (38.6)	1508 (38.1)	1.00	0.92–1.10	
CC	214 (7.0)	287 (7.3)	1.01	0.85–1.20	
2-df test					0.99
Per allele			1.01	0.94–1.08	0.89
<i>BRCA1</i>					
GG	1127 (56.0)	1399 (55.2)	1.00		
GC	748 (37.2)	947 (37.4)	1.01	0.90–1.13	
CC	138 (6.9)	188 (7.4)	1.03	0.84–1.27	
2-df test					0.96
Per allele			1.01	0.93–1.10	0.77
<i>BRCA2</i>					
GG	533 (51.3)	765 (53.7)	1.00		
GC	430 (41.4)	561 (39.4)	0.98	0.84–1.14	
CC	76 (7.3)	99 (6.9)	0.99	0.72–1.36	
2-df test					0.95
Per allele			0.99	0.87–1.12	0.83
<i>MDM2 309T&gt;G (rs2279744)</i>					
<i>BRCA1/2</i>					
TT	358 (40.3)	530 (39.8)	1.00		
TG	405 (45.6)	615 (46.1)	0.99	0.84–1.18	
GG	126 (14.2)	188 (14.1)	0.93	0.73–1.17	
2-df test					0.79
Per allele			0.97	0.87–1.08	0.60
<i>BRCA1</i>					
TT	275 (39.7)	369 (39.5)	1.00		
TG	323 (46.6)	443 (47.4)	0.98	0.81–1.19	
GG	95 (13.7)	123 (13.2)	0.91	0.67–1.19	
2-df test					0.78
Per allele			0.96	0.84–1.09	0.54
<i>BRCA2</i>					
TT	83 (42.4)	161 (40.5)	1.00		
TG	82 (41.8)	172 (43.2)	1.07	0.77–1.50	
GG	31 (15.8)	65 (16.3)	0.93	0.60–1.44	
2-df test					0.83
Per allele			0.98	0.80–1.21	0.88

for these SNPs suggest that they have little or no effect on *BRCA*-related breast cancer risk. These results are consistent with the absence of risk association in the recent *TP53* haplotype analysis, involving *Arg72Pro* and an intronic polymorphism c.97-147ins16 bp, in a series of 2932 *BRCA1* and *BRCA2* carriers from CIMBA (Osorio *et al*, 2008). Our sample of mutation carriers had power of approximately 75% for *TP53* and 40% for *MDM2* to detect significant associations ( $P < 0.05$ ) for a per-allele HR of 1.1 and power of 100 and 90% respectively for a HR of 1.2, suggesting that we can reliably dismiss previously suggested associations (Martin *et al*, 2003; Osorio *et al*, 2006; Yarden *et al*, 2008).

Yarden *et al* showed that the *MDM2* GG genotype among Ashkenazi *BRCA1/2* mutations carriers was significantly associated with breast cancer diagnosed < age 51 ( $P = 0.019$ ) (Yarden *et al*, 2008). However, we did not find any evidence of an increased risk for the GG homozygotes among the 217 carriers of the *BRCA1* Ashkenazi mutations 185delAG and 5382insC (HR = 0.98, 95%CI 0.48–2.01) in this series.

The BCAC study of 5191 cases and 3834 controls found no evidence of an association of *TP53 Arg72Pro* and *MDM2 309T>G* either with breast cancer overall or with oestrogen receptor (ER) status of tumours (Schmidt *et al*, 2007). As the majority of *BRCA1*

mutation-associated breast tumours are ER-negative (Lakhani *et al*, 2005), the absence of an association in our study of breast cancer with the *TP53* and *MDM2* SNPs in *BRCA1* mutation carriers is consistent with the lack of an association with ER-negative cancers in the general population.

**ACKNOWLEDGEMENTS**

The CIMBA data management, DE, LM, SP, MC and EMBRACE are supported by Cancer Research UK Grants C1287/A10118 and C1287/A8874. CL is supported by Cancer Research UK Grant C8197/A10123. ACA is a Cancer Research – UK Senior Cancer Research Fellow.

Jacques Simard is Chairholder of the Canada Research Chair in Oncogenetics. This work was supported by the Canadian Institutes of Health Research for the ‘CIHR Team of Prediction and Communication of Familial Risks of Breast Cancer’ program. Daniel Sinnett holds the François-Karl Viau Chair in Pediatric Oncogenomics and is a scholar of the Fonds de la Recherche en Santé du Québec (FRSQ).

We thank Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study for their contributions to this resource, and the many families who contribute to kConFab. kConFab is supported by grants from the National Breast Cancer Foundation, the National Health and Medical Research Council (NHMRC) and by the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. The kConFab Clinical Follow Up Study has been funded by NHMRC grants 145684, 288704 and 454508. ABS and GCT are NHMRC fellows.

We gratefully acknowledge the contribution of Dr Jeffery P Struewing and Marbin Pineda for their laboratory support of this project. Drs Greene and Loud were supported by funding from the Intramural Research Program of the US National Cancer Institute, and from research contracts NO2-CP-11019-50 and N02-CP-65504 with Westat, Rockville, MD, USA.

The NICCC thanks the laboratory technicians Mrs. Irena Rimon and Mrs. Ana Gurtovnik for their technical support.

OCGN: We thank Mona Gill and Nayana Weerasooriya for assistance and we acknowledge funding from Cancer Care Ontario.

We thank Alicia Barroso, Concepción Hernandez and Anna Gonzalez for their technical support. The CNIO study was partially funded by the Asociación Española Contra el Cáncer (AECC), the Fundación Marató and the project PI081120 from the Ministry of Science and Innovation.

We thank Diana Torres and Muhammad U Rashid for providing DNA samples and supplying data. We thank Antje Seidel-Renkert for expert technical assistance. The DKFZ study was supported by the DKFZ.

The HEBON study is supported by the Dutch Cancer Society grants NKI1998-1854, NKI2004-3088, NKI 2007-3756.

The investigators at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust are supported by an NIHR grant to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. RE, EB and L D’M are also supported by Cancer Research UK Grant C5047/A8385. DGE and FL are supported the NIHR Biomedical Research Centre, Manchester.

We wish to thank all the GEMO collaborating members (Cancer Genetics Network ‘Groupe Génétique et Cancer’, Fédération Nationale des Centres de Lutte Contre le Cancer, France) for their contribution to this study. The GEMO study was supported by the Ligue Nationale Contre le Cancer and the Association ‘Le cancer du sein, parlons-en!’ Award.

We thank Juliane Koehler for her excellent technical assistance and the 12 centers of the GC-HBOC (German Consortium of Hereditary Breast and Ovarian Cancer) for providing samples and clinical data. GC-HBOC is supported by a grant of the German Cancer Aid (grant 107054) and the Center for Molecular Medicine Cologne (grant TV93) to Rita K Schmutzler.

We thank Drs Kirsimari Aaltonen, Carl Blomqvist and RN Hanna Jäntti for their help with the patient data and Dr Johanna Tommiska

for her kind help with the genetic analyses. The Finnish Cancer registry is gratefully acknowledged for the cancer data. The HEBCS study has been financially supported by the Helsinki University Central Hospital Research Fund, Academy of Finland (110663), Finnish Cancer Society and the Sigrid Juselius Foundation.

Supplementary Information accompanies the paper on British Journal of Cancer website (<http://www.nature.com/bjc>)

## REFERENCES

- Antoniou AC, Sinilnikova OM, Simard J, Leone M, Dumont M, Neuhausen SL, Struewing JP, Stoppa-Lyonnet D, Barjhoux L, Hughes DJ, Coupier I, Belotti M, Lassot C, Bonadona V, Bignon YJ, Rebbeck TR, Wagner T, Lynch HT, Domchek SM, Nathanson KL, Garber JE, Weitzel J, Narod SA, Tomlinson G, Olopade OI, Godwin A, Isaacs C, Jakubowska A, Lubinski J, Gronwald J, Gorski B, Byrski T, Huzarski T, Peock S, Cook M, Baynes C, Murray A, Rogers M, Daly PA, Dorkins H, Schmutzler RK, Vermsold B, Engel C, Meindl A, Arnold N, Niederacher D, Deissler H, Spurdle AB, Chen X, Waddell N, Cloonan N, Kirchoff T, Offit K, Friedman E, Kaufmann B, Laitman Y, Galore G, Rennert G, Lejbkowitz F, Raskin L, Andrulis IL, Ilyushik E, Ozcelik H, Devilee P, Vreeswijk MP, Greene MH, Prindiville SA, Osorio A, Benitez J, Zikan M, Szabo CI, Kilpivaara O, Nevanlinna H, Hamann U, Durocher F, Arason A, Couch FJ, Easton DF, Chenevix-Trench G (2007) RAD51 135G->C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies. *Am J Hum Genet* **81**: 1186–1200
- Antoniou AC, Spurdle AB, Sinilnikova OM, Healey S, Pooley KA, Schmutzler RK, Vermsold B, Engel C, Meindl A, Arnold N, Hofmann W, Sutter C, Niederacher D, Deissler H, Caldes T, Kampjarvi K, Nevanlinna H, Simard J, Beesley J, Chen X, Neuhausen SL, Rebbeck TR, Wagner T, Lynch HT, Isaacs C, Weitzel J, Ganz PA, Daly MB, Tomlinson G, Olopade OI, Blum JL, Couch FJ, Peterlongo P, Manoukian S, Barile M, Radice P, Szabo CI, Pereira LH, Greene MH, Rennert G, Lejbkowitz F, Barnett-Griness O, Andrulis IL, Ozcelik H, Gerdes AM, Caligo MA, Laitman Y, Kaufman B, Milgrom R, Friedman E, Domchek SM, Nathanson KL, Osorio A, Llorca G, Milne RL, Benitez J, Hamann U, Hogervorst FB, Manders P, Ligtenberg MJ, van den Ouweland AM, Peock S, Cook M, Platte R, Evans DG, Eeles R, Pichert G, Chu C, Eccles D, Davidson R, Douglas F, Godwin AK, Barjhoux L, Mazoyer S, Sobol H, Bourdon V, Eisinger F, Chompret A, Capoulade C, Bressac-de PB, Lenoir GM, Gauthier-Villars M, Houdayer C, Stoppa-Lyonnet D, Chenevix-Trench G, Easton DF (2008) Common breast cancer-predisposition alleles are associated with breast cancer risk in BRCA1 and BRCA2 mutation carriers. *Am J Hum Genet* **82**: 937–948
- Bond GL, Hu W, Bond EE, Robins H, Lutzker SG, Arva NC, Bargonetti J, Bartel F, Taubert H, Wuertl P, Onel K, Yip L, Hwang SJ, Strong LC, Lozano G, Levine AJ (2004) A single nucleotide polymorphism in the MDM2 promoter attenuates the p53 tumor suppressor pathway and accelerates tumor formation in humans. *Cell* **119**: 591–602
- Bougeard G, Baert-Desurmont S, Tournier I, Vasseur S, Martin C, Brugieres L, Chompret A, Bressac-de PB, Stoppa-Lyonnet D, Bonaiti-Pellie C, Frebourg T (2006) Impact of the MDM2 SNP309 and p53 Arg72Pro polymorphism on age of tumour onset in Li-Fraumeni syndrome. *J Med Genet* **43**: 531–533
- Candeias MM, Malbert-Colas L, Powell DJ, Daskalogianni C, Maslon MM, Naski N, Bourougaa K, Calvo F, Fahraeus R (2008) p53 mRNA controls p53 activity by managing Mdm2 functions. *Nat Cell Biol* **10**: 1098–1105
- Chenevix-Trench G, Milne RL, Antoniou AC, Couch FJ, Easton DF, Goldgar DE (2007) An international initiative to identify genetic modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: the Consortium of Investigators of Modifiers of BRCA1 and BRCA2 (CIMBA). *Breast Cancer Res* **9**: 104
- Copson ER, White HE, Blaydes JP, Robinson DO, Johnson PW, Eccles DM (2006) Influence of the MDM2 single nucleotide polymorphism SNP309 on tumour development in BRCA1 mutation carriers. *BMC Cancer* **6**: 80
- Deng CX (2006) BRCA1: cell cycle checkpoint, genetic instability, DNA damage response and cancer evolution. *Nucleic Acids Res* **34**: 1416–1426
- Dumont P, Leu JI, Della III PA, George DL, Murphy M (2003) The codon 72 polymorphic variants of p53 have markedly different apoptotic potential. *Nat Genet* **33**: 357–365
- Evers B, Jonkers J (2006) Mouse models of BRCA1 and BRCA2 deficiency: past lessons, current understanding and future prospects. *Oncogene* **25**: 5885–5897
- Greenblatt MS, Chappuis PO, Bond JP, Hamel N, Foulkes WD (2001) TP53 mutations in breast cancer associated with BRCA1 or BRCA2 germ-line mutations: distinctive spectrum and structural distribution. *Cancer Res* **61**: 4092–4097
- Lakhani SR, Reis-Filho JS, Fulford L, Penault-Llorca F, van der Vijver M, Parry S, Bishop T, Benitez J, Rivas C, Bignon YJ, Chang-Claude J, Hamann U, Cornelisse CJ, Devilee P, Beckmann MW, Nestle-Kramling C, Daly PA, Haites N, Varley J, Lalloo F, Evans G, Maugard C, Meijers-Heijboer H, Klijn JG, Olah E, Gusterson BA, Piloni S, Radice P, Scherneck S, Sobol H, Jacquemier J, Wagner T, Peto J, Stratton MR, McGuffog L, Easton DF (2005) Prediction of BRCA1 status in patients with breast cancer using estrogen receptor and basal phenotype. *Clin Cancer Res* **11**: 5175–5180
- Manie E, Vincent-Salomon A, Lehmann-Che J, Pierron G, Turpin E, Warcoin M, Gruel N, Lebigot I, Sastre-Garau X, Lidereau R, Remenieras A, Feunteun J, Delattre O, de TH, Stoppa-Lyonnet D, Stern MH (2009) High frequency of TP53 mutation in BRCA1 and sporadic basal-like carcinomas but not in BRCA1 luminal breast tumors. *Cancer Res* **69**: 663–671
- Martin AM, Kanetsky PA, Amirimani B, Colligon TA, Athanasiadis G, Shih HA, Gerrero MR, Calzone K, Rebbeck TR, Weber BL (2003) Germline TP53 mutations in breast cancer families with multiple primary cancers: is TP53 a modifier of BRCA1? *J Med Genet* **40**: e34
- Osorio A, Martinez-Delgado B, Pollan M, Cuadros M, Urioste M, Torrenteras C, Melchor L, Diez O, De La HM, Velasco E, Gonzalez-Sarmiento R, Caldes T, Alonso C, Benitez J (2006) A haplotype containing the p53 polymorphisms Ins16bp and Arg72Pro modifies cancer risk in BRCA2 mutation carriers. *Hum Mutat* **27**: 242–248
- Osorio A, Pollan M, Pita G, Schmutzler RK, Vermsold B, Engel C, Meindl A, Arnold N, Preisler-Adams S, Niederacher D, Hofmann W, Gadzicki D, Jakubowska A, Hamann U, Lubinski J, Toloczko-Grabarek A, Cybulski C, Debnik T, Llorca G, Yannoukakos D, Diez O, Peissel B, Peterlongo P, Radice P, Heikkinen T, Nevanlinna H, Mai PL, Loud JT, McGuffog L, Antoniou AC, Benitez J (2008) An evaluation of the polymorphisms Ins16bp and Arg72Pro in p53 as breast cancer risk modifiers in BRCA1 and BRCA2 mutation carriers. *Br J Cancer* **99**: 974–977
- Ruijs MW, Schmidt MK, Nevanlinna H, Tommiska J, Aittomaki K, Pruntel R, Verhoef S, Van't Veer LJ (2007) The single-nucleotide polymorphism 309 in the MDM2 gene contributes to the Li-Fraumeni syndrome and related phenotypes. *Eur J Hum Genet* **15**: 110–114
- Schmidt MK, Reincke S, Broeks A, Braaf LM, Hogervorst FB, Tollenaar RA, Johnson N, Fletcher O, Peto J, Tommiska J, Blomqvist C, Nevanlinna H, Healey CS, Dunning AM, Pharoah PD, Easton DF, Dork T, Van't Veer LJ (2007) Do MDM2 SNP309 and TP53 R72P interact in breast cancer susceptibility? A large pooled series from the breast cancer association consortium. *Cancer Res* **67**: 9584–9590
- Toledo F, Wahl GM (2006) Regulating the p53 pathway: *in vitro* hypotheses, *in vivo* veritas. *Nat Rev Cancer* **6**: 909–923
- Tommiska J, Eerola H, Heinonen M, Salonen L, Kaare M, Tallila J, Ristimaki A, von SK, Aittomaki K, Heikkila P, Blomqvist C, Nevanlinna H (2005) Breast cancer patients with p53 Pro72 homozygous genotype have a poorer survival. *Clin Cancer Res* **11**: 5098–5103
- Wasielewski M, Nagel JH, Brekelmans C, Klijn JG, van den OA, Meijers-Heijboer H, Schutte M (2007) MDM2 SNP309 accelerates familial breast carcinogenesis independently of estrogen signaling. *Breast Cancer Res Treat* **104**: 153–157
- Yarden RI, Friedman E, Metsuyanin S, Olender T, Ben-Asher E, Papa MZ (2008) MDM2 SNP309 accelerates breast and ovarian carcinogenesis in BRCA1 and BRCA2 carriers of Jewish-Ashkenazi descent. *Breast Cancer Res Treat* **111**: 497–504