Fluorouracil: Biochemistry and Pharmacology

By Herbert M. Pinedo and Gode Fridus J. Peters

Fluorouracil (5FU) is still considered the most active antineoplastic agent in the treatment of advanced colorectal cancer. The drug needs to be converted to the nucleotide level in order to exert its effect. It can be incorporated into RNA leading to interference with the maturation of nuclear RNA. However, its conversion to 5-fluoro-2'-deoxy-5' monophosphate (FdUMP) leading to inhibition of thymidylate synthase (TS) and subsequently of DNA synthesis, is considered to be its main mechanism of action. In the presence of a folate cofactor a covalent ternary complex is formed, the stability of which is the main determinant of the action of 5FU. Resistance against 5FU can be mainly attributed to aberrations in its metabolism or to alterations of TS, eg, gene amplification, altered kinetics in respect to nucleotides or folates. Biochemical modulation of 5FU metabolism can be applied to overcome resistance against 5FU. A variety of normal purines, pyrimidines, and other antimetabolites have

been studied in this respect, but only some of them have been clinically successful. Delayed administration of uridine has recently been shown to "rescue" mice and patients from toxicity, while pretreatment with leucovorin is the most promising combination to enhance the therapeutic efficacy. 5FU is frequently administered in an intravenous (IV) injection, and shows a rapid distribution and a triphasic elimination. The nonlinearity of 5FU pharmacokinetics is related to saturation of its degradation. Continuous infusion of 5FU led to different kinetics. Regional administration, such as hepatic artery infusion, offers a way to achieve higher drug concentrations in liver metastases and is accompanied by lower systemic concentration. The current status of the biochemical and pharmacokinetic data is reviewed.

J Clin Oncol 6:1653-1664. © 1988 by American Society of Clinical Oncology.

FLUOROURACIL (5FU) has been used for several decades for the treatment of various types of cancer.^{1,2} The mechanism underlying the action of this compound is complex and depends on the type of tissue, ie, whether normal or tumor tissue is involved. In attempts to increase the antitumor activity and limit toxicity, various analogues of 5FU such as Ftorafur and 5'-deoxy-5-fluorouridine (5'dFUR) have been studied clinically³ and combinations of 5FU with other antimetabolites or natural compounds have been evaluated in efforts to improve the therapeutic index.⁴⁻¹² The present review deals with the biochemical activation and the inactivation of 5FU, various mechanisms underlying its action, resistance to and biochemical modulation of 5FU, metabolic aspects, and some new data on the pharmacology of the drug.

BIOCHEMICAL ACTIVATION

The initial metabolism of 5FU to nucleotides such as fluorouridine 5'-triphosphate (FUTP) and 5-fluoro-2'-deoxyuridine-5'-monophosphate (FdUMP) is essential for its action.^{1,3,4,13} Several enzymes belonging to pyrimidine metabolism are required for the conversion of 5FU to nucleotides (Fig 1). FdUMP can be formed from FUMP

via reduction of FUDP. The extent of growth inhibition by 5FU may be correlated with the activity of one or more of the enzymes catalyzing the initial metabolism of 5FU.³ For some cell lines orotate phosphoribosyl-transferase (OPRT) has been shown to play a major role in the initial metabolism, whereas for other cells uridine phosphorylase is more important.¹⁴ Nucleotides formed via the direct pathway (via OPRT) and the indirect pathway (via FUR) are incorporated into different RNA fractions.¹⁵ Sensitivity of cell lines and tumors to 5FU might also depend on the availability of cosubstrates required for the con-

---

From the Department of Oncology, Free University Hospital, Amsterdam.

Submitted September 30, 1987; accepted June 1, 1988.

Supported by the Netherlands Cancer Foundation "Koningin Wilhelmina Fonds," Grants No. AUKC VU 80-3, IKA VU 83-16, and 88-20, and by gifts from Hoffman-LaRoche, Mijdrecht, The Netherlands and Cyanamid-Lederle, Etten-Leur, The Netherlands. Dr Peters is a senior research fellow of the Royal Netherlands Academy of Sciences (KNAW).

Address reprint requests to G.J. Peters, PhD, Department of Oncology, Free University Hospital, PO Box 7057, 1007 MB Amsterdam, The Netherlands.

© 1988 by American Society of Clinical Oncology.

0732-183X/88/0610-0010\$3.00/0

Fig 1. Metabolism of 5FU. The enzymes involved are: **1, Orotate phosphoribosyl-transferase (OPRT); 2, uridine phosphorylase; 3, thymidine phosphorylase; 4, uridine kinase; 5, thymidine kinase; 6, ribonucleotide reductase; 7, thymidylate synthase (TMPsyn or TS); 5-fluorouridine, FUR; 5-fluoro-2'-deoxyuridine, FdUR; phosphoribosylpyrophosphate, PRPP; 5'-fluorouridine-5' monophosphate, FUMP; 5'-fluorouridine-5'-diphosphate FUDP; 5'-fluoro-2'-deoxyuridine-5'-fluoro-2'-deoxyuridine-5' triphosphate, FdUTP.**

version of 5FU to active nucleotides,¹³ as discussed in the section on modulation.

5FU INACTIVATION

5FU can be inactivated by degradation to 5-fluorodihydrouracil (F-DHU) (Fig 2). Further degradation of 5FU has been studied as far as

Catabolism of 5-Fluorouracil (FU)

Fig 2. Degradation of 5FU dihydrouracil, DHU; 5-fluoro-ureido propionate, FUPA.

5-fluoro-alanine (F-BAL), and further metabolism of F-BAL is not unlikely. β-alanine itself is a substrate for carnosine, but it can also be converted to acetate. Similarly, F-BAL can be converted to fluoroacetate, which has been related to neurotoxicity.¹⁶ 5FU degradation occurs in all tissues, but tumor tissue contains very small amounts of dihydrouracil dehydrogenase.¹⁷ The activity of this enzyme, while occurring in the kidney, is most intense in the liver,¹⁸ which means that the liver plays an important role in 5FU degradation and elimination.

It has been shown in patients that large amounts of 5FU are degraded to F-DHU,¹⁹ and fluorine-19 (19F) nuclear magnetic resonance (NMR) has been used to show that, in vivo, F-DHU is rapidly degraded further to F-BAL.²⁰ Recently, a new catabolite of 5FU was detected in bile, and was identified as an N-cholyl-2-fluoro-β-alanine conjugate by NMR²¹ and enzymatic methods.²² In vivo inhibition of 5FU degradation has been thought to increase the availability of 5FU to tumors. However, it has been reported that an improved therapeutic index was not observed clinically after the administration of thymidine and 5FU,²³ but in rats toxicity appeared to be increased.²⁴ Impaired 5FU degradation due to a deficiency of dihydrouracil dehydrogenase led to a dramatic and fatal increase of 5FU toxicity.²⁵

It may be concluded that inhibition of 5FU degradation probably will not improve therapeutic efficacy, since toxicity increases as much as or even more than the antitumor activity.

INHIBITION OF THYMIDYLATE SYNTHASE BY FdUMP

It has long been recognized²⁶⁻²⁸ that inhibition of TS by FdUMP is one of the main mechanisms underlying 5-FU action (Fig 3). Initial studies have been performed with the methotrexate (MTX)-resistant mutant from *Lactobacillus casei*.²⁶ Since the older studies have been reviewed extensively by Danenberg²⁷ and Danenberg and Lockshin, this review will be limited to clinically relevant aspects. Enzyme kinetic studies in various systems revealed a rather low Michaelis-Menten constant (Km) of about 2 μmol/L for dUMP.²⁷⁻²⁹ Without preincubation the inhibition constant (Ki) for FdUMP appeared to be competitive,²⁷ with a Km/Ki ratio of about 1,000.

Fig 3. Inhibition of TS by FdUMP (hatched bar) leading to accumulation of dUMP and FdUMP, depletion of TMP and TTP, and inhibition of DNA synthesis. TS catalyzes the conversion of dUMP to TMP. 5-CHO-THF, 5-formyl-tetrahydrofolate (leucovorin); DHF, dihydrofolate; 5,10-CH₂THF, 5,10-methylene-tetra hydrofolate.

FdUMP forms a reversible but tight-binding covalent bond with TS in the presence of 5,10-CH₂ tetrahydrofolate (THF),²⁷⁻³⁰ which is precipitable with trichloroacetic acid.³⁰ The *in vivo* recovery of uninhibited TS differed among various tumors and cell lines.³¹⁻³⁴ Retention of TS inhibition was also dependent on the ratio between free dUMP and FdUMP levels.³³ The presence of folate cofactor appeared to be correlated with the extent of enzyme inhibition and the retention of the complex.³³⁻³⁶ Not only enzyme inhibition, but also the enzyme level before treatment was related to growth inhibition by 5FU.³³ The degree of inhibition of TS and the persistence of inhibition are essential factors for maximal *in vivo* growth inhibition by either 5FU or FUdR. A relationship has been reported to exist between low sensitivity to 5FU and rapid disappearance of FdUMP.^{37,38} Retention of the inhibition of TS is dependent on the binding of FdUMP and the stabilization of the ternary complex by 5,10 CH₂ THF^{33,38-40} or one of its polyglutamates.^{33,41} The naturally occurring cofactor for TS is a polyglutamate.^{42,43} The *K_i* for FdUMP in the presence of folylpolyglutamate is lower than with the monoglutamate.⁴⁴ A noncovalent complex of FdUMP with TS that is less stable can also be formed.^{27,28} Folates appear to be essential for the formation of a covalent complex. Evidence has been presented that tumors of patients responding to 5FU show greater inhibition of TS than tumors of patients with progressive disease.⁴⁵

5FU INCORPORATION INTO RNA

In most cells and tissues 5FU will be converted not only to FdUMP, but also to FUTP which can

be incorporated into all classes of RNA in tumor cells, mainly into nuclear RNA.⁴⁶ Processing of nuclear RNA into cytoplasmic rRNA is in all probability the essential factor leading to cytotoxicity. Initially, it was demonstrated *in vitro* that the amount of 5FU incorporated into RNA correlated with the sensitivity to 5FU of various cell lines⁴⁷⁻⁵⁰ and *in vivo* with the antitumor effect of 5FU⁵¹ and with gastrointestinal cytotoxicity.⁵² Although 5FU is incorporated into most species of RNA, toxicity was not correlated with incorporation in all of these species.^{53,54} The cytotoxicity due to incorporation of 5FU into RNA is mainly determined by the incorporation of 5FU into nuclear RNA.^{51,55-57} Recently, more evidence has been presented that misincorporation into RNA might be associated with a block in processing and/or nuclear cytoplasmic transport.⁵⁸⁻⁶¹ Thus, in all likelihood, 5FU incorporation into RNA produces cytotoxicity by interference with the maturation of nuclear RNA.

5FU INCORPORATION INTO DNA

5FU incorporation into DNA has long been considered a very unlikely event, and not contributory to 5FU cytotoxicity. Intracellular FdUTP is hydrolyzed by dUTPase⁶²; FdUTP incorporated into DNA is thought to be removed by uracil-DNA glycosylase.⁶³ Despite these protective mechanisms, some 5FU residues can be incorporated into DNA.⁶³⁻⁷¹ 5FU incorporation has been shown to be enhanced by MTX,⁶⁹ but increased excision of 5FU residues⁶³ has also been reported. A relation between 5FU incorporation and cytotoxicity has been postulated.^{63,72} 5FU is capable of inducing DNA strand breaks,^{67,73} which might also be related to inefficient DNA repair of normally occurring defects.⁷³ Thus, the extent to which 5FU incorporation into DNA, the subsequent excision, its effect on DNA repair, and the induction of strand breaks are related to 5FU cytotoxicity is not yet clear.

5FU NUCLEOTIDE SUGARS

Uridine metabolites occur predominantly as nucleotide sugars, such as uridine 5'-diphosphate (UDP)-glucose and UDP-N-acetyl-hexoseamines. These sugars are substrates for glycosyltransferases, which catalyze the glycosylation of proteins and lipids, and are important for cellular functions involving, for example,

cell-surface glycoprotein and glycolipid receptors, differentiation markers, and recognition determinants. It has been shown that 5FU can be a substrate for the synthesis of FUDP sugars, such as FUDP-hexoses,⁷⁴⁻⁷⁶ FUDP-hexoseamines, and FdUDP-N-acetylhexoseamines.⁷⁷⁻⁷⁹ From these findings it is clear that FUDP sugars are formed, but their effect on the nucleotide sugar metabolism and glycosylation warrants further investigation.

RESISTANCE TO 5FU

5FU resistance, whether intrinsic or acquired, is usually caused by aberrations in the metabolism of 5FU or altered effects of 5FU-metabolites. The normal metabolism has been discussed in the preceding sections, and aberrations are summarized in Table 1. Generally, studies on 5FU resistance have been performed by comparison of several tumor cell lines with different sensitivity to 5FU or selection of a 5FU-resistant subpopulation from a sensitive tumor or cell line. It has already been shown by Reyes and Hall⁸⁰ and Kessel et al⁸¹ that tumors with a low level of anabolism have a low sensitivity to 5FU. OPRT may be the limiting enzyme for 5FU anabolism.⁸⁰ 5FU transport across the cell membrane might not limit its activity, but FUDR resistance was found to be related to deficiency in its transport.⁸² Depletion of cosubstrates, ie, (deoxy)Ribose-1-P or PRPP,⁴ seems to limit the anabolism of 5FU, as suggested by indirect evidence. Increased availability of Ribose-1-P,¹⁵ deoxy-Rib-1-P,^{39,83} or PRPP⁴ enhanced the sensitivity to 5FU. Enhanced nucleotide catabolism due to a high level

of alkaline phosphatase activity has been shown to affect FUDR toxicity.⁸⁴ Altered dTTP levels also affect FUDR toxicity.⁸⁵

Aberrations in TS kinetics can lead to resistance against 5FU. Several forms of aberration are summarized in Table 1. Altered enzyme kinetics²⁹ for TS were reflected by a higher dissociation constant for the ternary covalent complex, but also by a weaker binding of dUMP. Intrinsic resistance to 5FU has been associated with high accumulation of dUMP.³⁸ The turnover of TS was higher in a resistant sub-cell line than in the sensitive cells.²⁹ A higher activity of TS was found in an FUDR-resistant sub-cell line,³⁶ possibly due to gene amplification. Amplification of the gene coding for TS has been shown,⁸⁶ recently also in a case of human colon cancer with acquired resistance.⁸⁷ The stability of the ternary complex depends on the concentrations of dUMP and FdUMP and the kinetic parameters, but also on the availability of folates. Low total folate pools were associated with 5FU resistance,^{35,41} as well as a low proportion of polyglutamate derivatives.⁴¹ It may be concluded that resistance to 5FU can be due to a variety of aberrations in 5FU-metabolism, but factors affecting TS appear to be of major clinical relevance.

BIOCHEMICAL MODULATION OF 5FU

Biochemical modulation of anticancer agents involves the pharmacologic manipulation of the intracellular pathways of a drug. The aim is to improve the therapeutic index.^{10,88} Modulation can be used to overcome 5FU resistance. For cell culture systems and animal models, various combinations of 5FU with other drugs have been selected on a rational basis.^{3,88} A list of such combinations is given in Table 2, where a subdivision according to antimetabolites and naturally occurring purines and pyrimidines is made. The list is not exhaustive. Although certain combinations have only theoretical value, their use in *in vitro* studies has led to a better insight into the mechanism of action and resistance. Other combinations have more practical applications. The preclinical studies on combinations of 5FU with purines have thrown more light on 5FU activation and metabolism, but the findings have not yet been applied clinically. Purines can even provide protection against 5FU cytotoxicity by

Table 1. Resistance to 5FU

Deficiency of 5FU anabolism
Deficiency of 5FU transport
Depletion of essential cosubstrates
Enhanced catabolism of 5FU, FUMP, or FdUMP
Enhanced intracellular uridine concentrations
Altered dTTP levels
Alterations in thymidylate synthase
Altered enzyme kinetics
Enhanced dUMP accumulation
Decreased FdUMP retention
Rapid recovery of new enzyme synthesis
Gene amplification
Decreased stability of ternary complex
Depletion of folates
Decreased polyglutamylation of folates

Abbreviation: dTTP, deoxythymidine triphosphate.

Table 2. Biochemical Modulation of 5FU

Modulating Agent	Postulated Mechanism	Sequence of Addition	Tested Systems
Purines ^{13,83,89-93}			
Inosine	Increased rib-I-P might lead to increased anabolism	Pre, sim	In vitro, mice
Guanosine	Similar to inosine	Pre, sim	In vitro, mice
GMP	Similar to guanosine after conversion of GMP to guanosine	Pre, sim	In vitro, mice
Deoxyinosine	Increased dRib-I-P leads to enhanced FdUMP	Pre, sim	In vitro, mice
Pyrimidines			
Thymidine ^{7,23}	Inhibition of 5FU breakdown	Pre, sim	Mice, patients
Uridine ^{6,9,12,94-97,99,101}	Rescue of normal tissue by competition of UTP with FUTP	Delayed	Mice, patients, in vitro
Cytidine ^{12,96,99,101}	Similar to uridine	Delayed	Mice, in vitro
Antimetabolites			
Methotrexate ^{4,88}	Increased 5FU anabolism due to enhanced PRPP	Pre	In vitro, animals, patients
PALA ^{6,8,78,102}	Decrease of uracil nucleotides leads to enhanced anabolism	Pre	In vitro, animals, patients
Allopurinol ¹⁰³	Postulated differential metabolism of 5FU in tumors and normal tissues is used, decreased toxicity	Sim	In vitro, animals, patients
Hydroxyurea ¹⁰⁵	Inhibition of ribonucleotide reduction prevents rescue by normal nucleotides	Post	In vitro
Dipyridamol ¹⁰⁶	Prevention efflux of FUR and FdUR	Sim	In vitro
Leucovorin ^{5,33,35,38-41,107-113}	Enhanced retention of FdUMP binding to thymidylate synthase in tumors	Pre, sim	In vitro, animals, patients

Abbreviations: pre, pretreatment; sim, simultaneous; post, posttreatment; rib-I-P, ribose-I-P; dRib-I-P, deoxyribose-I-P; GMP, guanosine-5'-phosphate.

depletion of PRPP leading to inhibition of activation.^{89,90}

Combinations of 5FU with pyrimidines have been studied in both animal models and patients. 5FU combined with thymidine led to increased toxicity^{7,23} caused by enhancement of the anabolism of 5FU in normal tissues. An interesting scientifically based combination is that of 5FU and uridine,^{8,9} chosen on the hypothesis that 5FU antitumor activity consists mainly of inhibition of TS whereas 5FU toxicity is caused by incorporation of 5FU into RNA^{52,94} (Fig 4); if high-dose uridine is administered several hours after 5FU, the binding of FdUMP to TS will not

be affected but UTP will replace FUTP in RNA.^{8,9} Preliminary results of a phase I study^{11,12,95,96} indicated that delayed uridine administration prevented myelosuppression induced by 5FU.⁹⁵ Since cytidine, too, can prevent 5FU toxicity in mice,¹⁰ its clinical application should be considered.

The selectivity of uridine "rescue" might be related not only to a different mechanism of action in tumors and normal tissues, but also to the metabolism of uridine. Darnowsky and Handschumacher^{97,98} showed that uridine concentrations in murine tissues are much higher than in plasma, and suggested a concentrative mecha-

Fig 4. Proposed mechanism of uridine (UR) "rescue." FdUMP bound to TS will not be affected by delayed UR, while FUTP formed from UR will interact with FUTP for incorporation into RNA.

nism for uridine uptake by tissues and especially by tumors.⁹⁸ This might be related to a selective effect of delayed uridine administration on 5FU-induced myeloid toxicity. The relative increase in plasma uridine is much higher than that occurring in tumor tissue.^{11,12,99} If the concentrations are similar in plasma and bone marrow, uridine might have a stronger effect on bone marrow than on other tissues since it is an important precursor for pyrimidine nucleotide synthesis in lymphoid cells.¹⁰⁰ Uridine treatment was complicated by the development of fever in both humans and the rabbit,^{11,96} whereas mice developed hypothermia.⁹⁹ Fever in humans could be prevented by a complicated schedule of intermittent administration.^{11,96}

Although experimental data on MTX plus 5FU⁴ have encouraged clinical studies, the results of treatment have been disappointing. There are some data indicating that a longer interval between MTX and 5FU may increase the response rate in humans.^{10,88} The combination of N-phosphonacetyl-L-aspartate (PALA) and 5FU has given no clinical benefit.^{6,102}

5FU and allopurinol were combined because metabolites of allopurinol were expected to inhibit 5FU anabolism catalyzed by OPRT in normal tissues but not in tumor tissues.¹⁰³ It was postulated that uridine phosphorylase would be important for 5FU anabolism in tumor tissue.¹⁰⁴

This combination, which has not yet shown any clinical advantage, may require further evaluation when more is known about 5FU metabolism in human tumors and healthy tissues.

An interesting combination is that of 5FU and hydroxyurea,¹⁰⁵ which exploits the cell-phase specificity of 5FU at low concentrations.⁴⁸ The combination of 5FU and dipyridamole¹⁰⁶ is based on the nucleoside transport-inhibiting properties of dipyridamole, which does not affect 5FU uptake but inhibits efflux of FUR and FUDR. Neither of the last two combinations has received sufficient clinical investigation.

Selective rescue of healthy tissue and increased FdUMP binding to TS in tumors offer a favorable basis for combination with 5-FU. Foliates are known to prolong retention of the FdUMP-TS complex.^{33,38-41,107,108} Leucovorin has been used for this purpose. In two murine colon cancer lines pretreatment with leucovorin increased the therapeutic effect of 5FU.¹⁰⁹ Phase I and II clinical trials of 5FU-leucovorin in patients with advanced colorectal cancer showed response rates of up to 40%, which is considerably higher than those obtained with 5FU alone.^{5,110,111} The response rate in randomized trials comparing 5FU with 5FU plus leucovorin in colorectal cancer lay between 40% and 48% for the combination and between 10% and 15% for single-agent 5FU.^{112,113} Combination of 5FU and leucovorin with delayed uridine may also enhance antitumor activity and prevent toxicity.¹⁰⁹

Of the combinations with 5FU tested so far, that of leucovorin with 5FU is the most promising. Since a number of different schedules are in use results might be improved by using more appropriate schedules. Both preclinical¹⁰⁹ and clinical¹¹⁰⁻¹¹³ data justify the conclusion that pretreatment with leucovorin looks promising. The clinical schedule of a two-hour leucovorin infusion with a mid-infusion bolus injection of 5FU appears to give the best results, and this is in agreement with the preclinical data.

PHARMACOKINETICS

The pharmacokinetics of 5FU have been reviewed extensively by others.^{114,115} Initial assays of 5FU lacked either specificity or sensitivity.¹¹⁵ Currently, the most widely used method is high performance liquid chromatography (HPLC) combined with UV absorption with a detection

limit of 0.5 to 1.0 $\mu\text{mol/L}$ 5FU.^{12,115,116} A lower detection limit for 5FU (down to 3×10^{-9} mol/L; 0.3 ng/mL) can be achieved with gas-chromatography-mass-spectrometry (GC-MS).^{117,118} Most of the pharmacokinetic studies were restricted to three hours or less, but with the sensitive GC-MS method 5FU plasma concentrations can be followed for at least eight hours after the injection,¹¹⁸ which makes it possible to perform long-term pharmacokinetic studies.¹¹⁹

The pharmacokinetics of single-dose 5FU administered as an intravenous (IV) bolus injection in doses ranging between 300 and 600 mg/m² have been studied in detail,^{114,120} and the findings are summarized in Table 3. Rapid distribution over a large volume and rapid elimination have been reported to follow peak levels lying in the millimolar range (Fig 5). The total clearance was rather high (Table 3), and comparable to the liver flow, but hepatic extraction has been estimated to be 50%.¹²³ Yet the liver is the organ with the highest level of dihydrouracil dehydrogenase activity.¹⁸ The kidneys, in which the activity of this enzyme is also high, contribute to elimination by both degradation and active renal excretion, about 20% of 5FU being excreted as the parent drug.¹¹⁵ The lungs have also been reported to be a major site of 5FU clearance.^{114,115,124,125} Collins et al¹²⁰ have shown that a saturable two-compartment model can be used to describe the elimination kinetics of 5FU. Calculation gave an apparent K_m of 15 $\mu\text{mol/L}$ in plasma.

Nonlinearity of 5FU kinetics has been described by several authors,^{114,115,122,124,125} and is related to the saturation of 5FU catabolism. Studies on the pharmacokinetics of 5FU catabolites have been hampered by the lack of appropriate detection methods. A relatively insensitive method applied HPLC (Fig 5), and a more sensitive

Fig 5. Representative curves for 5FU and F-DHU post-IV bolus injection of 5FU (500 mg/m²) to a colorectal cancer patient. 5FU was determined with GC-MS.¹¹⁸ F-DHU was determined with HPLC with UV detection at 210 nm (Van Groeningen et al¹²¹); due to F-DHU instability in plasma^{19,122} extractions and measurements were performed as soon as possible.

method used GC with electron capture detection.¹⁹ With ¹⁹F-NMR, the other catabolites could also be demonstrated in human plasma.²⁰ Analysis of the cumulative urinary excretion of these catabolites showed that F-BAL was the major one followed by FUPA. F-DHU was a minor constituent of the urinary excretion products.¹²⁶ The high detection limit of 10 $\mu\text{mol/L}$, which is also in the range of the peak plasma concentration,¹²⁶ is the major limitation of this technique. Improvement might permit investigation of the dose-dependency of 5FU pharmacokinetics in humans in relation to the in vivo behavior of F-DHU.

There is no evidence that continuous IV administration of 5FU is associated with a higher antitumor efficacy than bolus administration. These two schedules give quite different types of toxicity, mucositis being dose-limiting for infusion and myelosuppression for the bolus injection.¹¹⁴ The pharmacokinetics of continuous 5FU infusion differ significantly from those of the IV bolus, the former having a much higher clearance value of 2 to 6 L/min,¹²⁰ which considerably exceeds the hepatic flow of 1.5 L/min and approaches the cardiac output. This high clearance level can be explained mainly by the high pulmonary extraction.^{114,115,120,124} Pulmonary extraction accounted for a clearance higher than the cardiac output.¹¹⁵ However, it has been shown that the liver and kidneys also contribute to clearance.

In the past, 5FU was administered orally; however, there is a marked variability in its bioavailability, ranging between 28% and

Table 3. Pharmacokinetic Parameters of 5FU Administered as an IV Bolus Injection

Parameters	Value
Peak levels	$10^{-4} - 10^{-3}$ mol/L
$T_{1/2}$, β	10-20 min
MRT	12-23 min
V_D	8-54 L
Total clearance	0.5-2.0 L/min

Note. Values are from doses varying between 300 and 600 mg/m².^{114,115,122,124,125}

Abbreviations: V_D , volume of distribution; MRT, mean residence time.

100%.^{113,114,127} This finding may be related to a saturable hepatic metabolism induced by dihydrouracil dehydrogenase¹²⁰ (Fig 2), but also to an additional first-pass effect arising from the rather high mucosal activity of dihydrouracil dehydrogenase.¹⁷ Because of the substantial variability observed, it is generally accepted that 5FU should not be administered orally.

5FU is also administered intrahepatically by portal or arterial infusion for the treatment of liver metastases. Specifically for this route of administration, hepatic extraction and the rate of infusion determine the systemic availability. The use of rapid intrahepatic arterial infusions at a high dose (1,000 mg/m²/d) gave relatively low hepatic extraction amounting from 20% to 60%,^{123,128} which led to a high systemic availability. With a slower infusion rate and/or lower doses (780 mg/m²/d), hepatic extraction exceeded 90%^{123,128} and this was accompanied by low systemic toxicity. More evidence pointing to hepatic saturation was provided by the observation that 5FU levels rose significantly during the infusion.¹²⁸ This new pharmacokinetics information makes it possible to design better 5FU schedules for the treatment of liver metastases.

Intraperitoneal infusions offer the possibility of achieving higher drug concentrations, give optimal exposure of tumor tissue within the abdominal cavity, and provide more effective treatment of not only the liver (via the portal vein) but also of peritoneal metastases. 5FU can be administered by intraperitoneal peritoneal dialysis¹²⁹ or via implantable devices.¹³⁰ Intraperitoneal 5FU was cleared at a rate of 14 mL/min, and 82% of the 5FU administered was absorbed within four hours. Hepatic extraction was calculated to be 67%.¹²⁹ A 2- to 3-log difference was observed between peritoneal and plasma 5FU concentrations.^{129,130} With continuous infusion the mean

steady-state level of 5FU in the intraperitoneal cavity was 622 μ mol/L.¹³⁰ Total body clearance ranged from 0.9 to 16.5 L/min,^{129,130} which is similar to the rate seen with continuous IV infusion of 5FU. Clearance decreased with increasing 5FU concentration, which is consistent with saturable or nonlinear 5FU pharmacokinetics. It might be worthwhile to study this method of administration in an adjuvant setting after surgical removal of Dukes B2 and C colorectal cancers.

CONCLUSIONS

Studies on the biochemistry of 5FU have yielded new detailed information on the mechanism of action of 5FU and about resistance to 5FU. Although valuable, most of this information was obtained from studies performed in vitro or in animals. Detailed biochemical studies in humans have been undertaken but are still scarce. Pharmacokinetic studies have supplied a basis for the application of new administration schedules, but improvement of the antitumor effect has not been achieved yet. Promising clinical results have been reported for treatment with 5FU plus leucovorin. Biochemical modulation seems to be the approach most likely to improve the therapeutic efficacy of 5FU; further research in this area is urgently needed.

More information about the intratumoral metabolism of 5FU is needed as well as in normal tissues. Analysis of the pharmacodynamic behavior of 5FU metabolites, such as FdUMP, and their binding to TS is essential, as is a more detailed analysis of the mechanisms leading to resistance and toxicity in humans. This information must be obtained before the selectivity of 5FU can be improved.

ACKNOWLEDGMENT

We thank C.J. van Groenigen for critical reading of the manuscript and valuable discussions.

REFERENCES

1. Myers CE, Diasio R, Eliot HM, et al: Pharmacokinetics of the fluoropyrimidines: Implications for their clinical use. *Cancer Treat Rev* 3:175-183, 1976
2. Gustavsson B, Hafström L: Adjuvant and palliative treatment of colorectal cancer with fluorinated pyrimidines. A pharmacologic and clinical review. *Acta Chir Scand* 504:1-28, 1981 (suppl)
3. Chabner BA: Pyrimidine Antagonists, in Chabner BA (ed): *Pharmacologic Principles of Cancer Treatment*. Philadelphia, Saunders, 1982, pp 183-212
4. Mini E, Bertino JR: Sequential methotrexate and 5-fluorouracil: Biochemical pharmacology and therapeutic use. *Chemotherapy* 11:147-162, 1983
5. Machover D, Schwarzenberg L, Goldschmidt E, et al: Treatment of advanced colorectal and gastric adenocarcinomas with 5-FU combined with high dose folinic acid: A pilot study. *Cancer Treat Rep* 66:1803-1807, 1982
6. Martin DS, Stolfi RL, Sawyer RC, et al: Application of biochemical modulation with therapeutically inactive modulating agent in clinical trials of cancer chemotherapy. *Cancer Treat Rep* 69:421-423, 1985
7. Young CW, Woodcock TM, Martin DS: Modulation of

5FU action by thymidine in murine and human tumors. *Cancer Treat Rep* 65:83-87, 1981 (suppl 3)

8. Martin DS, Stolfi RL, Sawyer RC, et al: High-dose 5-fluorouracil with delayed uridine "rescue" in mice. *Cancer Res* 42:3964-3970, 1982

9. Klubes P, Cerna I, Meldon MA: Uridine rescue from the lethal toxicity of 5-fluorouracil in mice. *Cancer Chemother Pharmacol* 8:17-21, 1982

10. Martin DS: Biochemical modulation: Perspectives and objectives, in Harrap KR, Connors TA (eds): *Proceedings of the Eighth Bristol-Myers Symposium on Cancer Research: New Avenues in Developmental Cancer Chemotherapy*. London, Academic, 1987, pp 113-162

11. Van Groeningen CJ, Leyva A, Kraal I, et al: Clinical and pharmacokinetic study of prolonged administration of high-dose uridine intended for rescue from 5-fluorouracil toxicity. *Cancer Treat Rep* 70:745-750, 1986

12. Leyva A, Van Groeningen CJ, Kraal I, et al: Phase I and pharmacokinetic studies of high-dose uridine intended for rescue from 5-fluorouracil toxicity. *Cancer Res* 44:5928-5933, 1984

13. Valeriote F, Santelli G: 5-Fluorouracil. *Pharmacol Ther* 24:107-132, 1984

14. Peters GJ, Laurensse E, Leyva A, et al: Sensitivity of human, murine, and rat cells to 5-fluorouracil and 5'-deoxy-5-fluorouridine in relation to drug-metabolizing enzymes. *Cancer Res* 46:20-28, 1986

15. Shani J, Danenberg PV: Evidence that intracellular synthesis of 5-fluorouridine-5'-phosphate from 5-fluorouracil and 5-fluorouridine is compartmentalized. *Biochem Biophys Res Commun* 122:439-445, 1984

16. Koenig H, Patel A: Biochemical basis for fluorouracil neurotoxicity. *Arch Neurol* 23:155-160, 1970

17. Ho DH, Townsend L, Luna MA, et al: Distribution and inhibition of dihydrouracil dehydrogenase activities in human tissues using 5-fluorouracil as a substrate. *Anticancer Res* 6:781-784, 1986

18. Yngner T, Engelbrecht C, Lewan C, et al: Anabolism versus catabolism of [5-³H] uridine and its relationship to ribonucleic acid labeling in mouse liver after partial hepatectomy. *Biochem J* 178:1-8, 1979

19. De Bruijn EA, Driessen O, Van den Bosch N, et al: A gas-chromatographic assay for the determination of 5, 6-dihydrofluorouracil and 5-fluorouracil in human plasma. *J Chromatogr* 278:283-289, 1983

20. Malet-Martino MC, Martino R, Lopez A, et al: New approach to metabolism of 5'-deoxy-5-fluorouridine in humans with fluorine-19 NMR. *Cancer Chemother Pharmacol* 13:31-35, 1984

21. Malet-Martino MC, Martino R, Bernadou J, et al: Doxifluridine mainly excreted in human bile as N-cholyl- α -fluoro- β -alanine, a novel metabolite. *Proc Am Assoc Cancer Res* 28:193, 1987 (abstr)

22. Sweeny DJ, Barnes S, Heggie GD, et al: Metabolism of 5-fluorouracil to an N-cholyl-2-fluoro- β -alanine conjugate: Previously unrecognized role for bile acids in drug conjugation. *Proc Natl Acad Sci USA* 84:5439-5443, 1987

23. Woodcock TM, Martin DS, Damin LAM, et al: Combination clinical trials with thymidine and fluorouracil: A phase I and clinical pharmacological evaluation. *Cancer* 45:1135-1143, 1980

24. Engelbrecht C, Lungquist I, Lewan L, et al: Modulation of 5-fluorouracil metabolism by thymidine in vivo and in vitro studies on RNA-directed effects in rat liver and rat hepatoma. *Biochem Pharmacol* 33:745-750, 1984

25. Tuchman M, Stoeckeler JS, Kiang DT, et al: Familial pyrimidinemia and pyrimiduria associated with fluorouracil toxicity. *N Engl J Med* 313:245-249, 1985

26. Santi DV, McHenry CA, Sommer H: Mechanism of interaction of thymidylate synthetase with 5-fluorodeoxyuridylate. *Biochemistry* 13:471-481, 1974

27. Danenberg PV: Thymidylate synthetase—a target enzyme in cancer chemotherapy. *Biochim Biophys Acta* 473:73-92, 1977

28. Danenberg PV, Lockshin A: Thymidylate synthetase—substrate complex formation. *Mol Cell Biochem* 43:49-57, 1982

29. Bapat AR, Zarov C, Danenberg PV: Human leukemic cells resistant to 5-fluoro-2'-deoxyuridine contain a thymidylate synthetase with lower affinity for nucleotides. *J Biol Chem* 258:4130-4136, 1983

30. Moore N, Ahmed F, Dunlap RB: Evidence for the existence of covalent nucleotide-thymidylate synthase complexes, identification of site of attachment, and enhancement by folates. *J Biol Chem* 261:12745-12749, 1986

31. Washtien WL, Santi DV: Assay of intracellular free and micromolecular bound metabolites of 5-fluorodeoxyuridine and 5-fluorouracil. *Cancer Res* 39:3397-3404, 1979

32. Spears CP, Shahinian AH, Moran RG, et al: In vivo kinetics of thymidylate synthetase inhibition in 5-fluorouracil sensitive and resistant murine colon adenocarcinomas. *Cancer Res* 42:450-456, 1982

33. Berger SH, Hakala MT: Relationship of dUMP and free FdUMP pools to inhibition of thymidylate synthase by 5-fluorouracil. *Mol Pharmacol* 25:303-309, 1984

34. Brockman RW, Chaddix SC, Rose LM: Biochemical aspects of chemotherapy of mouse colon carcinoma. Fluoropyrimidines and pyrazofurin. *Cancer* 40:2681-2691, 1977

35. Houghton JA, Maroda SJ, Phillips JO, et al: Biochemical determinants of responsiveness to 5-fluorouracil and its derivatives in xenografts of human colorectal adenocarcinomas in mice. *Cancer Res* 41:144-149, 1981

36. Priest DG, Ledford BE, Doig MT: Increased thymidylate synthetase in 5-fluorodeoxyuridine resistant cultured hepatoma cells. *Biochem Pharmacol* 29:1549-1553, 1980

37. Klubes P, Connely K, Cerna J, et al: Effects of 5-fluorouracil on 5-fluorodeoxyuridine-5'-monophosphate and 2'-deoxyuridine-5'-monophosphate pools and DNA synthesis in solid mouse L1210 and rat Walker 256 tumors. *Cancer Res* 38:2325-2331, 1978

38. Houghton JA, Weiss KD, Williams LG, et al: Relationship between 5-fluoro-5'-deoxyuridylate, 2'-deoxyuridylate and thymidylate synthase activity subsequent to 5-fluorouracil administration, in xenografts of human colon adenocarcinomas. *Biochem Pharmacol* 35:1351-1358, 1986

39. Evans RM, Laskin JD, Hakala MT: Effect of excess folates and deoxyinosine on the activity and site of action of 5-fluorouracil. *Cancer Res* 41:3288-3295, 1981

40. Houghton JA, Torrance PM, Radparvar S, et al: Binding of 5-fluorodeoxyuridylate to thymidylate synthase in human colon adenocarcinoma xenografts. *Eur J Cancer Clin Oncol* 22:505-510, 1986

41. Yin M-B, Zakrzewski SF, Hakala MT: Relationship of

cellular folate cofactor pools to the activity of 5-fluorouracil. *Mol Pharmacol* 23:190-197, 1983

42. Moran RG, Zakrewski S, Werheiser WC: Folate metabolism in mammalian cells in culture. Partial characterization of the folates present in mouse L1210 leukemia cells. *J Biol Chem* 251:3566-3575, 1976

43. Priest DB, Happel KK, Mangum M, et al: Tissue folyl-polyglutamate chain-length characterization by electrophoresis as thymidylate synthetase-5-fluorodeoxyuridylate ternary complexes. *Anal Biochem* 115:163-169, 1981

44. Radparvar S, Houghton PJ, Houghton JA: Characteristics of thymidylate synthase purified from a human colon adenocarcinoma. *Arch Biochem Biophys* 260:342-350, 1988

45. Spears CP, Gustavsson BG, Mitchell MS, et al: Thymidylate synthetase inhibition in malignant and normal liver of patients given intravenous 5-fluorouracil. *Cancer Res* 44:4144-4150, 1984

46. Mandel HG: The target cell determinants of the antitumor actions of 5-FU: Does FU incorporation into RNA play a role. *Cancer Treat Rep* 65:63-71, 1981 (suppl 3)

47. Wilkinson DS, Crumley J: Metabolism of 5-fluorouracil in sensitive and resistant Novikoff hepatoma cells. *J Biol Chem* 252:1051-1056, 1977

48. Maybaum J, Ullman B, Mandel HG, et al: Regulation of RNA- and DNA-directed actions of 5-fluoropyrimidines in mouse T-lymphoma (S-49) cells. *Cancer Res* 40:4209-4215, 1980

49. Cory JG, Breland JC, Carter GL: Effect of 5-fluorouracil on RNA metabolism in Novikoff Hepatoma cells. *Cancer Res* 39:4905-4913, 1979

50. Kufe DW, Major PP: 5-fluorouracil incorporation into human breast carcinoma RNA correlates with cytotoxicity. *J Biol Chem* 256:9802-9805, 1981

51. Spiegelman S, Sawyer R, Nayak R, et al: Improving the anti-tumor activity of 5-fluorouracil by increasing its incorporation into RNA via metabolic modulation. *Proc Natl Acad Sci USA* 77:4966-4970, 1980

52. Houghton JA, Houghton PJ, Wooten RS: Mechanism of induction of the gastrointestinal toxicity in the mouse by 5-fluorouracil, 5-fluorouridine and 5-fluoro-2'-deoxyuridine. *Cancer Res* 39:2406-2413, 1979

53. Glazer RI, Hartman KD: Analysis of the effect of 5-fluorouracil on the synthesis and translation of polysomal poly (A) RNA from Ehrlich ascites cells. *Mol Pharmacol* 19:117-121, 1981

54. Glazer RI, Hartman KD: In vitro translation of messenger RNA following exposure of human colon carcinoma cells in culture to 5-fluorouracil and 5-fluorouridine. *Mol Pharmacol* 23:540-546, 1983

55. Glazer RI, Lloyd LS: Association of cell lethality with incorporation of 5-fluorouracil and 5-fluorouridine into nuclear RNA in human colon carcinoma cells in culture. *Mol Pharmacol* 21:468-473, 1982

56. Glazer RI, Hartman KD: The effect of 5-fluorouracil on the synthesis and methylation of low molecular weight RNA in L1210 cells. *Mol Pharmacol* 17:245-249, 1980

57. Herrick P, Kufe DW: Lethality associated with incorporation of 5-fluorouracil into preribosomal RNA. *Mol Pharmacol* 26:135-140, 1984

58. Armstrong RD, Lewis M, Stern SG, et al: Acute effect of 5-fluorouracil on cytoplasmic and nuclear dihydrofolate reduc-

tase messenger RNA metabolism. *J Biol Chem* 261:7366-7371, 1986

59. Will CL, Dolnick BJ: 5-fluorouracil augmentation of dihydrofolate reductase gene transcripts containing intervening sequences in methotrexate-resistant KB cells. *Mol Pharmacol* 29:643-648, 1986

60. Kanamaru R, Kakuta H, Sato T, et al: The inhibitory effect of 5-fluorouracil on the metabolism of preribosomal and ribosomal RNA in L1210 cells in vitro. *Cancer Chemother Pharmacol* 17:43-46, 1986

61. Iwata T, Watanabe T, Kufe DW: Effects of 5-fluorouracil on globin mRNA synthesis in murine erythroleukemia cells. *Biochemistry* 25:2703-2707, 1985

62. Ingraham HA, Tseng BY, Goulian M: Mechanism for exclusion of 5-fluorouracil from DNA. *Cancer Res* 40:998-1001, 1980

63. Herrick DJ, Major PP, Kufe DW: Effect of methotrexate on incorporation and excision of 5-fluorouracil residue in human breast carcinoma DNA. *Cancer Res* 42:5015-5017, 1982

64. Major PP, Egan E, Herrick D, et al: 5-Fluorouracil incorporation in DNA of human breast carcinoma cells. *Cancer Res* 42:3005-3009, 1982

65. Danenberg PV, Heidelberger C, Mulkins MA, et al: The incorporation of 5-fluoro-2'-deoxyuridine into DNA of mammalian tumor cells. *Biochem Biophys Res Commun* 102:654-658, 1981

66. Kufe DW, Scott P, Fram R, et al: Biologic effect of 5-fluoro-2'-deoxyuridine incorporation in L1210 DNA. *Biochem Pharmacol* 32:1337-1340, 1983

67. Lönn U, Lönn S: Interaction between 5-fluorouracil and DNA of human colon adenocarcinoma. *Cancer Res* 44:3414-3418, 1984

68. Cheng YC, Nakayama K: Effects of 5-fluoro-2'-deoxyuridine on DNA metabolism in HeLa cells. *Mol Pharmacol* 23:171-174, 1983

69. Tanaka M, Kimura K, Yoshida S: Enhancement of the incorporation of 5-fluorodeoxyuridylate into DNA of HL-60 cells by metabolic modulations. *Cancer Res* 43:5145-5150, 1983

70. Kufe DW, Herrick D, Gunner L: Uracil enhancement of 5-fluorodeoxyuridine incorporation into human breast carcinoma deoxyribonucleic acid. *Biochem Pharmacol* 33:2329-2331, 1984

71. Sawyer RC, Stolfi RL, Martin DS, et al: Incorporation of 5-fluorouracil into murine bone marrow DNA in vivo. *Cancer Res* 44:1847-1851, 1984

72. Lönn U, Lönn S: DNA lesions in human neoplastic cells and cytotoxicity of 5-fluoropyrimidines. *Cancer Res* 46:3866-3871, 1986

73. Shuetz JD, Diasio RB: The effect of 5-fluorouracil on DNA chain elongation in intact bone marrow cells. *Biochem Biophys Res Commun* 133:361-367, 1985

74. Pogolotti AL, Nolan PA, Santi DV: Methods for the complete analysis of 5-fluorouracil metabolites in cell extracts. *Anal Biochem* 117:178-186, 1981

75. Peters GJ, Laurensse E, Lankelma J, et al: Separation of several 5-fluorouracil metabolites in various melanoma cell lines. Evidence for the synthesis of 5-fluorouracil nucleotide sugars. *Eur J Cancer Clin Oncol* 20:1425-1431, 1984

76. Iigo M, Kureitani K, Hoshi A: Relationship between anti-tumor effect and metabolites of 5-fluorouracil in combination

treatment with 5-fluorouracil and guanosine in ascites sarcoma 180 tumor system. *Cancer Res* 43:5687-5694, 1983

77. Peterson MS, Ingraham HA, Goulian M: 2'-Deoxyribosyl analogues of UDP-N-acetylglucosamine in cells treated with methotrexate or 5-fluoro-deoxy-uridine. *J Biol Chem* 258:10831-10834, 1983

78. Anukarahanonta T, Holstege A, Keppler DOR: Selective enhancement of 5-fluorouridine uptake and action in rat hepatomas in vivo following pretreatment with D-galactosamine and 6-azauridine or N-(phosphonacetyl)-L-aspartate. *Eur J Cancer* 10:1171-1180, 1980

79. Weckbecker G, Keppler DOR: Substrate properties of 5-fluorouridine diphospho sugars detected in hepatoma cells. *Biochem Pharmacol* 33:2291-2298, 1984

80. Reyes P, Hall TC: Synthesis of 5-fluorouridine 5'-phosphate by a pyrimidine transferase of mammalian origin. II. Correlation between the tumor levels of the enzyme and 5-fluorouracil-promoted increase in survival of tumor bearing mice. *Biochem Pharmacol* 18:2587-2590, 1969

81. Kessel D, Hall TC, Wodinsky I: Nucleotide formation as a determinant of 5-fluorouracil response in mouse leukemias. *Science* 145:911-913, 1966

82. Sobrero AF, Moir RD, Bertino JR, et al: Defective facilitated diffusion of nucleosides, a primary mechanism of resistance to 5-fluoro-2'-deoxyuridine in the HCT-8 human carcinoma lines. *Cancer Res* 45:3155-3160, 1985

83. Peters GJ, Laurensse E, Leyva A, et al: Purine nucleosides as cell-specific modulators of 5-fluorouracil metabolism and cytotoxicity. *Eur J Cancer Clin Oncol* 22:1869-1881, 1987

84. Fernandes DJ, Cranford SK: Resistance of CCRF-CEM cloned sublines to fluorodeoxyuridine associated with enhanced phosphatase activities. *Biochem Pharmacol* 34:125-132, 1985

85. Cohen A, Ullman B: Role of intracellular TTP levels in fluorodeoxyuridine toxicity. *Biochem Pharmacol* 33:3298-3301, 1984

86. Berger SH, Jenh CH, Johnson LF, et al: Thymidylate synthase overproduction and gene amplification in fluorodeoxyuridine-resistant human cells. *Mol Pharmacol* 28:461-467, 1985

87. Clark JL, Berger SH, Mittelman A, et al: Thymidylate synthase gene amplification in a colon tumor resistant to fluoropyrimidine chemotherapy. *Cancer Treat Rep* 71:261-265, 1987

88. Leyland-Jones B, O'Dwyer PJ: Biochemical modulation: Application of laboratory models to the clinic. *Cancer Treat Rep* 70:219-229, 1986

89. Ullman B, Kirsch J: Metabolism of 5-fluorouracil in cultured cells. Protection from 5-fluorouracil cytotoxicity by purines. *Mol Pharmacol* 15:357-366, 1979

90. Beltz RE, Waters RN, Hegarty TJ: Enhancement and depression by inosine of the growth inhibitory action of 5-fluorouracil on cultured Jensen tumor cells. *Biochem Biophys Res Commun* 112:235-241, 1983

91. Peters GJ, Laurensse E, Leyva A, et al: Fluctuations in phosphoribosyl-pyrophosphate levels. Effects of drugs. *FEBS Lett* 170:277-280, 1984

92. Santelli G, Valeriote F: In vivo potentiation of 5-fluorouracil cytotoxicity against AKR leukemia by purines, pyrimidines and their nucleosides and deoxynucleosides. *JNCI* 64:69-72, 1980

93. Iigo M, Ando N, Hoshi A, et al: Effect of pyrimidines, purines and their nucleosides on antitumor activity of 5-fluorouracil against L1210 leukemia. *J Pharm Dyn* 5:515-520, 1982

94. Sawyer RC, Stolfi RL, Spiegelman S, et al: Effect of uridine on the metabolism of 5-fluorouracil in the CD₈F₁ murine mammary carcinoma system. *Pharm Res* 2:69-75, 1984

95. Van Groeningen CJ, Leyva A, Peters GJ, et al: Reversal of 5-fluorouracil induced myelo-suppression by high-dose uridine. *Proc Am Assoc Cancer Res* 27:169, 1986 (abstr)

96. Peters GJ, Van Groeningen CJ, Laurensse E, et al: Effect of pyrimidine nucleosides on body temperatures of man and rabbit in relation to pharmacokinetic data. *Pharm Res* 4:113-119, 1987

97. Darnowsky JW, Handschumacher RE: Tissue specific enhancement of uridine utilization and 5-fluorouracil therapy in mice by benzylacyclouridine. *Cancer Res* 45:5364-5368, 1985

98. Darnowsky JW, Handschumacher RE: Tissue uridine pools: Evidence in vivo of a concentrative mechanism for uridine uptake. *Cancer Res* 46:3490-3494, 1986

99. Peters GJ, Van Groeningen CJ, Laurensse E, et al: Uridine-induced hypothermia in mice and rats in relation to plasma and tissue levels of uridine and its metabolites. *Cancer Chemother Pharmacol* 20:101-108, 1987

100. Peters GJ, Veerkamp JH: Purine and pyrimidine metabolism in peripheral blood lymphocytes. *Int J Biochem* 15:115-123, 1983

101. Peters GJ, van Dijk J, Laurensse EJ, et al: In vitro biochemical and in vivo biological studies of the uridine "rescue" of 5-fluorouracil. *Br J Cancer* 57:259-265, 1988

102. Casper ES, Vale K, Williams LJ, et al: Phase I and clinical pharmacological evaluation of biochemical modulation of 5-fluorouracil with N-(phosphonacetyl)-L-aspartate. *Cancer Res* 43:2324-2329, 1983

103. Schwartz PM, Handschumacher RE: Selective antagonism of 5-fluorouracil cytotoxicity by 4-hydroxypyrazolopyrimidine (allopurinol) in vitro. *Cancer Res* 39:3095-3101, 1979

104. Schwartz PM, Noir D, Hyde CM, et al: Role of uridine phosphorylase in the anabolism of 5-fluorouracil. *Biochem Pharmacol* 34:3585-3589, 1985

105. Kobayashi S, Hoshino T: Combined cytotoxic effect of low dose 5-fluorouracil and hydroxyurea on 9L cells in vitro. *Cancer Res* 43:5309-5313, 1983

106. Grem JL, Fischer PH: Augmentation of 5-fluorouracil cytotoxicity in human colon cancer cells by dipyrindamole. *Cancer Res* 45:2967-2972, 1985

107. Waxman S, Bruckner H: The enhancement of 5-fluorouracil antimetabolic activity by leucovorin, menadione and γ -tocopherol. *Eur J Cancer Clin Oncol* 18:685-692, 1982

108. Mini E, Moroson BA, Bertino JR: Cytotoxicity of floxuridine and 5-fluorouracil in human T-lymphoblast leukemia cells: Enhancement by leucovorin. *Cancer Treat Rep* 71:381-389, 1987

109. Nadal JC, Van Groeningen CJ, Pinedo HM, et al: Schedule-dependency of *in vivo* modulation of 5-fluorouracil by leucovorin and uridine in murine colon carcinoma. *Invest New Drugs* (in press)

110. Madajewics S, Petrelli N, Rustum YM, et al: Phase I-II trial of high dose calcium leucovorin and 5-fluorouracil in advanced colorectal cancer. *Cancer Res* 44:4667-4669, 1984

111. Cunningham J, Bukowsky RM, Budd GT, et al: 5-fluorouracil and folinic acid. A phase I-II trial in gastrointestinal malignancy. *Invest New Drugs* 2:391-395, 1984

112. Petrelli N, Herrera L, Rustum Y, et al: A prospective randomized trial of 5-fluorouracil versus 5-fluorouracil and

high-dose leucovorin versus 5-fluorouracil and methotrexate in previously untreated patients with advanced colorectal carcinoma. *J Clin Oncol* 5:1559-1565, 1987

113. Laufman LR, Kreczowski KA, Roach R, et al: Leucovorin plus 5-fluorouracil: An effective treatment for metastatic colon cancer. *J Clin Oncol* 5:1394-1400, 1987

114. Myers CE: The pharmacology of fluoropyrimidines. *Pharmacol Rev* 33:1-15, 1981

115. El Sayed YM, Sadée W: The fluoropyrimidines, in Ames MM, Powis G, Kovach JS (eds): *Pharmacokinetics of Anticancer Agents in Humans*. Amsterdam, Elsevier, 1983, pp 209-227

116. Peters GJ, Kraal I, Laurensse E, et al: Separation of 5-fluorouracil and uracil by ion-pair reversed-phase high performance liquid chromatography on a column with porous polymer packing. *J Chromatogr* 307:464-468, 1984

117. Finn C, Sadée W: Determination of 5-fluorouracil plasma levels in rats and man by isotope dilution-mass fragmentography. *Cancer Chemother Rep* 59:279-286, 1975

118. Kok RM, De Jong APJM, Van Groeningen CJ, et al: Highly sensitive determination of 5-fluorouracil in human plasma by capillary gas chromatography and negative ion chemical ionization mass spectrometry. *J Chromatogr* 343:59-66, 1985

119. Van Groeningen CJ, Pinedo HM, Heddes J, et al: Pharmacokinetics of 5-fluorouracil measured with a sensitive mass spectrometric method in patients during dose escalation schedule. *Cancer Res* (submitted)

120. Collins JM, Dedrick RL, King FG, et al: Non-linear pharmacokinetic models for 5-fluorouracil in man. *Clin Pharmacol Ther* 28:235-246, 1980

121. Van Groeningen CJ, Lankelma J, Kraal I, et al: Unpublished results

122. De Bruijn EA, Remeger L, Tjaden UR, et al: Non-linear pharmacokinetics of 5-fluorouracil as described by *in vivo* behavior of 5,6-dihydro-5-fluorouracil. *Biochem Pharmacol* 35:2461-2465, 1986

123. Ensminger WD, Rosowsky A, Raso V, et al: A clinical pharmacologic evaluation of hepatic arterial infusion of 5-fluoro-2'-deoxyuridine and 5-fluorouracil. *Cancer Res* 38:3784-3792, 1978

124. Collins JM: Pharmacokinetics of 5-fluorouracil infusions in the rat: Comparison with man and other species. *Cancer Chemother Pharmacol* 14:108-111, 1985

125. Schwartz PM, Turek PJ, Hyde CM, et al: Altered plasma kinetics of 5FU at high dosage in rat and man. *Cancer Treat Rep* 69:133-136, 1985

126. Malet-Martino MC, Martino R, Lopez A, et al: Metabolism of 5'-deoxy-5-fluorouridine in humans: Qualitative and quantitative analysis by fluorine-19 NMR. *Biochem Pharmacol* 34:429-430, 1985

127. Phillips TA, Howell A, Grieve RJ, et al: Pharmacokinetics of oral and intravenous fluorouracil in humans. *J Pharm Sci* 69:1428-1431, 1980

128. Boublil J-L, Milano G, Khater R, et al: Continuous 5-day regional chemotherapy by 5-fluorouracil in colon carcinoma: Pharmacokinetic evaluation. *Br J Cancer* 52:15-20, 1985

129. Speyer JL, Collins JM, Dedrick RL, et al: Phase I and pharmacological studies of 5-fluorouracil administered intraperitoneally. *Cancer Res* 40:567-572, 1980

130. Gyves J: Pharmacology of intraperitoneal infusion of 5-fluorouracil and mitomycin C. *Semin Oncol* 12:29-32, 1985 (suppl 4)