

Chapter

---

# 11

**Summary and General Discussion**



The aim of this thesis was threefold: i) to gain insight into the heterogeneity of depressive disorders by examining the diagnostic categories of Major Depressive Disorder (MDD), Dysthymic Disorder and Double Depression on a broad range of socio-demographic, biological and psychological characteristics, course trajectories and consequences for general well-being; ii) to explore alternative models for depressive disorders, based on data-driven techniques, and iii) to examine the association between depression and anxiety, and its implications for the nosology of depressive disorders. In the present chapter, the main findings of this thesis will be integratively summarized and discussed within the framework of outcomes of previous studies. Next, methodological issues, relevant for this thesis, will be addressed and some notes on possible clinical implications of our findings will be given. Finally, some topics for future research are identified.

## MAIN FINDINGS

### Part I. Depressive disorders: carving nature at its joints?

Plato stressed the importance of dividing things into classes along the ‘natural joints’, instead of recklessly ‘breaking any part’ in order to create boundaries between categories (*Phaedrus 265e*- around 370 BC). To what extent are the current depressive categories divided along natural occurring boundaries? In Chapter 2 through 5 we examined MDD, Dysthymic Disorder and Double Depression on a broad range of socio-demographic, biological and psychological characteristics, course trajectories and consequences for general well-being. If MDD, Dysthymia and Double Depression represent distinct clinical entities, differences in these parameters might be expected. Since symptoms used to classify Dysthymic Disorder and MDD in DSM-III and further editions are similar, albeit of longer duration and less severity in Dysthymic Disorder, significant differences in pathophysiological processes between MDD and Dysthymia may be attributed to these core clinical features of depression (severity and/or duration).

#### *Biology and Psychology*

In a review on the biological and characterological features of Dysthymia, Griffiths et al. (2000) concluded that there is scattered information on differences in the biological underpinnings of MDD and Dysthymic Disorder and stressed the importance to distinguish between pure Dysthymia and Double Depression in studies on the biological aspects of Dysthymia. However, studies that compare pure Dysthymic Disorder with Double Depression are limited. In **Chapter 2**, we compared persons with a 6-month diagnosis of MDD (n=853), pure Dysthymic Disorder (n=43) and Double Depression (n=262) on i) biological characteristics (including cortisol, inflammation markers and serum Brain Derived Neurotrophic Factor (sBDNF)) and ii) psychological characteristics (including personality measures and cognitive vulnerability). No differences were found on biological parameters suggesting that outcomes on biological parameters seem to be unrelated to depressive subtypes of MDD, Dysthymia and Double Depression, and, thus, seem to be unrelated to

severity and/or duration of depressive symptoms. This is in line with other reports, based on NESDA-data, in which severity and duration were unrelated to cortisol (Vreeburg et al., 2009), inflammatory response- with the exception of higher levels of TNF- $\alpha$  in women with more severe depressive symptoms- (Vogelzangs et al., 2012), and BDNF (Molendijk et al., 2011). Broader literature on the importance of severity and chronicity of depressive symptoms on biological parameters conflicts with both present (Brouwer et al., 2005; Dowlati et al., 2010; Oldehinkel et al., 2001) and absent associations (Anisman et al., 1999; Aydemir et al. 2007; Bhagwagar et al., 2005; Nelson et al., 1997; Marques-Deak et al., 2007; Watson et al., 2002). Note, the lack of differences does not imply that similar pathophysiological processes underlie MDD, Dysthymia and Double Depression. Possibly our findings on cortisol, inflammation and BDNF represent the endpoints of a final common pathway, with heterogeneity in underlying processes. Another explanation of lack of differences might be the insufficient discriminant validity of the depressive categories. As Carroll (1989) said: 'No biological measure can in principle do better than the clinical independent variable against which it is compared'. Hence, possible flaws in the concepts of MDD, Dysthymia and/or Double Depression limit the detection of actual differences between these subgroups of depressive disorders.

Persons with MDD, Dysthymia and Double Depression showed different psychological profiles, with higher levels of neuroticism in the latter two, and an absence of differences in cognitive styles between Dysthymic Disorder and Double Depression, suggesting that personality and cognitive styles are associated with chronicity. This is in line with previous research (Wiersma et al., 2011). Unfortunately, causality cannot be inferred from our cross-sectional data. Cognitive theorists have proposed that maladaptive cognitive processing becomes more likely as more or chronic depressive episodes are experienced (Moulds et al., 2008). To what extent differences in personality and cognitive styles represent a state or trait effect is topic of debate. Ormel et al. (2004) previously demonstrated that persons with (a history of) MDD report higher levels of neuroticism when they are depressed than when they are not depressed, whereas others concluded that personality traits assessed during depressive episodes are a valid reflection of personality pathology rather than an artifact of depressed mood (Morey et al., 2010). Recently, NESDA researchers examined associations between specific personality dimensions and depressive disorders. They demonstrated that neuroticism, extraversion, and conscientiousness are not only predispositions of affective disorders, but they also appear to be subject to change with onset and recovery of depression (Karsten et al., 2012). Further, prospective research is needed to unravel the interplay between personality and depressive subtypes.

*Course trajectories of depressive disorders and functioning*

Longitudinal observations are of great importance for classifying mental disorders: when naturalistic course differs across disorders, their distinction could be useful in clinical practice and psychiatric nosology. In **Chapter 3**, we investigated the course trajectories of persons with a Composite International Diagnostic Interview (CIDI) diagnosis of Major Depressive Disorder (n=225), Dysthymic Disorder (n=62) or Double Depression (n=113) at baseline, with outcome defined as i) a CIDI diagnosis of depression after 3-year follow-up, and ii) the 3-year course of mental functioning, measured by the SF-36 mental subscale. Data were derived from NEMESIS-1. Cox proportional hazards analyses and Linear Mixed Models showed that Dysthymia and Double Depression had similar course trajectories, which were worse than the course trajectories of MDD only. These results do not support a distinction between Dysthymia and Double Depression. Multivariate analyses demonstrated the importance of neuroticism and global functioning in predicting the course of depressive disorders. What do these results add to the debate on the nosology of depressive disorders? Since the presence of a superimposed MDD does not alter the course of a pure Dysthymic disorder, the validity of a distinction between Dysthymia and Double Depression is not supported. However, both chronic disorders differed from MDD, representing non-chronic depression. To conclude, our findings are in line with Wells' proposal to differentiate between depressive disorders on the basis of functional status and chronicity (Wells, 1992), instead on stringent DSM-categories.

**Chapter 4** further elaborates on the association between level of functioning and type of depressive disorder. It is well known that depressive disorders have a large impact on both social and physical domains of functioning and work performance (Coryell et al., 1993; Judd et al., 1996; Rapaport et al., 2005; De Graaf et al., in press), similar to or even exceeding the impact noted in common somatic illnesses (Buist-Bouwman et al., 2006; Merikangas et al., 2007; Ormel et al., 1994). Since persistence of a lower level of functioning predicts recurrence of a depressive episode, even after the symptoms of depression are alleviated (Faravelli et al., 1986; Judd et al., 2000; Judd and Akiskal, 2000; Solomon et al., 2004; Spijker et al., 2004), insight into the course of social and physical functioning of persons with depressive disorders, and into the determinants of an impaired recovery of functioning, may facilitate recurrence prevention, limit the burden of disease, and informs us on the nosological status of the concepts under study. If for example substantial differences between level of functioning between MDD, Dysthymia and Double Depression are encountered, this would support the validity of the distinction between these subtypes.

Using data from NEMESIS-1, we compared 3-year trajectories of functioning across depressive persons with MDD (n=102), Dysthymia (n=66) and Double Depression (n=73), who recovered during three year follow-up, and compared these with persons without any DSM-diagnosis. Functioning was assessed using the Groningen Social Disability Schedule (GSDS) (Wiersma et al., 1988) and the SF-36 physical health summary-scale (Ware et al.,

1995). Data were derived from NEMESIS-1. Linear Mixed Models showed that compared to persons without any diagnosis, all depressed groups were significantly impaired on social and physical functioning. Across depressive groups, Dysthymic Disorder and Double Depression had a lower level of post-morbid physical functioning compared to MDD. Determinants for impaired social functioning (neuroticism) and impaired physical functioning (older age, somatic comorbidity and neuroticism) were identified. To conclude, chronicity of symptoms appears to be associated with impaired recovery of functioning to a greater extent than severity of symptoms, since especially those with Dysthymia (either with or without a superimposed MDD) showed slower and less complete recovery of functioning.

*Is Dysthymic Disorder worth the classification as a specific DSM-disorder?*

The findings of Chapter 2 through 4 allude to the questionable discriminant validity of Dysthymic Disorder - and provide further fuel to the debate currently being considered for DSM-V that the term 'Dysthymia', 'Double Depression' and 'Chronic Major Depression' be abandoned and subsumed into an enveloping 'Chronic Depressive Disorder'. However, if this container concept overrides identification of intrinsically differing conditions that might argue for quite different interventions, its clinical relevance is compromised. Therefore, **Chapter 5** seeks to further elucidate the concept of Dysthymic Disorder, assuming it is a broad diagnostic domain subsuming multiple expressions of clinical, chronic depression, such as minor chronic as well as 'smouldering' depressive episodes, treatment resistant depressive conditions, and depression syndromes underpinned by psychological and social factors. 318 Patients, attending the Black Dog Institute Depression Clinic, Sydney, and fulfilling the MINI-criteria of Dysthymic Disorder, were examined on a broad range of characteristics, including socio-demographics, personality measures and clinical characteristics. Latent Class Analysis (LCA) and Latent Profile Analysis (LPA) were conducted, with the aim of detecting distinct classes within Dysthymic Disorder, based on depressive symptomatology and personality domains, respectively. However, due to large heterogeneity we could not identify distinct conditions (other than variation by severity). Finally, clinicians' formulations of the patients were examined. Remarkably, none of the clinicians diagnosed a Dysthymic Disorder. These findings further add fuel to the debates on the questionable discriminant validity and clinical utility of Dysthymia in a tertiary referral clinic.

## **Part II. Alternative models for depressive disorders**

In Chapter 2 through 5, we examined current DSM-categories of MDD, Dysthymic Disorder and its composite- Double Depression. Since 1980, course was incorporated into the diagnostic criteria of these depressive subtypes, by requiring a minimum duration of depressive symptoms of two weeks for MDD and two years for Dysthymic Disorder. However, do DSM-categories of MDD, Dysthymic Disorder and Double Depression indeed adequately represent clinically relevant course trajectories of depression? Data-driven

techniques, free from any a priori assumptions or diagnostic categories, can help to detect naturally occurring course trajectories among persons with depression. In **Chapter 6**, we compare DSM-categories (MDD, Dysthymic Disorder and Double Depression) with empirically derived prognostic categories, using a prospectively followed cohort of depressed patients (n=804). Latent Class Growth Analysis (LCGA) identified five distinct course trajectories, ranging from mild severity and rapid remission to high severity and chronic course trajectory. These identified classes were compared with DSM-diagnoses. Contrary to expectations, over 50% of persons with Dysthymia and Double Depression were allocated to classes with favorable course trajectories, suggesting that current DSM-categories do not adequately represent clinically relevant course trajectories. The class with the most favorable course trajectory differed on a number of characteristics from other classes (younger age, more females, less childhood adversity, less somatic illnesses, lower neuroticism, higher extraversion). Older age, earlier age of onset and lower extraversion predicted poorest course trajectory.

Thus far, we have focused on specifiers of depressive disorders such as severity and duration, reflected in the diagnostic categories of MDD, Dysthymic Disorder and Double Depression. However, since DSM-III, other specifiers such as melancholic or atypical features are available for further differentiation of depressive subtypes. Over the past several decades, there has been substantial research devoted to the identification of depressive subtypes based on distinct symptom profiles, treatment response, and clinical correlates. Using Latent Class models in a large cohort (n=818) of subjects with depression, NESDA researchers previously demonstrated the importance of depression severity (moderate versus severe) and the nature of depressive symptoms (melancholic versus atypical) as differentiators between depressive subtypes (Lamers et al., 2010). However, few studies have investigated the stability of depressive subtypes, an essential requirement for the validity of the delineation of subtypes. Studies that examined the longitudinal stability of depressive subtypes have shown low to moderate stability of subtypes (Young et al., 1987; Coryell et al., 1994; Angst et al., 2007), with greater stability between adjacent episodes (Coryell et al., 1994). **Chapter 7** concerns the stability of depressive subtypes of a sample of 488 persons from NESDA who had a Major Depressive Disorder at both baseline and at the 2-year follow-up assessment. A Latent Transition Analysis (LTA) was applied to examine the stability of depressive subtypes across time-points. Overall LTA showed 76% stability across 2-year follow-up, with the greatest stability of the Severe Atypical class (79%). Persons hardly changed from Moderate to Severe Typical, and from Severe Typical to Severe Atypical. Transitions from the Severe Atypical to Severe Typical subtype may represent the phenomenon that endogenous/melancholic symptoms increase with increasing age (Brodaty et al. 1997; Parker et al. 2001). Analyses of correlates in stable subtypes showed a preponderance of women and more overweight and obesity in the Severe Atypical subtype, and a greater number of negative life events, and higher neuroticism and functioning scores

in the Severe Typical subtype. Subtypes of Major Depressive Disorder were found to be stable across 2-year follow-up and to have distinct determinants, supporting the notion that the identified subtypes are clinically meaningful.

### **Part III. Depression and Anxiety: same side of the coin?**

**Chapter 8** illustrates the importance of anxiety for course trajectories of depressive disorders. In this chapter, 303 respondents, who participated both in NEMESIS-1 and NESDA, with a depressive and/or anxiety CIDI-disorder were interviewed, examining the 7-year course of depression (n=141), anxiety (n=102) and their comorbid state (n=60) and possible prognostic factors. After 7 years, 60.7% of the subjects were free from a 12-month CIDI depression or anxiety diagnosis. The risk for a depression or anxiety at follow-up was higher for subjects with anxiety only or with comorbidity compared to subjects with depression only. Furthermore, low physical functioning and high neuroticism predicted the presence of a diagnosis after 7 years. During 7-years follow-up, 37.3% of the subjects were free from depressive and anxiety symptoms according to the Life Chart Interview (Lyketsos et al., 1994). Again, (comorbid) anxiety resulted in a poorer course. High neuroticism and childhood adversity predicted more follow-up time with symptoms.

In Chapter 8, anxiety was broadly defined, hence limiting the insight into the impact of specific anxiety disorders on course trajectories. In **Chapter 9**, anxiety was defined as panic attacks, in accordance with DSM-V's proposal to rate panic as a dimension across all mental disorders. Since the main purpose of this thesis is to gain insight into the heterogeneity of depressive disorders, we here only consider the associations of panic with affective disorders. Results showed a considerable impact of current panic attacks, and- to a lesser extent- of a history of panic attacks on the prevalence, 3-year onset and persistence, and functioning of affective disorders. Further research is needed to unravel the nature of this association: Is panic a severity marker of depressive disorders, conceivably representing a distinct subtype (Kessler et al., 1998), or a comorbid condition? And is the occurrence of panic prior to the onset of depression a proxy for some underlying causal risk factor (Kessler et al., 1998), possibly a symptom of shared genetical vulnerability? Irrespective of the nature of the association between anxiety and depression, results of Chapter 8 and 9 strongly point at the relevance of anxiety for course trajectories of depressive disorders, and hence, its contribution to the heterogeneity of depressive disorders.

Finally, the recognition of the large comorbidity between depression and anxiety also questioned the validity of a categorical distinction between specific depressive and anxiety categories (Goldberg, 1996), in particular Generalized Anxiety Disorder (GAD) and depression. In **Chapter 10**, Latent Class Analysis was applied to anxious and depressive symptomatology of respondents, participating in i) three large, population based cohorts (NSMHWB-2007, NCS-R and NEMESIS-2) and ii) a multi-site naturalistic cohort of adults



(NESDA) to examine whether GAD and Dysthymic Disorder can be differentiated into an anxiety versus a depression symptom cluster based on data driven methodology. A three-class and two-class model respectively best fitted the data, reflecting mainly different levels of severity of symptoms. No division into specific anxious versus depressive profiles emerged. The absence of a clear demarcation of an anxiety versus depressive profile in empirically derived classes questions the current categories of GAD and Dysthymic Disorder.

## **IN SUMMARY**

This thesis demonstrated the heterogeneity of depressive disorders and the role of anxiety on (course trajectories of) depressive disorders. It provided insight into the correlates of chronicity, and alternative models for the classification of depressive disorders were explored. We demonstrated that our current depression categories seem to exist in a kind of no man's land, disconnected from the underlying clinical entities and imposing artificial boundaries without clinical utility (McGorry et al., 2007). Illustration of the latter was the remarkable finding, that none of the clinicians in a tertiary referral clinic diagnosed Dysthymic Disorder in a population defined by the MINI as fulfilling the criteria of Dysthymia (Chapter 5). Furthermore, the limited discriminant validity of Dysthymic Disorder versus Double Depression was demonstrated; demarcation from MDD was more evident. Duration and severity of depressive symptoms and anxiety were identified as clinically relevant dimensions, in combination with assessment of correlates of chronicity, they may be preferable over the existing various diagnostic categories. These findings are in line with previous studies that found that basic clinical factors, such as severity and duration of the index episode are among the most consistent and strongest predictors of course (Spijker et al., 2002, 2004; Vuorilehto et al., 2009).

## **METHODOLOGICAL CONSIDERATIONS**

In the following, we (re)consider the most important strengths and limitations of our conducted studies, such as thoughts on sampling and non-response, measurement of psychopathology and statistical methods.

### *Sampling and non-response*

This thesis was based on several large, nationally representative surveys. In NEMESIS-1 and -2 a multistage, stratified, random sampling procedure was followed. Likewise, in the NSMHWB-2007, Australia, and the NCS-R, the United States, respondents were selected from a multistage clustered area probability sample of households. These sampling procedures enabled generalizability into the general population. The Netherlands Study of Depression and Anxiety (NESDA) is a multi-site naturalistic cohort study of adults (aged 18 to 65 years) recruited from the general population, general practices, and mental health organizations. The Black Dog Institute (BDI) sample included depressed out-patients,

recruited through a Depression Clinic, Sydney. In all surveys, long-term institutionalized persons and persons, who are not sufficiently fluent in Dutch (NEMESIS/ NESDA) or English (NSMHWB-2007/ NCS-R/ BDI), were excluded. Hence, generalizations towards institutionalized persons and some ethnic minorities are of limited legitimacy. In addition, as the subjects in these studies were adults, results have limited value for the elderly.

Non-reponse patterns in the baseline samples of NEMESIS-1 revealed a response rate at the baseline measurement of 70% (Bijl et al., 1998). In total 7076 persons participated; 5618 could be re-interviewed after one year ( $T_1$ ) (79.4%) and 4796 after three year ( $T_2$ ) (85.4% of  $T_1$ ) (Graaf et al., 2000). Of NEMESIS-2 only baseline data were available for the present thesis, consisting of 6646 persons (response rate: 65%). The NSMHWB-2007 consisted of 8841 respondents (response rate: 60%); the total sample of NCS-R consisted of 9282 persons (response rate: 71%). Detailed analyses of non-response and attrition were provided in the previous chapters. In addition, Lamers et al. (2012) described attrition in NESDA in more detail. Overall, psychopathology only slightly increased the probability of loss to follow-up. Attrition rates in Chapter 8, presenting the results on 7-year follow-up of depression and anxiety, were considerable. This merits further attention: Of the 662 subjects approached, 359 (54.2%) refused to participate and 303 (45.8%) could be followed up. However, those who were followed up did not differ in terms of age ( $p=.77$ ), gender ( $p=.79$ ), or type of baseline disorder (anxiety, depression or comorbid disorder,  $p=.97$ ) from those not participating. To conclude, we feel that our results on depressive subtypes are not appreciably affected by attrition.

#### *Measurement of depression and anxiety*

In all surveys, psychopathology was measured using standardized, diagnostic instruments, like Composite Interview Diagnostic Instrument (CIDI)- version 1.1; 2.1; and 3.0 (respectively: World Health Organization, 1990; 1998; Kessler et al., 2004 ) and Mini-International Neuropsychiatric Interview (MINI) (Lecrubier et al., 1997). The use of a structured instrument in large-scale epidemiological studies is a major strength. Inter-rater reliability for the two core concepts of this thesis, MDD and Dysthymic Disorder, has shown to be excellent (using the CIDI:  $k=0.97$  for MDD;  $k=0.96$  for Dysthymia (Wittchen et al., 1991; Wittchen, 1994); using the MINI:  $k=1.00$  for MDD, Dysthymia: to our knowledge not available (Sheehan et al., 1998)). However, some have argued that the use of community-based diagnostic instruments, administered by lay interviewers (as the CIDI), instead of a clinical interview by a clinician, results in detection of subthreshold cases with limited clinical significance (Regier et al., 1998). Therefore, in this thesis, in addition to DSM-categories, course was additionally defined using a continuous mental health score (Chapter 2) or Life Chart Information (Chapter 6, Chapter 8) with similar results, illustrating that our results are unlikely due to the choice of our diagnostic instrument.

Furthermore, this thesis was mainly longitudinal in its approach, with follow-up periods ranging from 2 to 7 years. In the course of time, some changes were made in the different versions of the CIDI. However, the diagnostic criteria for MDD and Dysthymic Disorder remained largely unchanged. In DSM-IV, criterion C was added, requiring that the depressive symptoms should have an impact on social and occupational functioning, hence, the threshold for fulfilling the criteria of either MDD or Dysthymic Disorder became somewhat higher in DSM-IV. In addition, some changes were made in course specifiers for MDD, and the specifier of primary versus secondary Dysthymic Disorder was abandoned. However, these changes in specifiers did not influence the criteria for fulfilling either a MDD or Dysthymic Disorder, and, therefore, did not influence the findings of this thesis. Relevant for Chapter 9 was the modification of the criteria for panic disorder in CIDI 2.1, requiring recurrent unexpected panic attacks accompanied by a month or more of persistent concern of having attacks or about the implications of an attack, whereas in CIDI 1.1, based on DSM-III-R, either four attacks in 4 weeks, or one attack followed by a month of persistent fear for having another attack, was required (WHO 1990, 1998). These changes resulted in a stricter definition of panic disorder at follow-up and might, therefore, result in an underestimation of the long-term persistence of anxiety disorders. Hence, our conclusion that a (comorbid) anxiety disorder resulted in a poorer course trajectory would have been stronger when the same diagnostic criteria would have been used.

In our search for predictors of course, we were able to use an extensive assessment battery, facilitating multivariate analyses on a range of potential risk factors. However, treatment was not included in our multivariate analyses, although it may be associated with course trajectories. In our naturalistic study designs, the effect of treatment on course cannot be adequately evaluated. In addition, adjustment for treatment effect is complex because of the bias, that patients with the most severe problems receive the most treatment. Hence, a spurious relation between treatment and outcome can be found and its interpretation becomes very difficult.

### *Statistical methods*

Various data-driven methods were applied in order to explore alternative models for the classification of depression. Chapter 6 presents one of the largest studies to date, that longitudinally investigates the existence of depressive course types in a cohort of depressed subjects by data-driven methods (LCGA). In addition, to our knowledge, Chapter 7 is the first to evaluate the prospective stability of depressive subtypes by Latent Transition Analysis. Finally, Chapter 10 relies on the aggregation of three major, general population samples from three different continents, enabling us to perform Latent Class Analysis on a sufficient number of participants. Despite these innovative procedures, when interpreting the results of these studies, several limitations should be taken into consideration. It should be noted that data-driven methods draw upon mathematical models that cannot account for a great

variety of fluctuations in symptom levels. There is evidence that depressive symptoms wax and wane within the same patient and that these symptomatic periods are interspersed in the overall course within times when patients are remitted and symptom free (Judd and Akiskal, 2000). This pleiomorphic course cannot completely be described by LCGA. Also, some potential limitations of the LTA should be noted. We used 2 time-points and had relative stable subtypes. Although two time points suffice for LTA analysis, it is possible that if we had had more follow-up measurements, more transitions would have occurred, and that the current results are therefore an overestimation of the true stability. Second, we evaluated course trajectories (Chapter 6) and transitions (Chapter 7) across a rather short period of two years. Although it has been reported that most symptom changes generally occur in the first year after the start of an episode (Penninx et al., 2011), future research should evaluate whether the course trajectories of symptomatology is indeed similar when longer time periods are used.

### **IMPLICATIONS FOR CLINICAL PRACTICE AND THEORY**

What are the possible implications of our findings for clinical practice? The most important message of this thesis concerns the lack of validity and utility of our current diagnostic system. This has great implications for (the organization of) mental health care, since diagnostic categories are the back and bone of patient care and clinical management. For example, many mental health organizations have organized their out-patients care along the lines of DSM-diagnostic groups, such as mood disorders or anxiety disorders, artificially drawing a line between depression and anxiety. Likewise, clinical guidelines and studies into intervention strategies, such as randomized clinical trials, strictly adhere to diagnostic categories.

We demonstrated that dimensions of duration and severity of depressive symptoms, and possibly an additional, dimensional rating of panic across mental disorders, may be preferable over the existing depressive categories. This requires a major shift in psychiatric thinking, since it undermines current clinical taxonomy and provides fuel for the development of alternative clinical models. We elaborate on possible models in the next paragraph, but what is its impact on patient care? Currently, Procrustean diagnostic concepts guide clinical practice. The knowledge gained in this thesis showed that, instead of counting the number of diagnostic criteria to see which DSM-category fits the complaints of individual patients, a more prominent role for rating scales of severity of depression and panic, and a thorough assessment of the psychiatric history, including assessment of duration and recurrence of symptoms, may be more useful to identify persons at risk for chronic course trajectories.

Currently, similar strategies are already in use in mental health care. For example, clinical guidelines for Bipolar Disorder (Nolen et al., 2008) propagate the use of a so-called ‘action

plan' (Perry et al., 1999) in which the nature and timing of prodromal symptoms of a manic or depressive relapse are elicited in order to seek earlier treatment and to reduce the time to next relapse. In this action plan, dimensions of severity and duration of symptoms guide interventions. This personalized intervention strategy has proven to be very effective in relapse prevention of manic episodes and improved social functioning and performance in employment (Perry et al., 1999). Likewise, a patient oriented approach, with input of severity measures (such as Hamilton Depression Rating Scale) and a thorough assessment of psychiatric history providing insight in the age of onset, relapse rate and duration of symptoms thus far, may provide a framework for individualized 'action plans' for depressed persons and facilitates the identification of persons at risk for chronic course trajectories.

However, severity and duration dimensions alone might not suffice. Shankman and Klein (2002) demonstrated that a combination of symptom severity and prior course provided only a minimal increase in predictive power over and above DSM diagnoses of Major Depressive Disorder and Dysthymia. They propagated inclusion of additional features, like personality and cognitive measures, functioning or biological correlates. We identified some correlates of poor course trajectories that may ameliorate the predictive power of dimensional models, such as older age (Chapter 6), early age of onset (Chapter 6), lower level of functioning (Chapter 3, 8), somatic illnesses (Chapter 4, 6), high neuroticism and low extraversion (Chapter 4, 6, 8) and childhood adversity (Chapter 6, 8). Most correlates were previously also identified as predictors for chronicity (see also Sargeant et al., 1990; Ormel et al., 1993, Keitner et al., 1992; Wiersma et al., 2009). The diversity in correlates of chronic course illustrates, that treatment focused on depressive symptoms only will not suffice to reduce risk of recurrence or chronicity and to enhance functioning of depressed individuals. Successful treatment must go beyond ameliorating signs and symptoms to address the broader issue of restoration of health. Among others, in clinical care, attention should be paid to personality dimensions, level of functioning and childhood adversity as significant predictors of chronic course trajectories of depression in the general population.

#### *Theoretical implications: an alternative model for depression?*

After Michelangelo finished his masterpiece, the Pieta, in 1499, he commented: "I saw the angel in the marble and carved until I set him free." In psychiatry, during the last decades, researchers sought to detect clinical entities hidden behind symptoms of disturbed mood, anxiety, delusions and so on, and to create classification systems for psychopathology. A challenging endeavor. Some mental disorders were defined, to be abandoned again several years later. In this thesis we navigated through the landscape of depression using triangulation of data and methods and demonstrated the flaws in current DSM-categories. Where to go next? Instead of endlessly searching for our own Pieta's, we might as well wonder: Is there a Pieta hidden behind the symptoms of disturbed mood? Or should we adhere to a more dimensional approach?

Based on the work of Fava and Kellner (1993), a clinical staging framework as an alternative model for current DSM-classification has been previously proposed (Fava and Tossani, 2007; McGorry, 2007; Hetrick et al., 2008). Clinical staging, already a proven strategy in the treatment of malignancies, defines the progression of disease in time and where a person lies along this continuum of the course of illness (McGorry, 2007). Defining discrete stages according to the progression of disease creates a prevention-oriented framework. This requires an accurate understanding of the broad social, biological and personal risk and protective factors that influence movement across stages (McGorry, 2007). Fava and Tossani (2007) developed an initial clinical staging framework for depression. This model was still based on current depressive categories (see Table 1).

**Table 1.** Stages of primary unipolar depression proposed by Fava and Tossani (2007).

Stages
1. Prodromal phase (anxiety, irritable mood, anhedonia, sleep disorders)
a. No depressive symptoms
b. Minor Depression
2. Major Depressive Disorder
3. Residual Phase
a. No depressive symptoms
b. Dysthymia
4.      a. Recurrent Depression
b. Double Depression
5. Chronic Major Depressive Episode (lasting at least 2 years without interruptions)

(Source: Fava and Tossani, 2007)

Our findings demonstrated the lack of validity and clinical utility of the current depressive categories, hence questioning the model of Fava and Tossani (2007), since this model was based on these traditional concepts. A clinical staging model, incorporating duration and severity dimensions, as well as an additional dimension of anxiety, in association with correlates of chronicity, such as neuroticism and level of functioning may be more valid. Hetrick et al. (2008) provided a heuristic model for unipolar depressive disorders (see Table 2). This model employs a more dimension-based approach, including depression severity and functional status. In addition, the role of anxiety is reckoned, at least in stage 0 and 1.

**Table 2.** Clinical staging model framework for unipolar depressive disorders proposed by Hetrick et al. (2008).

Stages	
0.	Increased risk of anxiety or depressive disorder. No symptoms currently.
1.	<ol style="list-style-type: none"> <li>a. Mild or non-specific symptoms of anxiety or depression, including neurocognitive deficits of severe mood disorder. Mild functional change or decline.</li> <li>b. Ultra high risk: moderate but subthreshold symptoms of anxiety or depression, with moderate neurocognitive changes and functional decline to caseness (Global Assessment of Functioning [GAF] &lt; 70)</li> </ol>
2.	First episode of Major Depressive Disorder. Full threshold disorder with moderate to severe symptoms, neurocognitive deficits and functional decline (GAF 30-50)
3.	<ol style="list-style-type: none"> <li>a. Incomplete remission from first episode of care. Could be linked or fast-tracked to Stage 4.</li> <li>b. Recurrence of relapse of depressive disorder which stabilizes with treatment at a level of GAF, residual symptoms, or neurocognition below the best level achieved following remission from first episode</li> <li>c. Multiple relapses, provided worsening in clinical extent and impact of illness is objectively present.</li> </ol>
4.	Severe, persistent OR unremitting illness as judged on symptoms, neurocognition and disability criteria.

Note: The clinical staging model also includes target populations for recruitment, potential interventions and indicative biological and endophenotypic markers (see Hetrick et al., 2008).

According to the findings of this thesis, we would promote the inclusion of correlates of chronicity like personality measures (high neuroticism, low extraversion), older age, earlier age of onset, somatic comorbidity and childhood adversity as markers for further differentiation of persons at risk for progression from stage 2 to 3 or 4. Future research may further refine the clinical staging model for mood disorders, and may result in clearly quantifiable stages guiding the use of appropriate stage-specific, evidence-based treatment (Hetrick et al., 2008). First results are promising: Fournier et al. (2010) showed that the magnitude of benefit of antidepressant medication with placebo increases with the severity of depression symptoms, demonstrating the benefits of dimensional approaches above stringent DSM-categories.

Finally, the prominent role of anxiety in the classification of depression merits further attention. This thesis suggests that dimensionally rating anxiety, in casu panic, across all mental disorders might be of great value for clinical care. This approach is not new. In 1960s and 1970s depressed patients were subtyped according to the presence or absence of anxiety symptoms rather than being given separate diagnoses (Paykel, 1972; Overall et al., 1966). Recently, others have argued for a 3-dimensional structure (Clark and Watson, 1991) or a 2-dimensional structure of an internalizing dimension, encompassing Major Depressive Disorder, Dysthymia and anxiety disorders, and externalizing dimension (Krueger, 1999; Vollebergh et al., 2001; Olino et al., 2012). Chapter 10 contributes to this debate by showing that at least GAD and Dysthymic Disorder belong to one diagnostic domain. To what extent a dimensional rating of panic across mental disorders, including mood disorders, suffices, or

a reclassification of psychopathology into a tripartite model, or an internalizing versus externalizing dimensional model is needed, merits further research. Recently, Wardenaar et al. (2012) demonstrated the predictive value of dimensions of the tripartite model (general distress, anhedonic depression and anxious arousal) (Clark and Watson, 1991) on 2-year course trajectories of depression and anxiety on top of diagnosis and other prognostic factors. These findings warrant further replication. Knowledge to be gained by this research should be integrated into the staging framework for affective disorders.

## **FUTURE RESEARCH**

To date, the practical application of clinical staging of mood disorders largely eluded psychiatry (Hetrick et al., 2008). Hence, future research should further refine a clinical staging framework for mood disorders, and examine its validity and utility for clinical practice and research. Thorough knowledge on broad social, biological, genetic, and personal risk and protective factors, that influence progression from one stage to the next, is needed. The more information is included in the model, the stronger its predictive value (Fava et al., 2012). Brietzke et al. (2012) recently proposed a predictive algorithm for bipolar disorders integrating genetic (i.e. family history), environmental (e.g. childhood maltreatment) and biological markers (i.e. BDNF, inflammatory and oxidative stress markers). Similar proposals should be developed for unipolar depressive disorders. Furthermore, we need to know the relative potency of the identified risk factors and which of them may be responsive to current interventions. Some factors may operate across several or all stages, others may be stage specific (McGorry et al., 2007). For example, socio-demographics such as age, female gender and low education are all confirmed determinants of the onset of depressive disorders (Kessler et al., 1994; Seedat et al., 2009). However, their role in course prediction has been less consistent (Ramsawh et al., 2009; Spijker et al., 2002; Yonkers et al., 2003).

In oncology, clinical staging depends on the type of malignancy. For example, lung cancers are broadly classified into two types: small cell lung cancers (SCLC) and non-small cell lung cancers (NSCLC). This classification is based upon the microscopic appearance of the tumor cells themselves. These two types of cancers grow and spread in different ways and may have different treatment options, so a distinction between these two types is important. To what extent distinction between typical and atypical depression (Chapter 7), or a division into melancholic versus non-melancholic depression (Parker et al., 2010), warrants distinct clinical staging models, merits further research. Likewise, the subjects in this thesis were adults, participating in general population studies or out-patients settings. Knowledge from severely depressed in-patients as well as depressed elderly is needed to assess whether a clinical staging framework fits all types of depression, at all ages.



Research into intervention strategies can greatly benefit from this alternative approach. Thus far, our official diagnostic systems have impeded the search for neurobiological, genetic and psychosocial risk factors, and treatment. In clinical practice, dissemination of a clinical staging approach, such as treatment algorithms and 'action plans', can help to prevent mood disorders to progress, and to personalize treatment for individual patients along the lines indicated by the staging model.

In this thesis, data-driven methods, such as LCGA and LTA, were applied. Distinct course trajectories and depressive subtypes were identified. To examine whether these identified subtypes represent clinically relevant entities or dimension, an examination of these subtypes on a broad range of socio-demographic, biological, genetic and psychological characteristics, course trajectories and consequences for general well-being is needed. For example, in Chapter 6, our fifth class (chronic, severe) was characterized by greater severity, an early age of onset and a higher familial loading, possibly identifying a genetically more homogenous group. Lyons et al. (1998) and more recently Kendler et al. (2009) also suggested that early age-of-onset does typify a subtype of depressive disorders, with poorer course trajectories. This may be informative for etiological research. Likewise, Vogelzangs et al. (2012) for example demonstrated immune dysregulation has a role in a subgroup of depressed persons, in particular in men with a late-onset depression. Selection of persons, characterized by an early onset depression and a high familial loading for depressive disorders, for research into the pathophysiology and genetic background of mood disorders may facilitate the identification of etiological processes and genetic vulnerability.

## **EPILOGUE**

Knowledge gained by this thesis resulted in a critical appraisal of current depressive categories of MDD, Dysthymic Disorder and Double Depression. We demonstrated the possible benefits of dimensional approaches, with two cross-cutting dimensions of severity and duration of depressive symptoms, a dimension of anxiety, and additional features, such as functioning, personality measures and childhood adversity, above stringent DSM-categories. Findings provided the 'bricks' for building alternative models for depression, like the clinical staging framework. Psychiatry should move beyond our customary clinical taxonomy, and develop alternative models to differentiate between patients, who otherwise seem to be deceptively similar, since they share the same psychiatric diagnosis (Fava et al., 2012). Much is still there to be learned about the nature of depression, paraphrasing Father Sebastian Englert of Rapa Nui, Easter Island: "When the subject is depression, no man's knowledge is either complete or secure".

## REFERENCES

- Angst J, Gamma A, Benazzi F, Ajdacic V, Rossler W. Melancholia and atypical depression in the Zurich study: epidemiology, clinical characteristics, course, comorbidity and personality. *Acta Psychiatrica Scandinavica* 2007; 155: 72-84.
- Anisman H, Ravindran AV, Griffiths J, Merali Z. Endocrine and cytokine correlates of major depression and dysthymia with typical or atypical features. *Molecular Psychiatry* 1999; 4: 182-188.
- Aydemir O, Deveci A, Taskin OE, Taneli F, Esen-Danaci A. Serum brain-derived neurotrophic factor level in dysthymia: a comparative study with major depressive disorder. *Progress in Neuropsychopharmacology Biological Psychiatry* 2007; 31: 1023-1026.
- Bhagwagar Z, Hafizi S, Cowen PJ. Increased salivary cortisol after waking in depression. *Psychopharmacology (Berl)* 2005; 182: 54-57.
- Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: results of The Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Social Psychiatry and Psychiatric Epidemiology* 1998; 33: 587-595.
- Brietzke E, Mansur RB, Soczynska JK, Kapczinski F, Bressan RA, McIntyre RS. Towards a multifactorial approach for prediction of bipolar disorder in at risk populations, *Journal of Affective Disorders* 2012; doi:10.1016/j.jad.2012.02.016.
- Brodsky H, Luscombe G, Parker G, Wilhelm K, Hickie I, Austin MP, Mitchell P. Increased rate of psychosis and psychomotor change in depression with age. *Psychological Medicine* 1997; 27: 1205-1213.
- Brouwer JP, Appelhof BC, Hoogendijk WJ, Huyser J, Endert E, Zuckerman C, Schene AH, Tijssen JG, Van Dyck R, Wiersinga WM, Fliers E. Thyroid and adrenal axis in major depression: a controlled study in outpatients. *European Journal of Endocrinology* 2005; 152: 185-191.
- Buist-Bouwman MA, Graaf R de, Vollebergh WA, Alonso J, Bruffaerts R, Ormel J. Functional disability of mental disorders and comparison with physical disorders: a study among the general population of six European countries. *Acta Psychiatrica Scandinavica* 2006; 113: 492-500.
- Carroll BJ. Diagnostic validity and laboratory studies: rules of the game. In: *The Validity of Psychiatry Diagnosis* (eds. Robins LN and Barrett JA) 1989; pp. 229-245. APA Series, Raven Press: New York.
- Clark LA, Watson D. Tripartite model of anxiety and depression, psychometric evidence and taxonomic implications. *Journal of Abnormal Psychology* 1991; 100: 316-336.
- Coryell W, Scheftner W, Keller M, Endicott J, Maser J, Klerman GL. The enduring psychosocial consequences of mania and depression. *American Journal of Psychiatry* 1993; 150: 720-727.
- Coryell W, Winokur G, Shea T, Maser JD, Endicott J, Akiskal HS. The long-term stability of depressive subtypes. *American Journal of Psychiatry* 1994; 151: 199-204.
- Dowlati Y, Hermann N, Swardfager W, Liu H, Sham L, Reim EK, Lanctôt KL. A meta-analysis of cytokines in major depression. *Biological Psychiatry* 2010; 67: 446-457.
- Faravelli C, Ambonetti A, Pallanti S, Pazzagli A. Depressive relapses and incomplete recovery from index episode. *American Journal of Psychiatry* 1986; 143: 888-891.
- Fava GA, Kellner R. Staging: a neglected dimension in psychiatric classification. *Acta Psychiatrica Scandinavica* 1993; 87: 225-230.
- Fava GA, Tossani E. Prodromal stage of major depression. *Early Intervention in Psychiatry* 2007; 1: 9-18.

Fava GA, Rafanelli C, Tomba E. The clinical process in psychiatry: a clinimetric approach. *Journal of Clinical Psychiatry* 2012; 73: 177-184.

Fournier JC, DeRubeis RJ, Hollon SD, Dimidjian S, Amsterdam JD, Shelton RC, Fawcett J. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *Journal of the American Medical Association* 2010; 303: 47-53.

Goldberg D. A dimensional model for common mental disorders. *British Journal of Psychiatry* 1996; Suppl 44-49.

Graaf R de, Bijl RV, Smit F, Ravelli A, Vollebergh WA. Psychiatric and sociodemographic predictors of attrition in a longitudinal study: The Netherlands Mental Health Survey and Incidence Study (NEMESIS). *American Journal of Epidemiology* 2000; 152: 1039-1047.

Graaf R de, Tuithof M, Dorsselaer S van, Have M ten. Comparing the effects on work performance of mental and physical disorders. *Social Psychiatry and Psychiatric Epidemiology*. In press.

Griffiths J, Ravindran AV, Merali Z, Anisman H. Dysthymia: a review of pharmacological and behavioral factors. *Molecular Psychiatry* 2000; 5: 242-61.

Hetrick SE, Parker AG, Hickie IB, Purcell R, Yung AR, McGorry PD. Early identification and intervention in depressive disorders: towards a clinical staging model. *Psychotherapy and Psychosomatics* 2008; 77: 263-270.

Judd LL, Paulus MP, Wells KB, Rapaport MH. Socioeconomic burden of subsyndromal depressive symptoms and major depression in a sample of the general population. *American Journal of Psychiatry* 1996; 153: 1411-1417.

Judd LL, Akiskal HS. Delineating the longitudinal structure of depressive illness: beyond clinical subtypes and duration thresholds. *Pharmacopsychiatry* 2000; 33: 3-7.

Judd LL, Akiskal HS, Zeller PJ, Paulus M, Leon AC, Maser JD, Endicott J, Coryell W, Kunovac JL, Mueller TI, Rice JP, Keller MB. Psychosocial disability during the long-term course of unipolar major depressive disorder. *Archives of General Psychiatry* 2000; 7: 375-380.

Karsten J, Penninx BWJH, Riese H, Ormel H, Nolen WA, Hartman CA. The state effect of depressive and anxiety disorders on big five personality traits. *Journal of Psychiatric Research* 2012; 46: 644-650.

Keitner GI, Ryan CE, Miller IW, Norman WH. Recovery and major depression: factors associated with twelve-month outcome. *American Journal of Psychiatry* 1992; 149: 93-99.

Kendler KS, Fiske A, Gardner CO, Gatz M. Delineation of two genetic pathways to major depression. *Biological Psychiatry* 2009; 65: 808-811.

Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry* 1994; 51: 8-19.

Kessler RC, Stang PE, Wittchen HU, Ustun TB, Roy-Burne PP, Walters EE. Lifetime panic-depression comorbidity in the National Comorbidity Survey. *Archives of General Psychiatry* 1998; 55: 801-808.

Kessler RC, Ustun TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *International Journal of Methods in Psychiatric Research* 2004; 13: 93-121.

Kraepelin, E. *Psychiatrie. Ein Lehrbuch für Studierende und Aerzte. Vol II.* 1899. Leipzig: Verlag von Johann Ambrosius Barth.

Krueger RF. The structure of common mental disorders. *Archives of General Psychiatry* 1999; 56: 921-926.

Lamers F, de Jonge P, Nolen WA, Smit JH, Zitman FG, Beekman AT, Penninx BW. Identifying depressive subtypes in a large cohort study: results from the Netherlands Study of Depression and Anxiety (NESDA). *Journal of Clinical Psychiatry* 2010; 71: 1582-1589.

Lamers F, Hoogendoorn A, Smit J, van Dyck R, Zitman FG, Nolen WA, Penninx BW. Sociodemographic and psychiatric determinants of attrition in the Netherlands Study of Depression and Anxiety (NESDA). *Comprehensive Psychiatry* 2012; 53: 63-70.

Lecrubier Y, Sheehan D, Weiller E, Amorim P, Bonora I, Sheehan K, Janavs J, Dunbar G. The MINI International Neuropsychiatric Interview (M.I.N.I.) A Short Diagnostic Structured Interview: Reliability and Validity According to the CIDI. *European Psychiatry* 1997; 12: 224-231.

Lyons MJ, Eisen SA, Goldberg J, True W, Lin N, Meyer JM, Toomey R, Faraone SV, Merla-Ramos M, Tsuang MT. A registry-based twin study of depression in men. *Archives of General Psychiatry* 1998; 55: 468-472.

Lyketsois C, Nestadt G, Cwi J, Heithoff K, Eaton WW. The life-chart interview: a standardized method to describe the course of psychopathology. *International Journal of Methods in Psychiatric Research* 1994; 4: 143-155.

Marques-Deak AH, Neto FL, Dominguez WV, Solis AC, Kurciant D, Sato F, Ross JM, Prado EB. Cytokine profiles in women with different subtypes of major depressive disorder. *Journal of Psychiatric Research* 2007; 41: 152-159.

McGorry PD. Issues for DSM-V: clinical staging: a heuristic pathway to valid nosology and safer, more effective treatment in psychiatry. *American Journal of Psychiatry* 2007; 164: 859-860.

Merikangas KR, Ames M, Cui L, Stang PE, Ustun TB, Korff M von, Kessler RC. The impact of comorbidity of mental and physical conditions on role disability in the US adult household population. *Archives of General Psychiatry* 2007; 64: 1180-1188.

Molendijk ML, Bus BAA, Spinhoven Ph, Penninx BBWJH, Kenis G, Prickaerts J, Oude Voshaar RC, Elzinga BM. Serum levels of brain-derived neurotrophic factor in major depressive disorder: state-trait issues, clinical features and pharmacological treatment. *Molecular Psychiatry* 2011; 16: 1088-1095.

Morey LC, Shea MT, Markowitz JC, Stout RL, Hopwood CJ, Gunderson JG, Grilo CM, McGlashan TH, Yen S, Sanislow CA, Skodol AE. State effects of major depression on the assessment of personality and personality disorder. *American Journal of Psychiatry* 2010; 167: 528-535.

Moulds ML, Kandris E, Williams AD, Lang T, Yap C, Hoffmeister K. An investigation of the relationship between cognitive reactivity and rumination. *Behavioral Therapy* 2008; 39: 65-71.

Nelson JC, Davis JM. DST studies in psychotic depression: a meta-analysis. *American Journal of Psychiatry* 1997; 154: 1497-1503.

Nolen WA, Kupka RW, Schulte PFJ, Knoppert-van der Klein EAM, Honig A, Reichart CG, Goossens PJJ, Daemen P, Ravelli DP (Richtlijncommissie bipolaire stoornissen van de Commissie Kwaliteitszorg van de Nederlandse Vereniging voor Psychiatrie) Richtlijn Bipolaire Stoornissen- Tweede, herziene versie. 2008. De Tijdstroom, Utrecht.

Oldehinkel AJ, van den Berg MD, Flentge F, Bouhuys AL, ter Horst GJ, Ormel J. Urinary free cortisol excretion in elderly persons with minor and major depression. *Psychiatry Research* 2001; 104: 39-47.

Olino TM, Klein DN, Farmer RF, Seeley JR, Lewinsohn PM. Examination of the structure of psychopathology using latent class analysis. *Comprehensive Psychiatry* 2012; 53: 323-332.

Ormel J, Oldehinkel T, Brilman E, Van den Brink W. Outcome of depression and anxiety in primary care. A three-wave study of psychopathology and disability. *Archives of General Psychiatry* 1993; 50: 759-766.

Ormel J, Von Korff M, Ustun TB, Pini S, Korten A, Oldehinkel T. Common mental disorders and disability across cultures. Results from the WHO Collaborative Study on Psychological Problems in General Health Care. *Journal of the American Medical Association* 1994; 272: 1741-1748.

Ormel J, Rosmalen J, Farmer A. Neuroticism: a non-informative marker of vulnerability to psychopathology. *Social Psychiatry and Psychiatric Epidemiology* 2004; 39: 906-912.

Overall JE, Hollister LE, Johnson M, Pennington V. Nosology of depression and differential response to drugs. *Journal of the American Medical Association* 1966; 195: 946-948.

Parker G, Roy K, Hadzi-Pavlovic D, Wilhelm K, Mitchell P. The differential impact of age on the phenomenology of melancholia. *Psychological Medicine* 2001; 31: 1231-1236.

Parker G, Fink M, Shorter E, Taylor MA, Akiskal H, Berrios G, Bolwig T, Brown WA, Carroll B, Healy D, Klein DF, Koukopoulos A, Michels R, Paris J, Rubin RT, Spitzer R, Swartz C. Issues for DSM-5: whether melancholia? The case for its classification as a distinct mood disorder. *American Journal of Psychiatry* 2010; 167:745-747.

Perry A, Tarrrier N, Morriss R, McCarthy E, Limb K. Randomised controlled trial of efficacy of teaching patients with bipolar disorder to identify early symptoms of relapse and obtain treatment. *British Medical Journal* 1999; 318: 149-153.

Paykel ES. Depressive typologies and response to amitriptyline. *British Journal of Psychiatry* 1972; 120: 147-156.

Penninx BW, Nolen WA, Lamers F, Zitman FG, Smit JH, Spinhoven P, Cuijpers P, de Jong PJ, van Marwijk HW, van der Meer K, Verhaak P, Laurant MG, de Graaf R, Hoogendijk WJ, van der Wee N, Ormel J, van Dyck R, Beekman AT. Two-year course of depressive and anxiety disorders: results from the Netherlands Study of Depression and Anxiety (NESDA). *Journal of Affective Disorders* 2011; 133: 76-85.

Ramsawh HJ, Raffa SD, Edelen MO, Rende R, Keller MB. Anxiety in middle adulthood: effects of age and time on the 14-year course of panic disorder, social phobia and generalized anxiety disorder. *Psychological Medicine* 2009; 39: 615-624.

Rapaport MH, Clary C, Fayyad R, Endicott J. Quality-of-life impairment in depressive and anxiety disorders. *American Journal of Psychiatry* 2005; 162: 1171-1178.

Regier DA, Kaelber CT, Rae DS, Farmer ME, Knauper B, Kessler RC, Norquist GS. Limitations of diagnostic criteria and assessment instruments for mental disorders. *Archives of General Psychiatry* 1998; 47: 519-526.

Sargeant JK, Bruce ML, Florio LP, Weissman MM. Factors associated with 1-year outcome of major depression in the community. *Archives of General Psychiatry* 1990; 47: 519-526.

Seedat S, Scott KM, Angermeyer MC, Berglund P, Bromet EJ, Brugha TS, Demyttenaere K, de Girolamo G, Haro JM, Jin R, Karam EG, Kovess-Masfety V, Levinson D, Medina Mora ME, Ono Y, Ormel J, Pennell BE, Posada-Villa J, Sampson NA, Williams D, Kessler RC. Cross-national associations between gender and mental disorders in the World Health Organization World Mental Health Surveys. *Archives General Psychiatry* 2009; 66: 785-795.

Shankman SA, Klein DN. Dimensional diagnosis of depression: Adding the dimension of course to severity, and comparison to the DSM. *Comprehensive Psychiatry* 2002; 43: 420-426.

Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* 1998; 59 Suppl 20: 22-33.

Solomon DA, Leon AC, Endicott J, Mueller TI, Coryell W, Shea MT, Keller MB. Psychosocial impairment and recurrence of major depression. *Comprehensive Psychiatry* 2004; 45: 423-430.

Spijker J, de Graaf R, Bijl RV, Beekman ATF, Ormel J, Nolen WA. Duration of major depressive episodes in the general population: results from The Netherlands Mental Health Survey and Incidence Study (NEMESIS). *British Journal of Psychiatry* 2002; 181: 208-213.

Spijker J, Graaf R de, Bijl RV, Beekman AT, Ormel J, Nolen WA. Functional disability and depression in the general population. Results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Acta Psychiatrica Scandinavica* 2004; 110: 208-214.

Vogelzangs N, Duivis HE, Beekman ATF, Kluit C, Neuteboom J, Hoogendijk W, Smit JH, de Jonge P and Penninx BWJH. Association of depressive disorders, depression characteristics and antidepressant medication with inflammation. *Translational Psychiatry* 2012; 2, e79; doi:10.1038/tp.2012.8.

Vollebergh WAM, Iedema J, Bijl RV, De Graaf R, Smit F, Ormel J. The structure and stability of common mental disorders: the NEMESIS Study. *Archives of General Psychiatry* 2001; 58: 597-603.

Vreeburg SA, Hoogendijk WJG, van Pelt J, de Rijk RH, Verhagen JCM, van Dyck R, Smit JH, Zitman FG, Penninx BWJH. Major Depressive Disorder and hypothalamic-pituitary-adrenal axis activity: results from a large cohort study. *Archives of General Psychiatry* 2009; 66: 617-626.

Vuorilehto MS, Melartin TK, Isometsa ET. Course and outcome of depressive disorders in primary care: a prospective 18-month study. *Psychological Medicine* 2009; 39: 1697-1707.

Wardenaar KJ, Giltay EJ, van Veen T, Zitman FG, Penninx BW. Symptom dimensions as predictors of the two-year course of depressive and anxiety disorders. *Journal of Affective Disorders* 2012; 136: 1198-1203.

Ware JE, Kosinski M, Bayliss MS, McHorney CA, Rogers WH, Raczek A. Comparison of methods for the scoring and statistical analysis of SF-36 health profile and summary measures: summary of results from the Medical Outcomes Study. *Medical Care* 1995; 33(4 Suppl): AS264-AS279.

Watson S, Gallagher P, Del-Estal D, Hearn A, Ferrier IN, Young AH. Hypothalamic-pituitary- adrenal axis function in patients with chronic depression. *Psychological Medicine* 2002; 32: 1021-1028.

Wells KB, Burnam MA, Rogers W, Hays R, Camp P. The course of depression in adult outpatients. Results from the Medical Outcomes Study. *Archives of General Psychiatry* 1992; 49: 788-794.

Wiersma D, De Jong A, Ormel J. The Groningen Social Disabilities Schedule: development, relationship with I.C.I.D.H., and psychometric properties. *International Journal of Rehabilitation Research* 1988; 11: 213-224.

Wiersma JE, Hovens JGFM, Van Oppen P, Giltay EJ, Van Schaik DJF, Beekman ATF, Penninx BWJH. The importance of childhood trauma and childhood life events for chronicity of depression in adults. *Journal of Clinical Psychiatry* 2009; 70: 983-989.

Wiersma JE, van Oppen P, van Schaik DJ, van der Does AJ, Beekman AT, Penninx BW. Psychological characteristics of chronic depression: a longitudinal cohort study. *Journal of Clinical Psychiatry* 2011; 72: 288-294.

Wittchen HU, Robins LN, Cottler LB, Sartorius N, Burke JD, Regier D. Cross-cultural feasibility, reliability and sources of variance of the Composite International Diagnostic Interview (CIDI). The Multicentre WHO/ADAMHA Field Trials. *British Journal of Psychiatry* 1991; 159: 645-653, 658.

Wittchen HU. Reliability and validity studies of the WHO-Composite International Diagnostic Interview (CIDI): a critical review. *Journal of Psychiatric Research* 1994; 28: 57-84.

World Health Organization. Composite Diagnostic Interview (CIDI), version 1.0. 1990. Geneva, World Health Organization.

World Health Organization. Composite International Diagnostic Interview (CIDI), version 2.1. 1998. Geneva, World Health Organization.

Yonkers KA, Bruce SE, Dyck IR, Keller MB. Chronicity, relapse, and illness-course of panic disorder, social phobia, and generalized anxiety disorder: findings in men and women from 8 years of follow-up. *Depression and Anxiety* 2003; 17: 173-179.

Young MA, Keller MB, Lavori PW, Scheftner WA, Fawcett JA, Endicott J, Hirschfeld RM. Lack of stability of the RDC endogenous subtype in consecutive episodes of major depression. *Journal of Affective Disorders* 1987; 12: 139-143.





# Nederlandse Samenvatting

---



Depressies, ofwel 'depressieve stoornissen', zijn veel voorkomende ziektebeelden met een grote impact op het leven van patiënten en op de samenleving als geheel. De Wereldgezondheidsorganisatie (WHO) verwacht dat in 2030 depressie op de eerste plaats staat van ziekten met de hoogste ziektelast. Twintig procent van de Nederlandse bevolking krijgt in zijn of haar leven te maken met een stemmingsstoornis en per jaar lijdt ongeveer 6% van de volwassenen aan een stemmingsstoornis. Dit zijn enorme aantallen. Omgerekend betekent het dat rond 650 duizend volwassenen in Nederland in 2011 last hadden van een stemmingsstoornis. Inzicht in depressies is dan ook van groot belang.

Sinds 1980 wordt in de psychiatrie wereldwijd gebruik gemaakt van het 'Diagnostic and Statistical Manual' (DSM), een classificatiesysteem om psychiatrische ziektebeelden te ordenen. Dit classificatiesysteem heeft onder meer onderzoek naar het ontstaan, beloop en behandeling van depressies gefaciliteerd. De huidige versie van de DSM maakt onderscheid tussen een depressieve stoornis, gedefinieerd als een depressieve stemming of interesseverlies gedurende tenminste twee weken; een dysthyme stoornis, gedefinieerd als een depressieve stemming, milder van aard dan de depressieve stoornis, maar van langere duur (tenminste twee jaar); en een depressie 'niet anderszins omschreven'. In dit proefschrift onderzoeken wij de heterogeniteit van depressies. In het bijzonder verkennen wij de concepten 'depressieve stoornis' en 'dysthyme stoornis'.

Epidemiologische studies tonen aan dat een depressieve stoornis en dysthyme stoornis vaak tegelijkertijd voorkomen- in sommige onderzoeken heeft zelfs meer dan 90% van de patiënten met een dysthyme stoornis tevens een depressieve stoornis. Dit gelijktijdig voorkomen van beide stoornissen wordt 'dubbele depressie' genoemd. Tevens blijken de (chronische) depressieve stoornis en de dysthyme stoornis nauwelijks van elkaar te verschillen in socio-demografische en klinische kenmerken. Dit voedt de twijfel over de validiteit van beide concepten. Met het oog op de DSM-V, de nieuwe versie van de DSM die naar verwachting in mei 2013 uit zal komen, zijn alternatieve classificaties voor depressies voorgesteld. Hierin worden beide stoornissen samengevoegd en wordt er slechts een onderscheid gemaakt tussen een chronische en niet-chronische depressieve stoornis. De dysthyme stoornis zou dan uit de DSM verdwijnen.

Niet alleen wordt het onderscheid tussen de depressieve en de dysthyme stoornis betwist, ook de afgrenzing tussen depressies en angststoornissen is onderwerp van discussie. Al sinds de oude Grieken wordt erkend dat depressies vaak gepaard gaan met angstklachten. Desondanks worden sinds de DSM-III (1980) angst en depressie apart geclassificeerd. Een grote mate van comorbiditeit- het gezamenlijk voorkomen- van angst en depressie leidt echter tot de vraag of het huidige onderscheid tussen angst en depressie wel gerechtvaardigd is.

Gezien de hoge prevalentie van depressies in de bevolking en de grote ziektelast, is het van groot belang dat er helderheid wordt verkregen in de heterogeniteit van depressieve stoornissen. In dit proefschrift trachten wij dan ook i) inzicht te verkrijgen in de heterogeniteit binnen het grote containerbegrip ‘depressie’ en dan in het bijzonder in het onderscheid tussen de depressieve stoornis, de dysthyme stoornis en de combinatie van beide, de zogenaamde ‘dubbele depressie’; ii) alternatieve modellen voor het indelen van depressieve stoornissen te verkennen, en iii) het belang van angst(stoornissen) op de classificatie van depressieve stoornissen te onderzoeken.

### **Deel I: Classificatie van depressieve stoornissen**

De eerste vier hoofdstukken van dit proefschrift onderzoeken de huidige onderverdeling van depressies in de depressieve stoornis, de dysthyme stoornis en de ‘dubbele depressie’. In *hoofdstuk 2* wordt de vraag gesteld of er verschillen zijn in biologische en psychologische kenmerken van mensen- deelnemers aan de Nederlandse Studie naar Depressie en Angst (NESDA)- met een depressieve stoornis, dysthyme stoornis of ‘dubbele depressie’. Er werden geen verschillen gevonden in biologische kenmerken, bestaande uit cortisol, ontstekingsparameters en serum Brain Derived Neurotrophic Factor (sBDNF). Personen met een dysthyme stoornis en ‘dubbele depressie’ hebben wel een ander psychologisch profiel. Zij zijn onder andere neurotischer en minder consciëntieus dan personen met een depressieve stoornis. Dit doet vermoeden dat chroniciteit van depressieve symptomen geassocieerd is met dergelijke persoonlijkheidskenmerken.

Het beloop van een ziekte verschaft veel informatie over een ziektebeeld. In *hoofdstuk 3* en *4* wordt dan ook respectievelijk i) het 3-jaars beloop van depressies en ii) het 3-jaars beloop van het dagelijks functioneren van mensen met een doorgemaakte depressie beschreven. Beide hoofdstukken zijn gebaseerd op de Netherlands Mental Health Survey and Incidence Study (NEMESIS). Opnieuw werden drie groepen- depressieve stoornis, dysthyme stoornis en ‘dubbele depressie’- met elkaar vergeleken. Beide hoofdstukken tonen aan dat er welliswaar een verschil is tussen de depressieve stoornis en de dysthyme stoornis of ‘dubbele depressie’, maar dat de beide chronische vormen (de dysthyme stoornis en de ‘dubbele depressie’) nauwelijks van elkaar te onderscheiden zijn. Een onderverdeling in niet-chronisch versus chronisch- zoals voorgesteld voor de DSM-V lijkt dan ook gerechtvaardigd.

Tenslotte wordt in *hoofdstuk 5* de ‘dysthyme stoornis’ nader onder de loep genomen. Met diverse methoden wordt getracht het concept te ontrafelen, vanuit de gedachte dat dit wellicht een conglomeraat is van verschillende depressieve subtypes. 318 Depressieve patiënten, verwezen naar het ‘Black Dog Institute Depression Clinic’ in Sydney, Australië, werden geselecteerd. Statistische methoden (Latente Klasse Analyse en Latente Profiel Analyse), die zonder vooronderstellingen kijken of er bepaalde patronen of klassen in de data voorkomen, werden toegepast. Verschillende klassen werden geïdentificeerd. Echter, deze klassen bleken dermate verschillend, dat er geen eenduidige subtypes geïdentificeerd

konden worden, behoudens subtypen verschillend in ernst van de depressieve symptomen. Tenslotte werd onderzocht welke psychiatrische diagnoses psychiaters stelden bij een subgroep patiënten, die voldeden aan de criteria van een pure 'dysthyme stoornis' (n=42). Geen van de psychiaters stelde de diagnose 'dysthyme stoornis'. Deze resultaten bevragen de discriminante validiteit en klinische utiliteit van het concept dysthyme stoornis.

In dit eerste deel van het proefschrift baseren wij ons op DSM-definities. Het is echter zeer wel mogelijk dat onze resultaten tot dusver beïnvloed werden door de gehanteerde DSM-definities. Immers, indien er geen verschillen worden gevonden tussen de verschillende depressie-groepen, kan dit wijzen op ofwel een afwezigheid van diversiteit binnen de depressieve stoornissen, ofwel op een foutieve definiëring van de depressie-groepen. Werkelijke verschillen tussen de groepen worden dan niet gevonden - een gebrek aan validiteit van de gehanteerde diagnoses. Om dit nader te onderzoeken worden in deel 2 van het proefschrift alternatieve onderzoeksmethoden toegepast waarmee wordt gezocht naar een andere indeling voor depressies dan de huidige DSM-classificatie.

## **Deel II. Alternatieve classificatiemodellen voor depressies**

In het tweede deel van dit proefschrift hanteren we niet de DSM-classificatie om de heterogeniteit van depressieve stoornissen te onderzoeken, maar laten we de data spreken. *Hoofdstuk 6* toont de resultaten van een 'Latent Class Growth Analysis' (LCGA)- een extensie van de traditionele Latente Klasse Analyse, waarmee latente beloopstypen geïdentificeerd kunnen worden. Met behulp van het Life Chart Interview (LCI) werden gedurende twee jaar depressieve symptomen geregistreerd van een groep depressieve patiënten (n=804), die participeerden in NESDA. De LCI is een methode om het beloop van depressieve of angst klachten in kaart te brengen. Personen plaatsen allereerst belangrijke levensgebeurtenissen op een kalender. Vervolgens wordt gekeken of er in de afgelopen jaren sprake was van angst of depressieve klachten. De levensgebeurtenissen helpen om het geheugen op te frissen en daarmee tot een goede registratie van angst of depressieve klachten te komen. De LCGA identificeerde vijf verschillende beloopstypen van depressieve klachten. De beloopstypen varieerden in ernst en duur van de symptomen. Het was opmerkelijk, dat meer dan 50% van de personen met een dysthyme stoornis of 'dubbele depressie' bleken te behoren tot de gunstige beloopstypen, terwijl deze depressies in de DSM gedefinieerd worden als chronische depressies van tenminste twee jaar of langer. Dit doet vermoeden dat de DSM-diagnoses niet goed in staat zijn het beloop te voorspellen. Nader onderzoek van de verschillende beloopstypen liet zien, dat i) oudere leeftijd, ii) een op jongere leeftijd ontstaan van depressies en iii) minder extraversie voorspellend waren voor een slechter beloop.

*Hoofdstuk 7* presenteert de resultaten van een tweede data-gestuurde methode, de Latente Transitie Analyse (LTA). Hier werd gekeken naar de stabiliteit van depressieve subgroepen, geïdentificeerd naar ernst en aard- vitale kenmerken versus atypische kenmerken- van de

depressieve symptomen. Vitale kenmerken zijn onder meer vroeg ontwaken en afname van eetlust en gewichtsverlies. Atypische kenmerken zijn onder meer toegenomen slaapbehoefte en gewichtstoename. De LTA toonde aan dat de subtypes stabiel waren gedurende 2 jaar follow-up, hetgeen een indicatie is dat classificatie naar ernst en kenmerken van de depressieve symptomen (vitale kenmerken versus atypische kenmerken) een klinisch relevant onderscheid markeren.

### **Deel III. Depressie en angst, of angstige depressie?**

Het laatste deel van het proefschrift verkent de rol van angst bij depressies. In *hoofdstuk 8* worden 303 mensen, met een depressie en/of angststoornis, participierend in zowel NEMESIS als NESDA, gedurende 7 jaar gevolgd. Na 7 jaar bleek ruim 60% geen DSM-diagnose van een depressie of angststoornis meer te hebben. Het risico op het persisteren van de stoornis bleek vergroot bij mensen met een angst stoornis- al dan niet met een comorbide depressie- bij aanvang van het onderzoek, bij slechter lichamelijk functioneren en bij meer neuroticisme. Tevens werd gekeken naar het persisteren van depressieve en angstklachten, geregistreerd middels het 'Life Chart Interview' in plaats van DSM-diagnoses. Ook hier bleek dat (comorbide) angst een slechter beloop voorspelde. Een grote mate van neuroticisme en traumatische ervaringen in de jeugd waren geassocieerd met meer symptomen gedurende follow-up.

In DSM-V wordt de rol van angst op het ontstaan en beloop van diverse psychiatrische aandoeningen erkend. Het voorstel is 'paniek' als een overkoepelende dimensie bij elke vorm van psychopathologie te scoren. In *hoofdstuk 9* onderzoeken wij de rol van paniek op het ontstaan en persisteren van psychopathologie en op het dagelijks functioneren. Aangezien wij in dit proefschrift de heterogeniteit van depressies onderzoeken, bespreken we hier alleen het effect van paniek op depressies. Zoals verwacht bleek paniek een aanzienlijke invloed te hebben op het ontstaan en beloop van depressieve stoornissen en het dagelijks functioneren van mensen met een depressie. Dit gold voor huidige paniek aanvallen, maar ook- zij het in mindere mate- voor een voorgeschiedenis van paniekaanvallen, zonder huidige klachten.

Tenslotte presenteert *hoofdstuk 10* de resultaten van een Latente Klasse Analyse, waarin gekeken werd of mensen met een gegeneraliseerde angststoornis en een dysthyme stoornis verschillen in hun symptoomprofielen. Als beide stoornissen voldoende verschillen pleit dit voor een onderscheid in twee stoornissen in de DSM. Het blijkt uit deze studie dat de symptomen grotendeels overlappen en er geen sprake is van een duidelijk 'angst' profiel versus een 'depressief' profiel. Veeleer lijkt er sprake van 'sombere piekeraars' ofwel 'piekerende depressieven'. Deel 3 van dit proefschrift toont aan dat angst een grote rol speelt in het ontstaan en beloop van depressie en dat het dikwijls moeilijk is angst van depressie te onderscheiden. In de classificatie van depressie dient angst dan ook een evidente rol te krijgen.

Dit proefschrift eindigt met een algemene discussie (*hoofdstuk 11*) van de belangrijkste bevindingen en methodologische overwegingen. Samenvattend pleiten de resultaten voor een aanpassing van de huidige DSM classificatie voor depressieve stoornissen. Het onderscheid van depressies in een depressieve stoornis versus dysthyme stoornis lijkt niet gerechtvaardigd. Daarentegen lijken de parameters 'ernst' en 'duur' van de depressieve klachten wel onderscheidend te zijn voor depressieve subtypes. Tevens dient angst een meer prominente rol in de classificatie van depressies te krijgen. Hoofdstuk 11 presenteert een alternatief voor het huidige classificatiemodel geïnspireerd door stageringsmodellen, zoals reeds toegepast in onder meer de oncologie. Bij longkanker bijvoorbeeld wordt gekeken naar de grootte van de tumor, de mate van uitzaaiingen in lymfklieren en/of uitzaaiingen elders in het lichaam. Op grond hiervan wordt het stadium van de longkanker, en daarmee de behandeling en prognose, bepaald. Voor depressies kan een dergelijk stageringsmodel, met stadia gekenmerkt door verschillende mate van ernst en duur van depressieve klachten en angst, leidraad zijn voor klinisch handelen. Verdere identificatie van predictoren voor slechter beloop, zoals oudere leeftijd, jongere leeftijd van ontstaan van klachten, persoonlijkheidskenmerken (zoals neuroticisme), trauma en slechter functioneren allen geïdentificeerd in dit proefschrift als voorspellers voor slechter beloop- en mogelijk biologische of genetische parameters, dienen in het model geïncorporeerd te worden. Zo zou men tot snellere identificatie van personen met een groot risico voor een slechter beloop kunnen komen. Dit zou snel en doelgericht ingrijpen kunnen faciliteren en daarmee leiden tot preventie van het ontstaan of chronisch worden van depressieve stoornissen. Mogelijk zijn, zoals in de oncologie, verschillende stageringsmodellen nodig voor bijvoorbeeld melancholische depressie, of voor bepaalde depressies op oudere leeftijd. In hoeverre slechts één stageringsmodel voldoende blijkt voor het grote container begrip 'depressie' dient dan ook nader onderzocht te worden.





Dankwoord

---



En dan het dankwoord. Dit proefschrift was niet tot stand gekomen dankzij... Meer dan waar! Maar: het voelt persoonlijker dan deze standaard zin. Mijn weg door onderzoeksland is niet zonder pieken of dalen geweest. Jonglerend met de tijd tussen kliniek en onderzoek. En dan: balans vinden tussen moederschap en deadlines. Het is niet alleen een wetenschappelijke uitdaging geweest, maar tevens een ontdekkingstocht in wat ik wil. Daar heb je steun bij nodig. Praktische steun (datamanagement en automatisering: bedankt!), inhoudelijke steun en een sterk wetenschappelijk kader (NEMESIS en NESDA-consortium: bedankt!), en een sterke dataset (alle respondenten: dank!), maar bovenal persoonlijke steun.

Allereerst mijn promotoren en co-promotor, Brenda, Aartjan, Witte en Ron. Zonder jullie was dit proefschrift er niet geweest. Brenda, je bent een briljant onderzoekster, scherp analyticus, keiharde werkster met toch altijd tijd voor feedback, en iemand die oprecht meedenkt. Wat heb ik veel van je geleerd. Ik hoop dat ik de komende jaren mag genieten van de samenwerking met jou. Aartjan, je was er tijdens verschillende 'stadia' van mijn loopbaan: de opleiding tot psychiater, het onderzoek en nu tijdens de transitie naar post-doc. Bereikbaar wanneer ik je nodig had. Niet alleen voor wetenschap inhoudelijke vragen, maar ook voor verheldering van keuzes en als inspirator. Als opleider enthousiasmeerde je mij voor promotie-onderzoek, als wetenschapper (en clinicus) enthousiasmeerde je mij voor je idee van staging en profiling, zichtbaar in de general discussion van dit proefschrift. Jouw invloed gedurende elke fase toont aan, dat sommige factoren (profilers) in elk stadium van belang zijn! Witte, wat meer op een afstand zag je mijn vorderingen van begin tot einde. Ik hoop in de toekomst nog te mogen putten uit je grote kennis van de biologische psychiatrie. Ron, geweldige co-promotor en voorbeeld voor bewust keuzes maken. Altijd snel en precies in je commentaar op manuscripten, stimulerend bij dips en een fijne gesprekspartner. Ik bewonder je relativiseringsvermogen, en hoop nog nader onderzoek te kunnen doen naar onze stelling: 'Ten einde de impact van een sabbatical op het gemoed te minimaliseren, dient dit met enige regelmaat uitgevoerd te worden'.

Gordon: It was a great pleasure to work together, both in clinical care and research. I admire your passion for patients, your warmth, your excellent research and writing skills. Thanks for welcoming me in your institute and in your house; for sharing your thoughts with me. Be warned: I think I need a third visit to practice the art of both 'provocative' and poetic writing! Bec, thanks for your patience, guiding me through the datasets of the BDI. I really loved working with you, your humor, and your conscientiousness during the endless revisions of the paper (I counted 13 versions...). We'll keep in contact and hopefully, we'll finally have a 'diner' together ☺! Christine and Will, Karlyn and Pauline, all RAs, Matt, and all the others of the BDI: thanks for your great hospitality! You made my stay in Sydney unforgettable! Woof woof.

Verder uiteraard dank voor alle co-auteurs. Nicole: Veel te kort een kamer gedeeld. Ik geniet van je intelligentie, je betrokkenheid, je gezelligheid en onze beider link met tweelingen! Femke, het was een aangename gezamenlijke speurtocht naar de ins en outs van de LCGA. Wordt vervolgd! Matt and Gavin: having a cappuccino overlooking the Harbor Bridge and the Opera House, while brainstorming on dysthymia and GAD, is my definition of great research! I am looking forward to any replication study!

Wat is een ontdekkingstocht door onderzoeksland zonder mede-reizigers? Hoewel ik met regelmaat te druk was voor onze gezellige lunches met te dure kannen verse jus, heb ik de samenwerking en de kortstondige intervisie met jullie, mijn mede-promovendi, enorm gewaardeerd. Inger, Sophie, Carmilla, Jenneke en Adrie: bedankt! Hoewel nu verspreid over het brede landschap van onderzoek en kliniek, hoop ik dat we elkaar toch blijvend weten te vinden! Ook de overige promovendi en flex-kamergenoten: dank.

Ineke en Rianne: al vanaf de Lassus-tijd zijn jullie mijn rotsen in de branding. Marissa, Carla en Reen: jullie zijn onmisbare schakels! Tenslotte ook alle andere medewerkers van NESDA bedankt! Zonder publicaties, geen publicatie. Jopie en Marijke, jullie voorzagen mij van artikelen over de meest uiteenlopende onderwerpen, in papier of digitaal, vanaf de jaren 60 tot nu. Dank voor jullie speurwerk en jullie gezelligheid op de 4<sup>e</sup> etage. Lowijs. Hoe je het onderwijs hebt vormgegeven is uniek. Daarbij vergeet je niet dat er meer is en stapte je soms op mijn rem, voordat ik zover was. Dank voor je oprechte betrokkenheid. Willem, opleider aan het begin en stimulator tot het eind. Dank!

Beste Max, bevlogen en betrokken, groot clinicus, tot het uiterste je best doend een inspirerende en plezierige werkomgeving voor anderen te creëren. Mensen verbinden. Het lukt je grandioos! Veel dank voor je warmte, je openheid en de ruimte die je me geeft mijzelf te ontwikkelen. Ik hoop op nog vele jaren samenwerking in de wetenschap en in de kliniek.

Beste Piet: ik geloof dat jij eerder wist dat de wetenschap toch iets voor mij is dan ikzelf. Je hebt de gave om tegen de stroom in te denken, en mensen tot grotere hoogten te brengen. Hoezo glazen plafond? Het is een voorrecht met je samen te werken en je ideeën te horen. Gelukkig heb je niet altijd gelijk: het helemaal stoppen met werken is je gelukkig niet gelukt na je pensioen! Ik hoop dat ik nog lang van jouw originaliteit en genialiteit kan genieten (en ga nou niet tegensputteren...).

Jan Willem, veel dank voor de mogelijkheid om mijn wetenschappelijk kunnen verder te ontwikkelen down under. Hoewel de timing ongelukkig bleek, sta je voor je besluit, en heb je daardoor bijgedragen aan een prachtig internationaal netwerk (en aan de wens van Mare kangaroos eten te geven in plaats van geitjes...). Ik werk met veel plezier met je samen.

Audrey, niet alleen maar baas, maar met name ook koffie-en-klets-partner, Brokopondo-vaarder, oud-intervisie-genoot, en beiden trotse moeder van 2 meiden. Dank voor je vertrouwen in mij als onderzoekster, clinicus en manager. Tijd voor de speeltuin!

Team OO. Eric, al jaren intervisie-genoot, nu samen verantwoordelijk voor de afdeling. Dank voor de samenwerking. (Oud-) Arts-assistenten (Alies, Gerda, Mieke, Vivian, Sietske, Clara, Jojanneke, Sara en Sanne): fijn dat jullie op de toko pasten als ik een analyse zat te draaien! Team van Opname Ouderen: alle verpleegkundigen (ja ook Sigrid, Merle, Piet, Astrid, Ted, Harry, William en Cathy), Yvon, Margreet, Christine, Simone, Chris, Juliette, Marjan, Marilou, Michelle, Erik, Yuniel, Kitty, en de passanten (of inmiddels blijvers) Doreen, Anne, Derk, Annelies en Yvon: Heel fijn dat jullie mij de ruimte gaven onderzoek te doen of met mij onderzoek deden. Ook veel dank voor jullie oprechte meeleven als het tegen zat en jullie enthousiasme tijdens hoogtepunten. Anneke, jaren mijn duaal-partner. Tijdens mijn afwezigheid (niet zelden, en niet altijd kort), runde jij de afdeling. Samen hebben we veel meegemaakt, hier en in China. Veel dank voor je steun! Hoewel je nu op een andere locatie bent, we weten elkaar te vinden! Eric-Jan: net samen begonnen, nu al vertrouwd. We gaan er een goede tijd van maken.

Annemiek (maatje- en meer- op het plein: gauw weer een wijntje?!), Paul David, Caroline, Jeroen, Cora, Daniël, Annetje, Graziella, Vivian, Liesbeth, Irene, alle CMT-ers, Sigfried, Satish, Richard, Debbie, Cobie: jullie betrokkenheid is de kracht van het circuit ouderen! Hannie: je vond dat je niet in mijn dankwoord hoefde... fout gedacht! Heel fijn om op je terug te kunnen vallen en een weekend-planning door te nemen ☺. Mourien (en Margriet)- warm en vol interesse: ik geniet van onze 'omelet'-momenten.

Lieve vrienden, zonder jullie kon ik de klus niet klaren! Margreeth- mijn oudste vriendin. Inmiddels 27 jaar lief en leed. Laten we dit verveelvoudigen! Barbara- hartsvriendin! De kunst van het genieten is jou gegeven. Laten we dat samen blijven doen, voor altijd! Lindoor kinderdruckte en schrijfwerk helaas al jaren geen Pieterpad meer. Mijn schoenen staan weer klaar! Tim- veel margarita's, veel eitjes, veel sms-jes, veel jaren trouwe vriend. Nog lang niet genoeg! Michel en Daan- jullie hebben het vermogen mij in de grootste stress te laten ontspannen. Altijd een luisterend oor, verdiepend en fijne-soms eigenwijze-tips. Sjoerd- via hoge toppen in Nepal, over Friese wateren en in Franse tuinen. Ik geniet van ons samenzijn. Henri- Al jaren dierbare vriend. Heel fijn om mijn keuzes met jou te bevragen en te verhelderen (al is het dan in Heerenveen ☺). Frans - oudste vriend en kaas-fondue expert. Ik geniet van de ontelbare etentjes. Henri, Enno, Khalid, Elke, Khalil, Esther, Florine, Gertrude en Matt, Amba en Rudolf, Christel en Arnoud, Emmy, Marja, Judith, Maureen, Nikander en Leny, Chantal en Bram, Vincent en Sophie: jullie allen hielpen me mijn weg te kiezen, te relativeren en te genieten. Heel veel liefs! Ira, Howard, Ariane en Noara: jullie maakten Australië voor ons een nieuw thuis. Wanneer gaan we met een eski in the hute naar Straddie om alsnog die shrimps te BBQ-en? Marrit, Femke, Pieter en Sofie, en Martin,

Saskia, Fleur en Morris: down under is nog veel leuker als je er samen van kan genieten. Camilia, Alfred, Lemar and Nora: I think you invented 'hospitality'! We warmly remember the good times we had in Perth.

Lieve familie. Lieve Broer en Janke: dank voor de warme interesse (en het oppassen)! Het wordt weer tijd voor witjes! Lieve Welmoed, Reijnder, Jesse, Femke, Wietske, Roeland, Marloes, Minke en Sinne, Oer-Pake en Oer-Beppe. Betrokken en liefdevol. Dank. Phil en Nine, Flor, Emma en Damiaan, Doreth en Rob, Iwan, Annelies en Mirjam. Hoewel wat verder weg, dank voor het meelevens. Lieve Theo en Siets. Telkens weer wilden jullie weten hoe het met het onderzoek ging, maar bovenal hoe het met mij ging. Hoewel de materie soms ver van jullie bed was, waren jullie altijd geïnteresseerd. Lieve Diet en Tien. Als naamgenote trad ik in jullie voetsporen. Jullie waren er. Fijn me gesteund te voelen, in mijn beginjaren als arts-assistent, daarna als psychiater, en nu ook als onderzoeker, maar vooral als petekind. Lieve Bartje, ik hoop dat ik de komende tijd wat meer van jouw rust, ontspanning en bourgondische inslag weet te vinden. Fijn dat je mijn zwager bent!

Lieve papsje en mamsje. Hoor daar: de vogels fluiten. Kleine handjes, kleine voetjes. Jullie liefde is onvoorwaardelijk. Niet gebonden aan mooie publicaties of een hoge impact-factor. Jullie hebben me geleerd van het leven te genieten; te mogen zijn wie ik ben. Jullie vangnet (ook voor de meiden) is groots. Me door jullie gesteund voelen, brengt verre horizonnen dichtbij en maakt hoge bergtoppen klein. Altijd. Overal.

Lieve Neel, samen met een jeep afgelegen lepra-posten bezoeken is wel wat anders dan een zoektocht in een epidemiologische dataset, maar één ding blijft als onafhankelijk variabele staan: onze vriendschap door dik en dun. Dank voor al je meedenken, je trouw en je onvoorwaardelijk zijn. Ik ben er trots op dat je mijn paranimf bent!

Liefste zusje, al van het prilste begin, verkennen wij de wereld samen. Soms dichtbij elkaar, soms ver weg, maar altijd verbonden. Ik hoef je niets uit te leggen, want je begrijpt me zonder woorden. Jij vangt me als ik val. Ik sta er als jij struikelt. Samen genieten, samen ontdekken. Dat jij straks naast me staat is hoe het altijd hoort te zijn. Wij samen tegen de rest. TQM.

Lieve Duco. Dankzij jou telt dit proefschrift een hoofdstuk extra. Dankzij jou schitteren de Moai's op de voorkant. Dankzij jou bleef ik genieten naast de werkdrukke. 'No worries in the world'. De jaren samen blijven elkaar overtreffen. Maar zelfs dolfijnen in de nacht, een voorzichtige kangaroo bij de klif, of Pergolesi bij een verstild lava veld halen het niet bij wat ik voor je voel. Ik hoop dat we ons nog lang samen kunnen verwonderen, ons blijven verdiepen en trots zijn op 'de zusjes' en elkaar. Lieve PP LK X!

En tenslotte: allerliefste PD en allerliefste kleine kookaburra. Wat heeft dit proefschrift jullie gebracht? De computer op tafel, maar niet om Mega Mindy te kijken. Uit je vertrouwde huis gehaald worden om kangaroes te aaien. Maar jullie liefde is grenzeloos; mijn liefde voor jullie is grenzeloos. Jullie zijn de beste remedie tegen werkstress. Na een drukke dag 2 meisjes op schoot tegen mij aangeplakt. 's Avonds even kijken als jullie liggen te slapen. En wat is er nu mooier dan jullie lach? Dit proefschrift telt meer dan 80.000 woorden. Niet genoeg om te beschrijven hoeveel ik van jullie hou.





# Curriculum Vitae

---



Didericke (Didi) Rhebergen was born on February 8, 1973 in Groningen, the Netherlands. After graduation from high school (Christelijk Gymnasium, Leeuwarden), she joined a student exchange program and lived for one year in Jakarta, Indonesia. Since 1992, she studied Medicine at the VU University, Amsterdam and conducted a study into the knowledge, attitudes and practices of leprosy patients in Nepal. Since 1994, she combined her study Medicine, with a study Cultural Anthropology- specialization Medical Anthropology- at the VU University, Amsterdam. For her master thesis Anthropology, she spent six months in Surabaya, Indonesia, to study possible ways to empower female street sex workers in Surabaya. After she finished her medical internship cum laude, she started her residency in psychiatry at GGZ inGeest in 2002. Since 2007, she is working as an old age psychiatrist at an elderly in-patients ward of GGZinGeest. In 2006, she started working as a part-time PhD-student at the department of Psychiatry and EMGO+ at the VU University Medical Center, Amsterdam. She investigated the heterogeneity of depressive disorders, in particular the long-term course trajectories of various depressive subtypes. During her research, she finished her master in Epidemiology at the VU University. From October 2011 to April 2012 she visited the Black Dog Institute, Sydney, Australia, the Clinical Research Unit for Anxiety and Depression (CRUfAD) at the University of New South Wales in Sydney, Australia, and the University of Fremantle, Perth, Australia. Currently she is working as an old age psychiatrist, GGZ inGeest, Amsterdam, and combines clinical work and research in her daily practice. She is happy together with Duco Roolvink and they are the proud parents of Mare en Amaya.



## List of publications

---



Rhebergen D, Vogelzangs N, Hoogendijk W, Elzinga B, Van der Does W, Vreeburg S, Beekman ATF, Penninx BWJH. Can biological and psychological characteristics differentiate major depressive disorder, dysthymic disorder and double depression? Submitted.

Rhebergen D\*, Van der Steenstraten IM\*, Sunderland M, De Graaf R, Ten Have M, Lamers F, Penninx BWJH, Andrews G. An examination of generalized anxiety disorder and dysthymic disorder by latent class analysis. \*Joint first authorship. Submitted.

Rhebergen D, Huisman A, Bouckaert F, Kho K, Kok R, Sienaert P, Spaans HP, Stek M. Course trajectories of depressive disorders during ECT-treatment, identified by latent class growth analysis. Submitted.

Batelaan N, Rhebergen D, De Graaf R, Spijker J, Beekman ATF, Penninx BWJH. Panic attacks as a dimension across psychopathology: Evidence for associations with onset, course and functioning of mental disorders. *Journal of Clinical Psychiatry* 2012: In press.

Rhebergen D, Graham R, Hadzi-Pavlovic D, Stek ML, Friend P, Barrett M, Parker G. The categorisation of dysthymia: Can its constituents be meaningfully apportioned? *Journal of Affective Disorders* 2012: In press.

Rhebergen D, Arts DL, Comijs H, Beekman ATF, Terwee CB, Parker G, Stek ML. Psychometric properties of the Dutch version of the CORE measure of melancholia. *Journal of Affective Disorder* 2012: In press.

Dols A, Rhebergen D, Eikelenboom P, Stek M. Hypochondriacal delusion in an elderly woman recovers quickly with electroconvulsive therapy. *Clinics and Practice* 2012: In press.

Lamers F, Rhebergen D, Merikangas KR, De Jonge P, Beekman ATF, Penninx BWJH. Stability and transitions of depressive subtypes over 2-year follow-up. *Psychological Medicine* 2012; 17:1-11.

Attu SD, Rhebergen D, Comijs HC, Parker G, Stek ML. Psychomotor symptoms in depressed elderly patients: Assessment of the construct validity of the Dutch CORE by accelerometry. *Journal of Affective Disorders* 2012; 137: 146-50.

Rhebergen D, Lamers F, Spijker J, de Graaf R, Beekman AT, Penninx BWJ. Course trajectories of unipolar depressive disorders identified by latent class growth analysis. *Psychological Medicine* 2011; 7: 1-14.

Rhebergen D, Batelaan N, de Graaf R, Nolen W, Spijker J, Beekman AT, Penninx BW. The 7-year course of depression and anxiety in the general population. *Acta Psychiatrica Scandinavica* 2011; 123: 297-306.

Rhebergen D, Rouwenhorst A, Comijs HC, Dols A, Beekman ATF, Terwee CB, Arts D, Stek ML. De inter-beoordelaars betrouwbaarheid van de Nederlandse versie van de CORE. Een validatie studie bij opgenomen depressieve ouderen. *Tijdschrift voor Psychiatrie* 2010; 53: 49-55.

Rhebergen D, Beekman AT, De Graaf R, Nolen WA, Spijker J, Hoogendijk WJ, Penninx BW. Trajectories of recovery of social and physical functioning in major depression, dysthymic disorder and double depression: a 3-year follow-up. *Journal of Affective Disorders* 2010; 124: 148-156.

Dols A, Rhebergen D, Eikelenboom P, Stek ML. Melancholie; verstoorde beweging of ontstemd gemoed? Gevalsbeschrijving. *Tijdschrift voor Psychiatrie* 2010; 52: 583-588.

Rhebergen D, Klumpers UMH, Stek ML, Cath D. Katatonie en Clozapine. *Psyfar*, maart 2010.

Rhebergen D, Beekman AT, Graaf R, Nolen WA, Spijker J, Hoogendijk WJ, Penninx BWJ. The three-year naturalistic course of major depressive disorder, dysthymic disorder and double depression. *Journal of Affective Disorders* 2009; 115: 450-459.

Brunekreeft JA, Peerdeman SM, Rhebergen D. Subduraal hematoom en depressie. Gevalsbeschrijving. *Tijdschrift voor Psychiatrie* 2008; 50: 295-299.

Rhebergen D and Meesters PD. Chapter 1: Hollen of stilstaan. In: Meesters PD and Van Gool P (red.) *Neuropsychiatrie in de praktijk*. Maarssen: Elsevier Gezondheidszorg, 2007.

Rhebergen D, Stek ML. Reaction to 'Recent findings regarding the treatment of late-onset depression with vascular brain lesions'. *Tijdschrift voor Psychiatrie* 2006; 48: 602-603.

Rhebergen D, Yudho D, Susanti Hudiono E, Siyaranamual JR. Some Reflections on Street Sex Workers and STD/HIV/AIDS Services in Surabaya, Indonesia. *Research for Sex Work* 1999; 2: 20-21.

Rhebergen D. *Anak-Anak Jalan Diponegoro- Female Street Sex Workers in Surabaya, Indonesia*. Feministische Antropologie (11), Amsterdam, 1999. ISBN: 90-6875-064-X.



# Dissertation series

---



**DISSERTATION SERIES****Department of Psychiatry, VU University Medical Center**

N.M. (Neeltje) Batelaan (2010). Panic and Public Health: Diagnosis, Prognosis and Consequences. Vrije Universiteit Amsterdam . ISBN: 978 90 8659 411 5.

G.E. (Gideon) Anholt (2010). Obsessive-Compulsive Disorder: Spectrum Theory and Issues in Measurement. Vrije Universiteit Amsterdam.

N. (Nicole) Vogelzangs (2010). Depression and Metabolic Syndrome. Vrije Universiteit Amsterdam. ISBN: 978 90 8659 447 4.

C.M.M. (Carmilla) Licht (2010). Autonomic Nervous System Functioning in Major Depression and Anxiety Disorders. Vrije Universiteit Amsterdam. ISBN: 978 90 8659 487 0.

S.A. (Sophie) Vreeburg (2010). Hypothalamic-Pituitary-Adrenal Axis Activity in Depressive and Anxiety Disorders. Vrije Universiteit Amsterdam. ISBN: 978 90 8659 491 7.

S.N.T.M. (Sigfried) Schouws (2011). Cognitive Impairment in Older Persons with Bipolar Disorder. Vrije Universiteit Amsterdam. ISBN:978 90 9025 904 8.

P.L. (Peters) Remijnse (2011). Cognitive Flexibility in Obsessive-Compulsive Disorder and Major Depression – Functional Neuroimaging Studies on Reversal Learning and Task Switching. Vrije Universiteit Amsterdam. ISBN: 978 90 6464 449 8.

S.P. (Saskia) Wolfensberger (2011). Functional, Structural, and Molecular Imaging of the Risk for Anxiety and Depression. Vrije Universiteit Amsterdam. ISBN: 978 90 8659 536 5.

J.E. (Jenneke) Wiersma (2011). Psychological Characteristics and Treatment of Chronic Depression. Vrije Universiteit van Amsterdam. ISBN: 978 94 9121 150 8.

P.D. (Paul David) Meesters (2011). Schizophrenia in Later Life. Studies on Prevalence, Phenomenology and Care Needs (SOUL Study). Vrije Universiteit Amsterdam. ISBN: 978 90 8659 563 1.

R. (Ritsaert) Lieveise (2011). Chronobiopsychosocial Perspectives of Old Age Major Depression: a Randomized Placebo Controlled Trial with Bright Light. Vrije Universiteit Amsterdam. ISBN: 978 90 8570 858 2.

A. (Adrie) Seldenrijk (2011) Depression, Anxiety and Subclinical Cardiovascular Disease. Vrije Universiteit Amsterdam. ISBN: 978 94 6191 052 3.

Dissertation series | 252

Y. (Yuri) Milaneschi (2012) Biological Aspects of Late-life Depression. Vrije Universiteit Amsterdam. ISBN: 978 90 8659 608 9

L. (Lynn) Boschloo (2012) The Co-occurrence of Depression and Anxiety with Alcohol Use Disorders. Vrije Universiteit Amsterdam. ISBN: 978 94 6191 327 2.