

CONTENTS

1	General introduction	9
2	Simple wound exudate collection method identifies bioactive cytokines and chemokines in (arterio) venous ulcers <i>Wound Repair and Regeneration; In Press</i>	37
3	Chemokine profiles differ between intracutaneous and subcutaneous locations in early wound healing after breast reduction	57
4	Comparison of cytokine, chemokine and growth factor profiles in burn wounds, chronic wounds and surgical excision wounds	71
5	Skin substitute optimized for healing chronic wounds causes hypergranulation in excision wounds; differential roles for wound bed-derived IL-1 α and TNF- α	87
6	Chemokine-mediated migration of skin-derived stem cells: predominant role for CCL5/ RANTES <i>J Invest Dermatol, 2009;129:1569-1581</i>	107
7	Autocrine regulation of re-epithelialization after wounding by chemokine receptors CCR1, CCR10, CXCR1, CXCR2 and CXCR3 <i>J Invest Dermatol, 2011;132:215-225</i>	135
8	Keratinocyte, monocyte and endothelial cell-derived CCL27 promotes cell migration, proliferation and chemokine release in wound healing <i>Submitted for publication</i>	155
9	Structure-activity of histatin, a potent wound healing peptide from human saliva: cyclization of histatin potentiates molar activity 1,000 fold <i>FASEB, 2009;23:3928-3935</i>	175
10	Summary with options for chemokine therapies of burn and chronic wounds	193
	Nederlandse samenvatting	209
	List of publications	215
	Dankwoord	217
	Curriculum vitae	219