

References

- Agirre-Arrizubieta Z, Huiskamp GJ, Ferrier CH, van Huffelen AC, and Leijten FS (2009). Interictal magnetoencephalography and the irritative zone in the electrocorticogram. *Brain* 132.Pt 11, pp. 3060–71.
- Aguirre GK (2006). Experimental design and data analysis for fMRI. In: *Functional MRI*. Ed. by Faro S.H. and Mohamed F.B. 1st ed. Springer-Verlag. Chap. 3, pp. 59–74.
- Aguirre GK, Zarahn E, and D’Esposito M (1997). Empirical analyses of BOLD fMRI statistics. II. Spatially smoothed data collected under null-hypothesis and experimental conditions. *Neuroimage* 5.3, pp. 199–212.
- Aguirre GK, Zarahn ZE, and D’Esposito M (1998). The variability of human, BOLD hemodynamic responses. *Neuroimage* 8.4, pp. 360–369.
- Al-Asmi A, Bénar CG, Gross DW, Khani YA, Andermann F, Pike B, Dubeau F, and Gotman J (2003). fMRI activation in continuous and spike-triggered EEG-fMRI studies of epileptic spikes. *Epilepsia* 44.10, pp. 1328–39.
- Allen PJ (2010). EEG instrumentation and Safety. In: *EEG-fMRI: Physiological basis, technique, and applications*. Ed. by Mulert C; Lemieux L. 1st ed. Springer-Verlag. Chap. 7, pp. 115–133.
- Allen PJ, Polizzi G, Krakow K, Fish DR, and Lemieux L (1998). Identification of EEG events in the MR scanner: the problem of pulse artifact and a method for its subtraction. *Neuroimage* 8.3, pp. 229–39.
- Allen PJ, Josephs O, and Turner R (2000). A method for removing imaging artifact from continuous EEG recorded during functional MRI. *Neuroimage* 12.2, pp. 230–9.
- Anami K, Mori T, Tanaka F, Kawagoe Y, Okamoto J, Yarita M, Ohnishi T, Yumoto M, Matsuda H, and Saitoh O (2003). Stepping stone sampling for retrieving artifact-free electroencephalogram during functional magnetic resonance imaging. *NeuroImage* 19.2, pp. 281–295.
- Andrade-Valenca LP, Dubeau F, Mari F, Zelmann R, and Gotman J (2011). Interictal scalp fast oscillations as a marker of the seizure onset zone. *Neurology* 77.6, pp. 524–531.
- Arimoto H (2006). Multispectral polarization imaging for observing blood oxygen saturation in skin tissue. *Applied spectroscopy* 60, pp. 459–464.
- Arthuis M, Valton L, Regis J, Chauvel P, Wendling F, Naccache L, Bernard C, and Bartolomei F (2009). Impaired consciousness during temporal lobe seizures is related to increased long-distance cortical-subcortical synchronization. *Brain* 132.Pt 8, pp. 2091–101.
- Bagshaw AP, Aghakhani Y, Bénar CG, Kobayashi E, Hawco C, Dubeau F, Pike GB, and Gotman J (2004). EEG-fMRI of focal epileptic spikes: analysis with multiple haemodynamic functions and comparison with gadolinium-enhanced MR angiograms. *Hum Brain Mapp* 22.3, pp. 179–92.
- Bagshaw AP, Hawco C, Bénar CG, Kobayashi E, Aghakhani Y, Dubeau F, Pike GB, and Gotman J (2005). Analysis of the EEG-fMRI response to prolonged bursts of interictal epileptiform activity. *Neuroimage* 24.4, pp. 1099–112.
- Bagshaw AP, Kobayashi E, Dubeau F, Pike GB, and Gotman J (2006). Correspondence between EEG-fMRI and EEG dipole localisation of interictal discharges in focal epilepsy. *Neuroimage* 30.2, pp. 417–25.
- Barrat A, Barthelemy M, Pastor-Satorras R, and Vespignani A (2004). The architecture of complex weighted networks. *Proc Natl Acad Sci U S A* 101.11, pp. 3747–52.

- Bartolomei F, Wendling F, Bellanger JJ, Regis J, and Chauvel P (2001). Neural networks involving the medial temporal structures in temporal lobe epilepsy. *Clin Neurophysiol* 112.9, pp. 1746–60.
- Bartolomei F, Chauvel P, and Wendling F (2008). Epileptogenicity of brain structures in human temporal lobe epilepsy: a quantified study from intracerebral EEG. *Brain* 131.Pt 7, pp. 1818–30.
- Beall EB and Lowe MJ (2010). The non-separability of physiologic noise in functional connectivity MRI with spatial ICA at 3T. *J Neurosci Methods* 191.2, pp. 263–76.
- Becker R, Ritter P, Moosmann M, and Villringer A (2005). Visual evoked potentials recovered from fMRI scan periods. *Hum Brain Mapp* 26, pp. 221–230.
- Beckmann CF and Smith SM (2004). Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans Med Imaging* 23.2, pp. 137–52.
- Beckmann CF, DeLuca M, Devlin JT, and Smith S.M. (2005). Investigations into resting-state connectivity using independent component analysis. *Philos Trans R Soc Lond B Biol Sci* 360, pp. 1001–1013.
- Bénar CG, Gross DW, Wang Y, Petre V, Pike B, Dubeau F, and Gotman J (2002). The BOLD response to interictal epileptiform discharges. *Neuroimage* 17.3, pp. 1182–92.
- Bénar CG, Aghakhani Y, Wang Y, Izenberg A, Al-Asmi A, Dubeau F, and Gotman J (2003). Quality of EEG in simultaneous EEG-fMRI for epilepsy. *Clin Neurophysiol* 114.3, pp. 569–580.
- Bénar CG, Grova C, Kobayashi E, Bagshaw AP, Aghakhani Y, Dubeau F, and Gotman J (2006). EEG-fMRI of epileptic spikes: concordance with EEG source localization and intracranial EEG. *Neuroimage* 30.4, pp. 1161–70.
- Bénar CG, Baghsaw AP, and Lemieux L (2010). Experimental design and data analysis strategies. In: *EEG-fMRI: Physiological basis, technique, and applications*. Ed. by Mulert C and Lemieux L. 1st ed. Springer-Verlag. Chap. 12, pp. 221–257.
- Bettus G, Wendling F, Guye M, Valton L, Regis J, Chauvel P, and Bartolomei F (2008). Enhanced EEG functional connectivity in mesial temporal lobe epilepsy. *Epilepsy Res* 81.1, pp. 58–68.
- Bettus G, Bartolomei F, Confort-Gouny S, Guedj E, Chauvel P, Cozzone PJ, Ranjeva JP, and Guye M (2010). Role of resting state functional connectivity MRI in presurgical investigation of mesial temporal lobe epilepsy. *J Neurol Neurosurg Psychiatry* 81.10, pp. 1147–54.
- Bettus G, Ranjeva J, Wendling F, Bénar CG, Confort-Gouny S, Chauvel P, Cozzone PJ, Lemieux L, Bartolomei F, and Guye M (2011). Interictal Functional Connectivity of Human Epileptic Networks Assessed by Intracerebral EEG and BOLD Signal Fluctuations. *PLoS One* 6, e20071.
- Binder JR and Detre JA (2005). Functional MRI in epilepsy. In: *Magnetic resonance in epilepsy: Neuroimaging techniques*. Ed. by Kuzniecky R.I. and Jackson G.D. 2nd ed. Elsevier Academic Press. Chap. 10, pp. 281–298.
- Binnie CD and Stefan H (1999). Modern electroencephalography: its role in epilepsy management. *Clinical neurophysiology* 110, pp. 1671–1697.
- Birn RM, Cox RW, and Bandettini PA (2002). Detection versus estimation in event-related fMRI: choosing the optimal stimulus timing. *Neuroimage* 15.1, pp. 252–64.
- Birn RM, Diamond JB, Smith MA, and Bandettini PA (2006). Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. *Neuroimage* 31.4, pp. 1536–48.
- Birn RM, Smith MA, Jones TB, and Bandettini PA (2008). The respiration response function: the temporal dynamics of fMRI signal fluctuations related to changes in respiration. *Neuroimage* 40.2, pp. 644–54.
- Bromfield EB, Cavazos JE, and Sirven JI (2006). Epilepsy Surgery. In: *An introduction to epilepsy*. Ed. by Bromfield E.B., Cavazos J.E., and Sirven J.I. Vol. 1. West Hartford (CT): American Epilepsy Society. Chap. 4.
- Burock MA and Dale AM (2000). Estimation and detection of event-related fMRI signals with temporally correlated noise: a statistically efficient and unbiased approach. *Hum Brain Mapp* 11.4, pp. 249–60.
- Buxton RB, Wong EC, and Frank LR (1998). Dynamics of Blood Flow and Oxygenation Changes During Brain Activation: The Balloon Model. *Magn Reson Med* 39, pp. 855–864.
- Buxton RB, Uludag K, Dubowitz DJ, and Liu TT (2004). Modeling the hemodynamic response to brain activation. *Neuroimage* 23 Suppl 1, S220–33.
- Buzsaki G and Traub RD (2008). Physiological Basis fo the electroencephalogram and local field potentials. In: ed. by Engel J. and Pedley T. 2nd ed. Vol. 1. Lippincott Williams & Wilkins. Chap. 72.

- Carmichael DW, Thornton JS, Rodionov R, Thornton R, McEvoy A, Allen PJ, and Lemieux L (2008). Safety of localizing epilepsy monitoring intracranial electroencephalograph electrodes using MRI: radiofrequency-induced heating. eng. *J Magn Reson Imaging* 28.5, pp. 1233–1244.
- Carmichael DW, Thornton JS, Rodionov R, Thornton R, McEvoy AW, Ordidge RJ, Allen PJ, and Lemieux L (2010). Feasibility of simultaneous intracranial EEG-fMRI in humans: a safety study. eng. *Neuroimage* 49.1, pp. 379–390.
- Carmichael DW, Vulliemoz S, Rodionov R, Thornton JS, McEvoy AW, and Lemieux L (2012). Simultaneous intracranial EEG-fMRI in humans: Data quality. eng. *Neuroimage*.
- Carreno M and Luders HO (2008). General principles of presurgical evaluation. In: *Textbook of epilepsy surgery*. Ed. by Luders H.O. Vol. 1. Informa Healthcare. Chap. 47, pp. 409–422.
- Chakraborty A and McEvoy AW (2008). Presurgical functional mapping with functional MRI. *Curr Opin Neurology* 21, pp. 446–451.
- Chang C, Cunningham JP, and Glover GH (2009). Influence of heart rate on the BOLD signal: the cardiac response function. *Neuroimage* 44.3, pp. 857–69.
- Chatrin GE, Bergamini L, Dondey M, Klass DW, Lennox-Buchthal M, and Petersén I (1974). A glossary of terms most commonly used by clinical electroencephalographers. *Electroencephalography and clinical neurophysiology* 37.5, pp. 538–548.
- Chaudhary UJ, Kokkinos V, Carmichael DW, Rodionov R, Gasston D, Duncan JS, and Lemieux L (2010). Implementation and evaluation of simultaneous video-electroencephalography and functional magnetic resonance imaging. eng. *Magn Reson Imaging* 28.8, pp. 1192–1199.
- Chaudhary UJ, Duncan JS, and Lemieux L (2011). Mapping hemodynamic correlates of seizures using fMRI: A review. *Hum Brain Mapp*.
- Chaudhary UJ, Rodionov R, Carmichael DW, Thornton RC, Duncan JS, and Lemieux L (2012). Improving the sensitivity of EEG-fMRI studies of epileptic activity by modelling eye blinks, swallowing and other video-EEG detected physiological confounds. eng. *Neuroimage*.
- Cole DM, Smith SM, and Beckmann CF (2010). Advances and pitfalls in the analysis and interpretation of resting-state FMRI data. *Front Syst Neurosci* 4, p. 8.
- Colon AJ, Ossenblok P, Nieuwenhuis L, Stam KJ, and Boon P (2009). Use of routine MEG in the primary diagnostic process of epilepsy. *J Clin Neurophysiol* 26.5, pp. 326–32.
- Conner CR, Ellmore TM, Pieters TA, DiSano MA, and Tandon N (2011). Variability of the relationship between electrophysiology and BOLD-fMRI across cortical regions in humans. *J Neurosci* 31.36, pp. 12855–65.
- Crespel A, Gélisse P, Bureau M, and Genton P (2006). *The Epilepsies: EEG and epileptic syndromes*. Vol. 2. Atlas of electroencephalography. John Libbey Eurotext, p. XV.
- Cunningham CJ, Zaamout Mel F, Goodyear B, and Federico P (2008). Simultaneous EEG-fMRI in human epilepsy. *Can J Neurol Sci* 35.4, pp. 420–35.
- Dale AM (1999). Optimal experimental design for event-related fMRI. *Hum Brain Mapp* 8.2-3, pp. 109–14.
- Damoiseaux JS, Rombouts SA, Barkhof F, Scheltens P, Stam CJ, Smith SM, and Beckmann CF (2006). Consistent resting-state networks across healthy subjects. *Proc Natl Acad Sci U S A* 103.37, pp. 13848–53.
- de Munck JC, Gonçalves SI, Huijboom L, Kuijer JP, Pouwels PJ, Heethaar RM, and Lopes da Silva FH (2007). The hemodynamic response of the alpha rhythm: an EEG/fMRI study. *Neuroimage* 35.3, pp. 1142–51.
- de Munck JC, Gonçalves SI, Faes TJ, Kuijer JP, Pouwels PJ, Heethaar RM, and Lopes da Silva FH (2008). A study of the brain's resting state based on alpha band power, heart rate and fMRI. *Neuroimage* 42.1, pp. 112–21.
- de Munck JC, Gonçalves SI, Mammoliti R, Heethaar RM, and Lopes da Silva FH (2009a). Interactions between different EEG frequency bands and their effect on alpha-fMRI correlations. *Neuroimage* 47.1, pp. 69–76.
- de Munck JC and Ossenblok PPW (2009b). Towards clinical standards for EEG/fMRI. eng. *Clin Neurophysiol* 120.9, pp. 1617–1618.

- de Munck JC, Gonçalves SI, van Houdt PJ, Mammoliti R, Ossenblok P, and Lopes da Silva FH (2010). The hemodynamic response of EEG features. In: *EEG-fMRI: Physiological basis, technique, and applications*. Ed. by Mulert C and Lemieux L. Springer-Verlag. Chap. 3.6, pp. 195–208.
- de Munck JC, van Houdt PJ, Verdaasdonk RM, and Ossenblok PP (2012). A semi-automatic method to determine electrode positions and labels from gel artifacts in EEG/fMRI-studies. *Neuroimage* 59.1, pp. 399–403.
- de Munck JC, van Houdt PJ, Gonçalves SI, van Wegen E, and Ossenblok PPW (2013). Novel artefact removal algorithms for co-registered EEG/fMRI based on selective averaging and subtraction. *Neuroimage* 64, pp. 407–415.
- De Martino F, Gentile F, Esposito F, Balsi M, Di Salle F, Goebel R, and Formisano E (2007). Classification of fMRI independent components using IC-fingerprints and support vector machine classifiers. *Neuroimage* 34.1, pp. 177–94.
- Debener S, Mullinger KJ, Niazy RK, and Bowtell RW (2008). Properties of the ballistocardiogram artefact as revealed by EEG recordings at 1.5, 3 and 7 T static magnetic field strength. *Int J Psychophysiol* 67.3, pp. 189–99.
- Debener S, Kranczioch C, and Gutberlet I (2010a). EEG quality: origin and reduction of the EEG cardiac-related artefact. In: *EEG-fMRI: Physiological basis, technique, and applications*. Ed. by Mulert C. and Lemieux L. Springer-Verlag. Chap. 8, pp. 135–151.
- Debener S, Thorne J, Schneider TR, and Campos Viola F (2010b). Using ICA for the analysis of Multi-Channel EEG data. In: *Simultaneous EEG and fMRI: recording, analysis and application*. Ed. by Ullsperger M.; Debener S. Oxford University Press. Chap. 3.1, pp. 121–133.
- Deichmann R, Noth U, and Weiskopf N (2010). The basics of functional magnetic resonance imaging. In: *EEG-fMRI: physiological basis, technique, and applications*. Ed. by Mulert C. and Lemieux L. 1st ed. Springer-Verlag. Chap. 3, pp. 39–62.
- Diehl B, Tkach J, Piao Z, Ruggieri P, LaPresto E, Liu P, Fisher E, Bingaman W, and Najm I (2010). Diffusion tensor imaging in patients with focal epilepsy due to cortical dysplasia in the temporo-occipital region: electro-clinico-pathological correlations. *eng. Epilepsy Res* 90.3, pp. 178–187.
- Duncan JS (2010). Imaging in the surgical treatment of epilepsy. *Nat Rev Neurol* 6, pp. 537–550.
- Eccher M and Nair D (2008). The irritative zone: general principles. In: *Textbook of epilepsy surgery*. Ed. by Luders H.O. Informa Healthcare. Chap. 55, pp. 503–511.
- Eichele T, Specht K, Moosmann M, Jongsma MLA, Quiroga RQ, Nordby H, and Hugdahl K (2005). Assessing the spatiotemporal evolution of neuronal activation with single-trial event-related potentials and functional MRI. *eng. Proc Natl Acad Sci U S A* 102.49, pp. 17798–17803.
- Eichele T, Moosmann M, L Wu, Gutberlet I, and Debener S (2010). Removal of MRI artifacts from EEG recordings. In: *Simultaneous EEG and fMRI: recording, analysis, and application*. Ed. by Ullsperger M. and Debener S. Oxford University Press. Chap. 2.3, pp. 95–106.
- Ellingson ML, Liebenthal E, Spanaki MV, Prieto TE, Binder JR, and Ropella KM (2004). Ballistocardiogram artifact reduction in the simultaneous acquisition of auditory ERPS and fMRI. *NeuroImage* 22, pp. 1534–1542.
- Federico P, Abbott DF, Briellmann RS, Harvey AS, and Jackson GD (2005). Functional MRI of the pre-ictal state. *Brain* 128.Pt 8, pp. 1811–7.
- Fisher RS, van Emde Boas W, Blume W, Elger C, Genton P, Lee P, and Engel J Jr (2005). Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 46.4, pp. 470–2.
- Flanagan D, Abbott DF, and Jackson GD (2009). How wrong can we be? The effect of inaccurate mark-up of EEG/fMRI studies in epilepsy. *Clin Neurophysiol* 120.9, pp. 1637–47.
- Freyer F, Becker R, Anami K, Curio G, Villringer A, and Ritter P (2009). Ultrahigh-frequency EEG during fMRI: pushing the limits of imaging-artifact correction. *eng. Neuroimage* 48.1, pp. 94–108.
- Friston KJ, Williams S, Howard R, Frackowiak RS, and Turner R (1996). Movement-related effects in fMRI time-series. *eng. Magn Reson Med* 35.3, pp. 346–355.
- Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg MD, and Turner R (1998). Event-related fMRI: characterizing differential responses. *Neuroimage* 7.1, pp. 30–40.

- Genovese CR, Lazar NA, and Nichols T (2002). Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage* 15.4, pp. 870–8.
- Gholipour T, Moeller F, Pittau F, Dubeau F, and Gotman J (2011). Reproducibility of interictal EEG-fMRI results in patients with epilepsy. *Epilepsia* 52, pp. 433–442.
- Glover GH (1999). Deconvolution of impulse response in event-related BOLD fMRI. *Neuroimage* 9.4, pp. 416–29.
- Glover GH, Li TQ, and Ress D (2000). Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magn Reson Med* 44.1, pp. 162–7.
- Goldman RI, Stern JM, Engel J Jr, and Cohen MS (2002). Simultaneous EEG and fMRI of the alpha rhythm. *Neuroreport* 13.18, pp. 2487–92.
- Gonçalves S.I., de Munck J.C., Pouwels P.J., Schoonhoven R., Kuijser J.P., Maurits N.M., Hoogduin J.M., Van Someren E.J., Heethaar R.M., and Lopes da Silva F.H. (2006). Correlating the alpha rhythm to BOLD using simultaneous EEG/fMRI: inter-subject variability. *Neuroimage* 30.1, pp. 203–13.
- Gonçalves S.I., Pouwels P.J., Kuijser J.P., Heethaar R.M., and de Munck J.C. (2007). Artifact removal in co-registered EEG/fMRI by selective average subtraction. *Clin Neurophysiol* 118.11, pp. 2437–50.
- Gonçalves SI, Bijma F, Pouwels PJW, Jonker MA, Kuijser JPA, Heethaar RM, Lopes da Silva FH, and De Munck JC (2008). A data and model-driven approach to explore inter-subject variability of resting-state brain activity using EEG-fMRI. *IEEE J Sel Top SP* 2.6, pp. 944–953.
- Gotman J, Bénar CG, and Dubeau F (2004). Combining EEG and FMRI in epilepsy: methodological challenges and clinical results. *J Clin Neurophysiol* 21.4, pp. 229–40.
- Gotman J, Grova C, Bagshaw A, Kobayashi E, Aghakhani Y, and Dubeau F (2005). Generalized epileptic discharges show thalamocortical activation and suspension of the default state of the brain. *Proc Natl Acad Sci U S A* 102.42, pp. 15236–40.
- Gotman J, Kobayashi E, Bagshaw AP, Bénar CG, and Dubeau F (2006). Combining EEG and fMRI: a multimodal tool for epilepsy research. *J Magn Reson Imaging* 23.6, pp. 906–20.
- Grouiller F, Vercueil L, Krainik A, Segebarth C, Kahane P, and David O (2007). A comparative study of different artefact removal algorithms for EEG signals acquired during functional MRI. *Neuroimage* 38.1, pp. 124–37.
- Grouiller F, Vercueil L, Krainik A, Segebarth C, Kahane P, and David O (2010). Characterization of the hemodynamic modes associated with interictal epileptic activity using a deformable model-based analysis of combined EEG and functional MRI recordings. *Hum Brain Mapp* 31.8, pp. 1157–1173.
- Grouiller F, Thornton RC, Groening K, Spinelli L, Duncan JS, Schaller K, Siniatchkin M, Lemieux L, Seeck M, Michel CM, and Vulliemoz S (2011). With or without spikes: localization of focal epileptic activity by simultaneous electroencephalography and functional magnetic resonance imaging. *Brain* 134, pp. 2867–2886.
- Grova C, Daunizeau J, Kobayashi E, Bagshaw AP, Lina JM, Dubeau F, and Gotman J (2008). Concordance between distributed EEG source localization and simultaneous EEG-fMRI studies of epileptic spikes. *Neuroimage* 39.2, pp. 755–74.
- Gutberlet I (2010). Recording EEG signals inside the MRI. In: *Simultaneous EEG and fMRI: Recording, analysis, and application*. Ed. by Ullsperger M.; Debener S. Oxford university press. Chap. 2.1, pp. 69–83.
- Hamandi K, Powell HW, Laufs H, Symms MR, Barker GJ, Parker GJ, Lemieux L, and Duncan JS (2008). Combined EEG-fMRI and tractography to visualise propagation of epileptic activity. *J Neurol Neurosurg Psychiatry* 79.5, pp. 594–7.
- Hamer HM (2008). Noninvasive electroencephalography for the evaluation of the irritative zone. In: *Textbook of epilepsy surgery*. Ed. by Luders H.O. Vol. 1. Informa Healthcare. Chap. 56, pp. 512–520.
- Handwerker DA, Ollinger JM, and D'Esposito M (2004). Variation of BOLD hemodynamic responses across subjects and brain regions and their effects on statistical analyses. *Neuroimage* 21.4, pp. 1639–51.
- Hauf M, Jann K, Schindler K, Scheidegger O, Meyer C, Rummel C, Mariani C, Koenig T, and Wiest R (2012). Localizing Seizure-Onset Zones in Presurgical Evaluation of Drug-Resistant Epilepsy by Electroencephalography/fMRI: Effectiveness of Alternative Thresholding Strategies. *eng. AJNR Am J Neuroradiol*.

- Hawco CS, Bagshaw AP, Lu Y, Dubeau F, and Gotman J (2007). BOLD changes occur prior to epileptic spikes seen on scalp EEG. *Neuroimage* 35.4, pp. 1450–8.
- Heller R, Stanley D, Yekutieli D, Rubin N, and Benjamini Y (2005). Cluster-based analysis of fMRI data. *Neuroimage* 33, pp. 599–608.
- Herrmann CS and Debener S (2008). Simultaneous recording of EEG and BOLD responses: a historical perspective. *Int J Psychophysiol* 67.3, pp. 161–8.
- Hjorth B (1975). An on-line transformation of EEG scalp potentials into orthogonal source derivations. *Electroencephalogr Clin Neurophysiol* 39.5, pp. 526–30.
- Huettel SA, Song AW, and McCarthy G (2009a). BOLD fMRI: origin and properties. In: *Functional magnetic resonance imaging*. Ed. by Huettel S.A., Song A.W., and McCarthy G. 2nd ed. Sinauer Associates, In.c. Chap. 7, pp. 193–242.
- Huettel SA, Song AW, and McCarthy G (2009b). Experimental design. In: *Functional magnetic resonance imaging*. Ed. by Huettel S.A., Song A.W., and McCarthy G. Sinauer Associates, In.c. Chap. 9, pp. 293–329.
- Huettel SA, Song AW, and McCarthy G (2009c). From neuronal to hemodynamic activity. In: *Functional magnetic resonance imaging*. Ed. by Huettel S.A., Song A.W., and McCarthy G. 2nd ed. Sinauer Associates, In.c. Chap. 6, pp. 159–192.
- Huettel SA, Song AW, and McCarthy G (2009d). MR contrast mechanisms and pulse sequences. In: *Functional magnetic resonance imaging*. Ed. by Huettel S.A., Song A.W., and McCarthy G. 2nd ed. Sinauer Associates, In.c. Chap. 5, pp. 121–157.
- Huiskamp GJM (2005). Reduction of the Ballistocardiogram Artifact in Simultaneous EEG-fMRI using ICA. eng. *Conf Proc IEEE Eng Med Biol Soc* 4, pp. 3691–3694.
- Ives JR, Warach S, Schmitt F, Edelman RR, and Schomer DL (1993). Monitoring the patient's EEG during echo planar MRI. *Electroencephalogr Clin Neurophysiol* 87, pp. 417–420.
- Jackson G (2008). EEG-fMRI in the presurgical work-up of patients with partial epilepsy and unclear seizure focus. *Nat Clin Pract Neurol* 4.3, pp. 128–9.
- Jacobs J, Kobayashi E, Boor R, Muhle H, Stephan W, Hawco C, Dubeau F, Jansen O, Stephani U, Gotman J, and Siniatchkin M (2007). Hemodynamic responses to interictal epileptiform discharges in children with symptomatic epilepsy. *Epilepsia* 48.11, pp. 2068–78.
- Jacobs J, Hawco C, Kobayashi E, Boor R, LeVan P, Stephani U, Siniatchkin M, and Gotman J (2008). Variability of the hemodynamic response as a function of age and frequency of epileptic discharge in children with epilepsy. *Neuroimage* 40.2, pp. 601–14.
- Jacobs J, Levan P, Moeller F, Boor R, Stephani U, Gotman J, and Siniatchkin M (2009). Hemodynamic changes preceding the interictal EEG spike in patients with focal epilepsy investigated using simultaneous EEG-fMRI. *Neuroimage* 45.4, pp. 1220–31.
- Jacobs J, Staba R, Asano E, Otsubo H, Wu JW, Zijlmans M, Mohamed I, Kahane P, Dubeau F, Navarro V, and Gotman J (2012). High-frequency oscillations (HFOs) in clinical epilepsy. eng. *Prog Neurobiol* in press.
- Josephs O, Turner R, and Friston K (1997). Event-Related fMRI. *Hum Brain Mapp* 5.4, pp. 243–248.
- Kaufmann C, Wehrle R, Wetter TC, Holsboer F, Auer DP, Pollmacher T, and Czisch M (2006). Brain activation and hypothalamic functional connectivity during human non-rapid eye movement sleep: an EEG/fMRI study. *Brain* 129.Pt 3, pp. 655–67.
- Kilner JM, Mattout J, Henson R, and Friston KJ (2005). Hemodynamic correlates of EEG: A heuristic. *Neuroimage* 28, pp. 280–286.
- Kim SG and Bandettini PA (2006). Principles of functional MRI. In: *Functional MRI*. Ed. by Faro S.H. and Mohamed F.B. 1st ed. Springer-Verlag. Chap. 1, pp. 3–23.
- Kobayashi E, Bagshaw AP, Grova C, Dubeau F, and Gotman J (2006a). Negative BOLD responses to epileptic spikes. *Hum Brain Mapp* 27.6, pp. 488–97.
- Kobayashi E, Bagshaw AP, Bénar CG, Aghakhani Y, Andermann F, Dubeau F, and Gotman J (2006b). Temporal and extratemporal BOLD responses to temporal lobe interictal spikes. *Epilepsia* 47.2, pp. 343–54.

- Kobayashi E, Hawco CS, Grova C, Dubeau F, and Gotman J (2006c). Widespread and intense BOLD changes during brief focal electrographic seizures. *Neurology* 66.7, pp. 1049–55.
- Krakow K (2008). Imaging epileptic activity using functional MRI. *Neurodegener Dis* 5.5, pp. 286–95.
- Krakow K, Allen PJ, Lemieux L, Symms MR, and Fish DR (2000). Methodology: EEG-correlated fMRI. In: *Functional imaging in the epilepsies*. Ed. by Henry T.R., J.S. Duncan, and S.F. Berkovic. Lipp. Chap. 18, pp. 187–201.
- Krakow K, Messina D, Lemieux L, Duncan JS, and Fish DR (2001). Functional MRI activation of individual interictal epileptiform spikes. *Neuroimage* 13.3, pp. 502–5.
- Krakow K and Lengler U (2007). Presurgical functional MRI in epilepsy. In: *Clinical functional MRI: presurgical functional neuroimaging*. Ed. by Stippich C. and Sartor K. 1st ed. Springer-Verlag. Chap. 5, pp. 135–148.
- Lardizabal DV (2008). Medical Intractability in epilepsy. In: *Textbook of epilepsy surgery*. Ed. by Luders H.O. Vol. 1. Informa Healthcare. Chap. 25, pp. 203–204.
- Laufs H (2008). Endogenous brain oscillations and related networks detected by surface EEG-combined fMRI. *Hum Brain Mapp* 29.7, pp. 762–9.
- Laufs H (2012). Functional imaging of seizures and epilepsy: evolution from zones to networks. *Curr Opin Neurol*.
- Laufs H, Kleinschmidt A, Beyerle A, Eger E, Salek-Haddadi A, Preibisch C, and Krakow K (2003a). EEG-correlated fMRI of human alpha activity. *Neuroimage* 19.4, pp. 1463–76.
- Laufs H, Krakow K, Sterzer P, Eger E, Beyerle A, Salek-Haddadi A, and Kleinschmidt A (2003b). Electroencephalographic signatures of attentional and cognitive default modes in spontaneous brain activity fluctuations at rest. *Proc Natl Acad Sci U S A* 100.19, pp. 11053–8.
- Laufs H, Walker MC, and Lund TE (2007). 'Brain activation and hypothalamic functional connectivity during human non-rapid eye movement sleep: an EEG/fMRI study'—its limitations and an alternative approach. *Brain* 130.Pt 7, e75; author reply e76.
- Laufs H, Daunizeau J, Carmichael DW, and Kleinschmidt A (2008). Recent advances in recording electrophysiological data simultaneously with magnetic resonance imaging. *Neuroimage* 40.2, pp. 515–28.
- Laufs H and Thornton R (2010). Clinical applications: epilepsy. In: *Simultaneous EEG and fMRI*. Ed. by Ullsperger M. and Debener S. Oxford University Press. Chap. 4.2, pp. 295–310.
- Lazeyras F, Blanke O, Perrig S, Zimine I, Golay X, Delavelle J, Michel CM, de Tribolet N, Villemure JG, and Seeck M (2000). EEG-triggered functional MRI in patients with pharmacoresistant epilepsy. *J Magn Reson Imaging* 12.1, pp. 177–85.
- Lemieux L, Allen PJ, Franconi F, Symms MR, and Fish DR (1997). Recording of EEG during fMRI experiments: patient safety. *MRM* 38, pp. 943–952.
- Lemieux L, Salek-Haddadi A, Lund TE, Laufs H, and Carmichael D (2007). Modelling large motion events in fMRI studies of patients with epilepsy. *Magn Reson Imaging* 25.6, pp. 894–901.
- Lemieux L, Laufs H, Carmichael D, Paul JS, Walker MC, and Duncan JS (2008). Noncanonical spike-related BOLD responses in focal epilepsy. *Hum Brain Mapp* 29.3, pp. 329–45.
- Lemieux L, Daunizeau J, and Walker MC (2011). Concepts of connectivity and human epileptic activity. *Front Syst Neurosci* 5, p. 12.
- LeVan P and Gotman J (2009). Independent component analysis as a model-free approach for the detection of BOLD changes related to epileptic spikes: a simulation study. *Hum Brain Mapp* 30.7, pp. 2021–31.
- LeVan P, Tyvaert L, Moeller F, and Gotman J (2010a). Independent component analysis reveals dynamic ictal BOLD responses in EEG-fMRI data from focal epilepsy patients. *Neuroimage* 49.1, pp. 366–78.
- LeVan P, Zahneisen B, Grotz T, and Hennig J (2010b). Robust hemodynamic parameter estimation within visual cortex using high temporal resolution fMRI. In: *16th Annual Meeting of the Organization for Human Brain Mapping*.
- Lindberg LG, Tamura T, and Oberg PA (1991). Photoplethysmography. Part 1. Comparison with laser Doppler flowmetry. *Med Biol Eng Comput* 29.1, pp. 40–47.
- Linder SP, Wendelken SM, Wei E, and McGrath SP (2006). Using the morphology of photoplethysmogram peaks to detect changes in posture. *J Clin Monit Comput* 20.3, pp. 151–8.

- Lindquist MA and Wager TD (2007). Validity and power in hemodynamic response modeling: a comparison study and a new approach. *Hum Brain Mapp* 28.8, pp. 764–84.
- Liu TT, Frank LR, Wong EC, and Buxton RB (2001). Detection power, estimation efficiency, and predictability in event-related fMRI. *Neuroimage* 13.4, pp. 759–73.
- Liu TT and Frank LR (2004). Efficiency, power, and entropy in event-related FMRI with multiple trial types. Part I: theory. *Neuroimage* 21.1, pp. 387–400.
- Liu Y, Yang T, Yang X, Liu I, Liao W, Lui S, Huang X, Chen H, Gong Q, and Zhou D (2008). EEG-fMRI study of the interictal epileptic activity in patients with partial epilepsy. *J Neurol Sci* 268.1-2, pp. 117–23.
- Logothetis NK, Pauls J, Augath M, Trinath T, and Oeltermann A (2001). Neurophysiological investigation of the basis of the fMRI signal. *Nature* 412.6843, pp. 150–7.
- Lopes da Silva F, Pijn JP, and Boeijinga P (1989). Interdependence of EEG signals: linear vs. nonlinear associations and the significance of time delays and phase shifts. *Brain Topogr* 2.1-2, pp. 9–18.
- Lopes da Silva F and van Rotterdam A (2005). Biophysical aspects of EEG and Magnetoencephalogram generation. In: *Electroencephalography*. Ed. by Niedermeyer E. and Lopes da Silva F. 5th ed. Lippincott Williams & Wilkins. Chap. 5.
- Lu Y, Bagshaw AP, Grova C, Kobayashi E, Dubeau F, and Gotman J (2006). Using voxel-specific hemodynamic response function in EEG-fMRI data analysis. *Neuroimage* 32.1, pp. 238–47.
- Lu Y, Grova C, Kobayashi E, Dubeau F, and Gotman J (2007). Using voxel-specific hemodynamic response function in EEG-fMRI data analysis: An estimation and detection model. *Neuroimage* 34.1, pp. 195–203.
- Lund TE, Madsen KH, Sidaros K, Luo WL, and Nichols TE (2006). Non-white noise in fMRI: does modelling have an impact? *Neuroimage* 29.1, pp. 54–66.
- Luo C, Li Q, Lai Y, Xia Y, Qin Y, Liao W, Li S, Zhou D, Yao D, and Gong Q (2011). Altered functional connectivity in default mode network in absence epilepsy: a resting-state fMRI study. *eng. Hum Brain Mapp* 32.3, pp. 438–449.
- Mandelkow H, Halder P, Boesiger P, and Brandeis D (2006). Synchronization facilitates removal of MRI artefacts from concurrent EEG recordings and increases usable bandwidth. *NeuroImage* 32.3, pp. 1120–1126.
- Manganotti P, Formaggio E, Gasparini A, Cerini R, Bongiovanni LG, Storti SF, Mucelli RP, Fiaschi A, and Avesani M (2008). Continuous EEG-fMRI in patients with partial epilepsy and focal interictal slow-wave discharges on EEG. *Magn Reson Imaging* 26.8, pp. 1089–100.
- Mantini D, Perrucci MG, Del Gratta C, Romani GL, and Corbetta M (2007). Electrophysiological signatures of resting state networks in the human brain. *Proc Natl Acad Sci U S A* 104.32, pp. 13170–5.
- Masterton RA, Abbott DF, Fleming SW, and Jackson GD (2007). Measurement and reduction of motion and ballistocardiogram artefacts from simultaneous EEG and fMRI recordings. *Neuroimage* 37.1, pp. 202–11.
- Mathern GW and Sperling MR (2008). Presurgical evaluation: General principles and methods. In: *Epilepsy: A comprehensive textbook*. Ed. by Engel J. and Pedley T.A. Vol. 2. Lippincott Williams & Wilkins. Chap. 169.
- McGill ML, Devinsky O, Kelly C, Milham M, Castellanos FX, Quinn BT, DuBois J, Young JR, Carlson C, French J, Kuzniecky R, Halgren E, and Thesen T (2012). Default mode network abnormalities in idiopathic generalized epilepsy. *eng. Epilepsy Behav* 23.3, pp. 353–359. doi: 10.1016/j.yebeh.2012.01.013.
- McKeown MJ, Makeig S, Brown GG, Jung TP, Kindermann SS, Bell AJ, and Sejnowski TJ (1998). Analysis of fMRI data by blind separation into independent spatial components. *Hum Brain Mapp* 6.3, pp. 160–88.
- Meeren HK, Pijn JP, Van Luijckelaar EL, Coenen AM, and Lopes da Silva FH (2002). Cortical focus drives widespread corticothalamic networks during spontaneous absence seizures in rats. *J Neurosci* 22.4, pp. 1480–95.
- Moehring J, Coropceanu D, Galka A, Moeller F, Wolff S, Boor R, Jansen O, Stephani U, and Siniatchkin M (2011). Improving sensitivity of EEG-fMRI studies in epilepsy: The role of sleep-specific activity. *eng. Neurosci Lett* 505.2, pp. 211–215.

- Moeller F, Siebner HR, Wolff S, Muhle H, Boor R, Granert O, Jansen O, Stephani U, and Siniatchkin M (2008). Changes in activity of striato-thalamo-cortical network precede generalized spike wave discharges. *Neuroimage* 39.4, pp. 1839–49.
- Moeller F, Tyvaert L, Nguyen DK, LeVan P, Bouthillier A, Kobayashi E, Tampieri D, Dubeau F, and Gotman J (2009). EEG-fMRI: adding to standard evaluations of patients with nonlesional frontal lobe epilepsy. *Neurology* 73.23, pp. 2023–30.
- Moeller F, Levan P, and Gotman J (2011). Independent component analysis (ICA) of generalized spike wave discharges in fMRI: Comparison with general linear model-based EEG-fMRI. *Hum Brain Mapp* 32.2, pp. 209–217.
- Moosmann M, Ritter P, Krastel I, Brink A, Thees S, Blankenburg F, Taskin B, Obrig H, and Villringer A (2003). Correlates of alpha rhythm in functional magnetic resonance imaging and near infrared spectroscopy. *Neuroimage* 20.1, pp. 145–58.
- Moosmann M, Schönfelder V.H., Specht K., Scheeringa R., Nordby H., and Hugdahl K. (2009). Realignment parameter-informed artefact correction for simultaneous EEG-fMRI recording. *Neuroimage* 45, 1144–1150.
- Mullinger KJ, Morgan PS, and Bowtell RW (2008). Improved artifact correction for combined electroencephalography/functional MRI by means of synchronization and use of vectorcardiogram recordings. *Journal of Magnetic Resonance Imaging* 27, pp. 607–616.
- Murray WB and Foster PA (1996). The peripheral pulse wave: information overlooked. *J Clin Monit* 12.5, pp. 365–77.
- Nakamura W, Anami K, Mori T, Saitoh O, Cichocki A, and Amari S (2006). Removal of ballistocardiogram artifacts from simultaneously recorded EEG and fMRI data using independent component analysis. *eng. IEEE Trans Biomed Eng* 53.7, pp. 1294–1308.
- Napadow V, Dhond R, Conti G, Makris N, Brown EN, and Barbieri R (2008). Brain correlates of autonomic modulation: combining heart rate variability with fMRI. *Neuroimage* 42.1, pp. 169–77.
- Negishi M, Abildgaard M, Nixon T, and Constable RT (2004). Removal of time-varying gradient artefacts during continuous fMRI. *Clin Neurophysiol* 115, pp. 2181–2192.
- Negishi M, Martuzzi R, Novotny EJ, Spencer DD, and Constable RT (2011). Functional MRI connectivity as a predictor of the surgical outcome of epilepsy. *Epilepsia* 52.9, pp. 1733–40.
- Nguyen DK and Spencer SS (2004). Invasive EEG in presurgical evaluation of epilepsy. In: *The treatment of epilepsy*. Ed. by Shorvon S., Perruca E., Fish D.R., and Dodson E. 2nd ed. Vol. 1. Blackwell Science. Chap. 53, pp. 609–635.
- Niazy RK, Beckmann CF, Iannetti GD, Brady JM, and Smith SM (2005). Removal of fMRI environment artifacts from EEG data using optimal basis sets. *NeuroImage* 28, pp. 720–737.
- Noachtar S and Borggraefe I (2009a). Epilepsy surgery: a critical review. *Epilepsy Behav* 15.1, pp. 66–72.
- Noachtar S and Remi J (2009b). The role of EEG in epilepsy: a critical review. *Epilepsy Behav* 15.1, pp. 22–33.
- Noll DC and Scheider D (1994). Theory, simulation, and compensation of physiological motion artifacts in functional MRI. In: *Proceedings of IEEE International Conference on Image Processing*. Austin, Texas, pp. 40–44.
- Oostenveld R, Fries P, Maris E, and Schoffelen JM (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intell Neurosci* 2011, p. 156869.
- Ortega GJ, Menendez de la Prida L, Sola RG, and Pastor J (2008). Synchronization clusters of interictal activity in the lateral temporal cortex of epileptic patients: intraoperative electrocorticographic analysis. *Epilepsia* 49.2, pp. 269–80.
- Ossenblok P, de Munck JC, Colon A, Drolsbach W, and Boon P (2007). Magnetoencephalography is more successful for screening and localizing frontal lobe epilepsy than electroencephalography. *Epilepsia* 48.11, pp. 2139–49.
- Palmini A (2008). The irritative zone evaluated with invasive recordings. In: *Textbook of epilepsy surgery*. Ed. by Luders H.O. Vol. 1. Informa Healthcare. Chap. 57, pp. 521–529.
- Panayiotopoulos CP (2007). Principles of therapy in the epilepsies. In: *A clinical guide to epileptic syndromes and their treatment*. Ed. by Panayiotopoulos C.P. 2nd ed. Vol. 1. Springer-Verlag. Chap. 7, pp. 155–179.
- Passaro EA (2006). *Presurgical evaluation of medically intractable epilepsy*.

- Pijn JP, Vijn PC, Lopes da Silva FH, Van Ende Boas W, and Blanes W (1990). Localization of epileptogenic foci using a new signal analytical approach. *Neurophysiol Clin* 20.1, pp. 1–11.
- Pittau F, Levan P, Moeller F, Gholipour T, Haegelen C, Zelmann R, Dubeau F, and Gotman J (2011). Changes preceding interictal epileptic EEG abnormalities: Comparison between EEG/fMRI and intracerebral EEG. *Epilepsia* 52.6, pp. 1120–9.
- Pittau F, Dubeau F, and Gotman J (2012). Contribution of EEG/fMRI to the definition of the epileptic focus. *eng. Neurology* 78.19, pp. 1479–1487.
- Press WH, Flannery BP, Teukolsky SA, and Wetterling WT (2007). *Numerical Recipes in C++*. Ed. by Press WH., Flannery B.P., Teukolsky S.A., and Wetterling W.T. 3rd ed. Cambridge University Press, pp. 82–93.
- Rathakrishnan R, Moeller F, Levan P, Dubeau F, and Gotman J (2010). BOLD signal changes preceding negative responses in EEG-fMRI in patients with focal epilepsy. *Epilepsia* 51.9, pp. 1837–45.
- Richardson M (2003). Epilepsy and surgical mapping. *British Medical Bulletin* 65, pp. 179–192.
- Richardson M (2010). Current themes in neuroimaging of epilepsy: brain networks, dynamic phenomena, and clinical relevance. *Clin Neurophysiol* 121.8, pp. 1153–75.
- Rodionov R, De Martino F, Laufs H, Carmichael DW, Formisano E, Walker M, Duncan JS, and Lemieux L (2007). Independent component analysis of interictal fMRI in focal epilepsy: comparison with general linear model-based EEG-correlated fMRI. *Neuroimage* 38.3, pp. 488–500.
- Rosa MJ, Kilner J, Blankenburg F, Josephs O, and Penny W (2010). Estimating the transfer function from neuronal activity to BOLD using simultaneous EEG-fMRI. *Neuroimage* 49.2, pp. 1496–509.
- Rosenkranz K and Lemieux L (2010). Present and future of simultaneous EEG-fMRI. *Magma* 23.5-6, pp. 309–16.
- Rosenow F and Luders H (2001). Presurgical evaluation of epilepsy. *Brain* 124.Pt 9, pp. 1683–700.
- Rusch TL, Sankar R, and Scharf JE (1996). Signal processing methods for pulse oximetry. *Comput Biol Med* 26.2, pp. 143–59.
- Salek-Haddadi A, Friston KJ, Lemieux L, and Fish DR (2003). Studying spontaneous EEG activity with fMRI. *Brain Res Brain Res Rev* 43.1, pp. 110–33.
- Salek-Haddadi A, Diehl B, Hamandi K, Merschhemke M, Liston A, Friston K, Duncan JS, Fish DR, and Lemieux L (2006). Hemodynamic correlates of epileptiform discharges: an EEG-fMRI study of 63 patients with focal epilepsy. *Brain Res* 1088.1, pp. 148–66.
- Schabus M, Dang-Vu TT, Albouy G, Baletau E, Boly M, Carrier J, Darsaud A, Degueldre C, Desseilles M, Gais S, Phillips C, Rauchs G, Schnakers C, Sterpenich V, Vandewalle G, Luxen A, and Maquet P (2007). Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. *Proc Natl Acad Sci U S A* 104.32, pp. 13164–9.
- Schridde U, Khubchandani M, Motelow JE, Sanganahalli BG, Hyder F, and Blumenfeld H (2008). Negative BOLD with large increases in neuronal activity. *Cereb Cortex* 18.8, pp. 1814–27.
- Shamir M, Eidelman LA, Floman Y, Kaplan L, and Pizov R (1999). Pulse oximetry plethysmographic waveform during changes in blood volume. *Br J Anaesth* 82.2, pp. 178–81.
- Shelley KH (2007). Photoplethysmography: beyond the calculation of arterial oxygen saturation and heart rate. *Anesth Analg* 105.6 Suppl, S31–6, tables of contents.
- Shirer WR, Ryali S, Rykhlevskaia E, Menon V, and Greicius M (2012). Decoding Subject-Driven Cognitive States with Whole-Brain Connectivity Patterns. *Cereb Cortex* 22, pp. 158–165.
- Shmuel A (2010). Locally measured neuronal correlates of functional MRI signals. In: *EEG-fMRI: Physiological basis, technique, and applications*. Ed. by Mulert C. and Lemieux L. 1st ed. Springer-Verlag. Chap. 4, pp. 63–82.
- Shmuel A, Augath M, Oeltermann A, and Logothetis NK (2006). Negative functional MRI response correlates with decreases in neuronal activity in monkey visual area V1. *Nature Neuroscience* 9, pp. 569–577.
- Shmueli K, van Gelderen P, de Zwart JA, Horovitz SG, Fukunaga M, Jansma JM, and Duyn JH (2007). Low-frequency fluctuations in the cardiac rate as a source of variance in the resting-state fMRI BOLD signal. *Neuroimage* 38.2, pp. 306–20.

- Siero JC, Petridou N, Hoogduin H, Luijten PR, and Ramsey NF (2011). Cortical depth-dependent temporal dynamics of the BOLD response in the human brain. *J Cereb Blood Flow Metab* 31.10, pp. 1999–2008.
- Sijbers J, Van Audekerke J, Verhoye M, Van der Linden A, and Van Dyck D (2000). Reduction of ECG and gradient related artifacts in simultaneously recorded human EEG/fMRI data. *Magn Reson Imaging* 18, pp. 881–886.
- Smith AT, Williams AL, and Singh KD (2004). Negative BOLD in the visual cortex: evidence against blood stealing. *Hum Brain Mapp* 21, pp. 213–220.
- Smith SP, Fox PT, KL Miller, Glahn DC, Mickle Fox P, Mackay CE, Fillippini N, Watkins KE, Toro R, Laird AR, and Beckmann CF (2009). Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci U S A* 106, pp. 13040–13045.
- Spencer SS, Sperling MR, Shewmon DA, and Kahane P (2008). Intracranial Electrodes. In: *Epilepsy: A comprehensive Textbook*. Ed. by Engel J. and Pedley T.A. 2nd ed. Lippincott Williams & Wilkins. Chap. 171.
- Sperling MR and Clancy RR (2008). Ictal Electroencephalography. In: *Epilepsy: a comprehensive textbook*. Ed. by Engel J. and Pedley T.A. 2nd ed. Vol. 1. Lippincott Williams & Wilkins. Chap. 74.
- Srivastava G, Crottaz-Herbette S, Lau KM, Glover GH, and Menon V (2005). ICA-based procedures for removing ballistocardiogram artifacts from EEG data acquired in the MRI scanner. *Neuroimage* 24, pp. 50–60.
- Stam CJ (2005). Nonlinear dynamical analysis of EEG and MEG: review of an emerging field. *Clin Neurophysiol* 116.10, pp. 2266–301.
- Stern JM (2006). Simultaneous electroencephalography and functional magnetic resonance imaging applied to epilepsy. *Epilepsy Behav* 8.4, pp. 683–92.
- Sun L and Hinrichs H (2009). Simultaneously recorded EEG/fMRI: removal of gradient artifacts by subtraction of head movement related average artifact waveforms. *Hum Brain Mapp* 30, 3361–3377.
- Thomas CG, Harshman RA, and Menon RS (2002). Noise reduction in BOLD-based fMRI using component analysis. *Neuroimage* 17.3, pp. 1521–1537.
- Thornton R, Laufs H, Rodionov R, Cannadathu S, Carmichael DW, Vulliemoz S, Salek-Haddadi A, McEvoy AW, Smith SM, Lhatoo S, Elwes RD, Guye M, Walker MC, Lemieux L, and Duncan JS (2010a). EEG correlated functional MRI and postoperative outcome in focal epilepsy. *J Neurol Neurosurg Psychiatry* 81.8, pp. 922–7.
- Thornton R, Vulliemoz S, Rodionov R, Carmichael DW, Chaudhary UJ, Diehl B, Laufs H, Vollmar C, McEvoy AW, Walker MC, Bartolomei F, Guye M, Chauvel P, Duncan JS, and Lemieux L (2011). Epileptic networks in focal cortical dysplasia revealed using electroencephalography-functional magnetic resonance imaging. *Ann Neurol* 70.5, pp. 822–37.
- Thornton RC, Rodionov R, Laufs H, Vulliemoz S, Vaudano A, Carmichael D, Cannadathu S, Guye M, McEvoy A, Lhatoo S, Bartolomei F, Chauvel P, Diehl B, De Martino F, Elwes RD, Walker MC, Duncan JS, and Lemieux L (2010b). Imaging haemodynamic changes related to seizures: comparison of EEG-based general linear model, independent component analysis of fMRI and intracranial EEG. *Neuroimage* 53.1, pp. 196–205.
- Tohka J, Foerde K, Aron AR, Tom SM, Toga AW, and Poldrack RA (2008). Automatic independent component labeling for artifact removal in fMRI. *Neuroimage* 39.3, pp. 1227–1245.
- Tyvaert L, Hawco C, Kobayashi E, LeVan P, Dubeau F, and Gotman J (2008a). Different structures involved during ictal and interictal epileptic activity in malformations of cortical development: an EEG-fMRI study. *Brain* 131.Pt 8, pp. 2042–60.
- Tyvaert L, LeVan P, Grova C, Dubeau F, and Gotman J (2008b). Effects of fluctuating physiological rhythms during prolonged EEG-fMRI studies. *Clin Neurophysiol* 119.12, pp. 2762–2774.
- Tyvaert L, LeVan P, Dubeau F, and Gotman J (2009). Noninvasive dynamic imaging of seizures in epileptic patients. *Hum Brain Mapp* 30.12, pp. 3993–4011.
- Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B, and Joliot M (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15.1, pp. 273–89.

- Valente G, Esposito F, de Martino F, Goebel R, and Formisano E (2010). Using ICA for the analysis of fMRI data. In: *Simultaneous EEG and fMRI: recording, analysis, and application*. Ed. by Ullsperger M. and Debener S. Oxford University Press. Chap. 3.2, pp. 135–151.
- van Houdt PJ, de Munck JC, Zijlmans M, Huiskamp G, Leijten FS, Boon PA, and Ossenblok PP (2010a). Comparison of analytical strategies for EEG-correlated fMRI data in patients with epilepsy. *Magn Reson Imaging* 28.8, pp. 1078–86.
- van Houdt PJ, Ossenblok PP, Boon PA, Leijten FS, Velis DN, Stam CJ, and de Munck JC (2010b). Correction for pulse height variability reduces physiological noise in functional MRI when studying spontaneous brain activity. *Hum Brain Mapp* 31.2, pp. 311–25.
- van Houdt PJ, Ossenblok PP, van Erp MG, Schreuder KE, Krijn RJ, Boon PA, and Cluitmans PJ (2011). Automatic breath-to-breath analysis of nocturnal polysomnographic recordings. *Med Biol Eng Comput* 49.7, pp. 819–30.
- van Houdt PJ, Ossenblok PP, Colon AJ, Boon PA, and de Munck JC (2012). A framework to integrate EEG-correlated fMRI and intracerebral recordings. *Neuroimage* 60.4, pp. 2042–53.
- van Houdt PJ, Ossenblok PPW, Colon AJ, Leijten FSS, Verdaasdonk R, Boon PAJM, and de Munck JC (in prep). An explorative study to the existence of an epileptic resting-state fMRI network.
- van Houdt PJ, de Munck JC, Leijten FSS, Huiskamp GJM, Colon AJ, Boon PAJM, and Ossenblok PPW (submitted). Understanding EEG-fMRI correlation patterns in presurgical evaluation of focal epilepsy.
- Van der Meer JN, Tijssen MAJ, Bour LJ, Van Rootselaar AF, and Nederveen AJ (2010). Robust EMG-fMRI artifact reduction for motion (FARM). *Clin Neurophysiol* 121.5, pp. 766–776.
- Vincent JL, Larson-Prior LJ, Zempel JM, and Snyder AZ (2007). Moving GLM ballistocardiogram artifact reduction for EEG acquired simultaneously with fMRI. *Clin Neurophysiol* 118, pp. 981–998.
- Voets NL, Adcock JE, Stacey R, Hart Y, Carpenter K, Matthews PM, and Beckmann CF (2009). Functional and structural changes in the memory network associated with left temporal lobe epilepsy. *Hum Brain Mapp* 30.12, pp. 4070–81.
- Vulliemoz S, Thornton R, Rodionov R, Carmichael DW, Guye M, Lhatoo S, McEvoy AW, Spinelli L, Michel CM, Duncan JS, and Lemieux L (2009). The spatio-temporal mapping of epileptic networks: combination of EEG-fMRI and EEG source imaging. *Neuroimage* 46.3, pp. 834–43.
- Vulliemoz S, Rodionov R, Carmichael DW, Thornton R, Guye M, Lhatoo SD, Michel CM, Duncan JS, and Lemieux L (2010). Continuous EEG source imaging enhances analysis of EEG-fMRI in focal epilepsy. *Neuroimage* 49.4, pp. 3219–29.
- Vulliemoz S, Carmichael DW, Rosenkranz K, Diehl B, Rodionov R, Walker MC, McEvoy AW, and Lemieux L (2011). Simultaneous intracranial EEG and fMRI of interictal epileptic discharges in humans. *Neuroimage* 54.1, pp. 182–190.
- Waites AB, Shaw ME, Briellmann RS, Labate A, Abbott DF, and Jackson GD (2005). How reliable are fMRI-EEG studies of epilepsy? A nonparametric approach to analysis validation and optimization. *Neuroimage* 24.1, pp. 192–9.
- Walczak TS, Jayakar P, and Maizrahi EM (2008). Interictal Electroencephalography. In: *Epilepsy: a comprehensive textbook*. Ed. by Engel J. and Pedley T. 2nd ed. Vol. 1. Lippincott Williams & Wilkins. Chap. 73.
- Walker MC, Chaudhary UJ, and Lemieux L (2010). EEG-fMRI in adults with focal epilepsy. In: *EEG-fMRI: Physiological Basis, Technique, and Applications*. Ed. by Mulert C and Lemieux L. Springer-Verlag. Chap. 15, pp. 309–331.
- Wan X, Iwata K, Riera J, Kitamura M, Ozai T, Kitamura M, and Kawashima R (2006). Artifact reduction for EEG/fMRI recording: Nonlinear reduction of ballistocardiogram artifacts. *Clin Neurophysiol* 117.3, pp. 668–680.
- Ward JH (1963). Hierarchical grouping to optimize an objective function. *J Am Stat Assoc* 58, pp. 236–244.
- Wendling F, Bartolomei F, Bellanger JJ, and Chauvel P (2001). Interpretation of interdependencies in epileptic signals using a macroscopic physiological model of the EEG. *Clin Neurophysiol* 112.7, pp. 1201–18.
- Westmijse I, Ossenblok P, Gunning B, and van Luijtelaar G (2009). Onset and propagation of spike and slow wave discharges in human absence epilepsy: A MEG study. *Epilepsia* 50.12, pp. 2538–2548.

- Windischberger C, Langenberger H, Sycha T, Tschernko EM, Fuchsjaeger-Mayerl G, Schmetterer L, and Moser E (2002). On the origin of respiratory artifacts in BOLD-EPI of the human brain. *Magn Reson Imaging* 20.8, pp. 575–82.
- Worrell G (2012). High-frequency oscillations recorded on scalp EEG. *Eng. Epilepsy Curr* 12.2, pp. 57–58.
- Wu H, Rubinstein M, Shih E, Guttag J, Durand F, and Freeman W (2012). Eulerian video magnification for revealing subtle changes in the world. *ACM Trans Graph (Proceedings SIGGRAPH 2012)* 31.
- Yan WX, Mullinger KJ, Geirsdottir GB, and Bowtell R (2010). Physical modeling of pulse artefact sources in simultaneous EEG/fMRI. *Eng. Hum Brain Mapp* 31.4, pp. 604–620.
- Zahneisen B, Grotz T, Lee KJ, Ohlendorf S, Reisert M, Zaitsev M, and Hennig J (2011). Three-dimensional MR-encephalography: fast volumetric brain imaging using rosette trajectories. *Magn Reson Med* 65, pp. 1260–1268.
- Zhang Z, Lu G, Zhong Y, Tan Q, Yang Z, Liao W, Chen Z, Shi J, and Liu Y (2009a). Impaired attention network in temporal lobe epilepsy: a resting fMRI study. *Neurosci Lett* 458.3, pp. 97–101.
- Zhang Z, Lu G, Zhong Y, Tan Q, Liao W, Chen Z, Shi J, and Liu Y (2009b). Impaired perceptual networks in temporal lobe epilepsy revealed by resting fMRI. *J Neurol* 256.10, pp. 1705–13.
- Zijlmans M, Huiskamp G, Hersevoort M, Seppenwoolde JH, van Huffelen AC, and Leijten FS (2007). EEG-fMRI in the preoperative work-up for epilepsy surgery. *Brain* 130.Pt 9, pp. 2343–53.
- Zijlmans M, Jiruska P, Zelmann R, Leijten FSS, Jefferys JG, and Gotman J (2012). High-frequency oscillations as a new biomarker in epilepsy. *Eng. Ann Neurol* 71.2, pp. 169–178.

List of Abbreviations

AAL	anatomic automatic labeling
AEDs	anti-epileptic drugs
BOLD	blood oxygenation level-dependent
CBF	cerebral blood flow
CBV	cerebral blood volume
CMRO₂	cerebral metabolic rate for oxygen
CT	computed tomography
ECG	electrocardiogram
ECoG	electrocorticography
EEG	electroencephalography
EPI	echo planar imaging
fMRI	functional magnetic resonance imaging
FDR	false discovery rate
FIR	finite impulse response model
FOV	field of view
GLM	general linear model
HFO	high frequency oscillations
HRF	hemodynamic response function
IC	independent component
ICA	independent component analysis
IED	interictal epileptiform discharge
IRF	impulse response function
MEG	magnetoencephalography
MRI	magnetic resonance imaging

PAA	period amplitude analysis
PET	positron emission tomography
PPG	photoplethysmogram
RF	radiofrequency
RVT	respiratory volume per time
SEEG	stereoelectroencephalography
SPECT	single-photon emission computed tomography
TR	repetition time
VIPH	variation in pulse height

Summary

Epilepsy is a neurological disease characterized by epileptic seizures. In the Netherlands about 120.000 persons have epilepsy. Most of them can be treated successfully with anti-epileptic drugs. However, 20 - 30% of the patients does not respond to medication. These patients are potential candidates for epilepsy surgery, which aims to remove those brain regions that are responsible for the generation of seizures (i.e. epileptogenic zone). To create hypotheses about the location and extent of the epileptogenic zone several (non)invasive techniques are combined, ranging from electroencephalography (EEG) and magnetic resonance imaging (MRI) to neuropsychological tests.

In this context, we investigated the added value of a relatively new technique: simultaneous EEG and functional MRI (EEG-fMRI). EEG and fMRI are both techniques to measure neuronal activity. EEG measures electrical activity through electrodes at the scalp and is able to indicate the occurrence of epileptic activity [seizures or interictal epileptiform discharges (IEDs)] with good temporal resolution. However, the spatial resolution is relatively low, because the EEG signals represent a mixture of activity from different sources. fMRI, on the other hand, has a good spatial resolution. This technique provides an indirect measure for neuronal activity, which is based on changes in the ratio of oxy- and deoxyhemoglobin in the blood related to the occurrence of neuronal activity. Co-registration of EEG is necessary to determine whether changes in fMRI are related to epileptic activity. The epileptic events that are visible in the EEG during scanning are correlated with the fMRI time series yielding a correlation pattern indicating which brain regions were significantly more active during IEDs than in periods without visible IEDs in the EEG. The combination of EEG and fMRI provides a unique way to investigate epileptic networks with a high spatial precision.

The technical problems involved with the recording of EEG in the MRI environment are largely solved; under certain conditions and with special equipment it is safe to record EEG inside the MRI. However, the analysis and interpretation of the data are not straightforward. Therefore, the general aim of this thesis was to improve and to evaluate the potential of EEG-fMRI as a noninvasive technique in the presurgical evaluation of patients with medically intractable epilepsy.

Chapter 2 and 3 focus on the analysis of the fMRI data. In chapter 2 different analytical strategies for the correlation of EEG and fMRI were compared. The results suggest that the yield of EEG-fMRI is increased when a flexible approach is used, in which no assumptions are made about the precise underlying relation between EEG and fMRI, the hemodynamic response function (HRF). Since the HRF can vary over subjects and brain regions, the use of a more flexible approach, where the HRF is estimated from the data, increases the yield of fMRI.

Chapter 3 demonstrates that it is also important to correct the fMRI signals for variations in physiological functioning. These variations can be measured by a pulse oximeter at one of the subject's fingers. The variations in pulse height (VIPH) are related to respiration and cardiac output. VIPH appears to explain a large part of the fMRI signals, both in healthy volunteers and patients with epilepsy. Therefore, it is important to correct for these effects, especially when the subject has an abnormal breathing pattern during the experiment, as one of the patients with epilepsy in our study. When this physiological effect is taken into account the correlation pattern reduces from a very widespread correlation pattern to a pattern with a maximal BOLD response in the right temporal lobe. This region coincided with the resection area. The inclusion of VIPH regressors in the correlation model is not only important for the analysis of EEG-fMRI data of patients with epilepsy, but is relevant in general for resting-state fMRI studies, during which spontaneous brain activity is measured.

The yield of the EEG-fMRI analysis is dependent on the IEDs that are visible in the EEG. The detection of epileptic events is hampered by the presence of gradient and pulse artifacts in the EEG. Gradient artifacts result from switching of the gradients for slice selection, whereas pulse artifacts are related to the pulsation of the heart. Both types of artifacts can be removed off-line from the EEG with dedicated algorithms. Most of these algorithms are based on the average artifact subtraction principle, in which a template is created by averaging the artifact that is subsequently subtracted from the raw EEG data. In chapter 4 extensions of this approach are presented to further improve the correction of the EEG data. First, it appears that the gradient artifacts may differ slightly between scans. By clustering the artifacts and creating a template for each cluster, the correction was improved. Second, the pulsation artifacts appear to overlap from heartbeat to heartbeat. Therefore, the correction of the EEG data is improved when this overlap is taken into account.

Chapter 5 and 6 focus on the evaluation of the EEG-fMRI results. Most EEG-fMRI analyses yield a correlation pattern consisting of multiple activated brain regions. Our hypothesis was that these regions form an epileptic network including the onset and propagation of epilepsy related activity. For validation, the EEG-fMRI findings were compared to the results of invasive EEG data. These data are acquired either by depth electrodes implanted inside the brain tissue or by electrode grids placed on the subdural surface of the brain. In chapter 5 a semi-automatic procedure for the analysis of the invasive data is presented such that the EEG-fMRI findings can be

compared systematically with the invasive EEG data. This method was tested for five patients who were implanted with depth electrodes. In chapter 6 the application of the same method was described for the EEG data of sixteen patients recorded from subdural grids. In all data sets there was at least one region that was concordant with active invasive electrodes during IEDs. Furthermore, the EEG-fMRI results were compared to the location of the seizure onset zone and resection area: the EEG-fMRI correlation pattern included the seizure onset zone in 83 % of the data sets and the resection area in 93 % of the data sets. For some patients, we were able to validate an epileptic network of onset and propagation regions.

Although in most patients relevant results are obtained with EEG-fMRI, in some patients no IEDs are detected in the EEG during scanning. This may have several explanations, for instance the limited scanning period (about 45 minutes) or an origin located in deep lying structures. When no IEDs are present, the data cannot be analyzed with the current EEG-fMRI approach. Chapter 7 is an explorative study to the use of independent component analysis (ICA), a data-driven approach independent of the EEG. The preliminary results suggest that it is possible to find epileptic networks with ICA, even in the absence of visible IEDs in the EEG. This finding provides further perspective for the use of resting-state fMRI in clinical practice.

In conclusion, the yield of EEG-fMRI increases with the methodological improvements presented in this thesis. Furthermore, the results of EEG-fMRI are clinically relevant, because EEG-fMRI is usually concordant with IEDs in the invasive EEG data and includes the seizure onset zone and resection area. For this reason, there is an important role for EEG-fMRI in the surgical planning, especially with regard to the determination of the implantation strategy. In comparison to other noninvasive techniques in the presurgical evaluation, EEG-fMRI will have an added value in patients in whom a focus from deeply lying structures or multifocal epilepsy is expected. In addition, EEG-fMRI is able to reveal the whole network from onset to propagation, which is more difficult to achieve with EEG alone.

Nederlandse Samenvatting

Co-registratie van EEG en functionele MRI een niet-invasieve techniek in de prechirurgische evaluatie van focale epilepsie

Epilepsie is een neurologische aandoening die wordt gekenmerkt door epileptische aanvallen, die het dagelijkse functioneren van de patiënt substantieel kunnen beïnvloeden. In Nederland hebben ongeveer 120.000 mensen epilepsie. Twintig tot dertig procent van de patiënten reageert onvoldoende op medicatie. Zij kunnen in aanmerking komen voor een operatie waarbij het hersengebied verwijderd wordt dat cruciaal is voor het veroorzaken van aanvallen, het epileptogene gebied. Om de precieze locatie van het epileptogene gebied te bepalen, ondergaat de patiënt voorafgaand aan deze operatie een breed scala aan onderzoeken, van electroencefalografie (EEG) en magnetic resonance imaging (MRI) tot aan invasieve EEG onderzoeken.

In dit proefschrift wordt de toegevoegde waarde onderzocht van een relatief nieuwe techniek, namelijk de co-registratie van EEG en functionele MRI (EEG-fMRI). EEG en fMRI zijn beiden technieken om hersenactiviteit te meten. Met EEG wordt elektrische neuronale activiteit met behulp van elektroden geplaatst op de schedel. Op dit moment is EEG de belangrijkste techniek voor de diagnose van epilepsie. EEG is in staat om epileptische activiteit (aanvallen of interictale epileptische ontladingen) met een goede temporele precisie te meten. De spatiële resolutie is echter relatief laag, omdat het EEG een weergave is van verschillende bronnen. fMRI heeft juist een zeer goede spatiële resolutie. Deze techniek meet hersenactiviteit gebaseerd op veranderingen in zuurstofconcentratie in het bloed, die het gevolg zijn van zuurstofopname na neuronale activatie. Co-registratie van EEG is nodig om te controleren of de waargenomen veranderingen in het fMRI signaal het gevolg zijn van epileptische ontladingen. Het EEG wordt dus alleen gebruikt om te bepalen op welke momenten er epileptische ontladingen plaats vindt tijdens het scannen. Deze momenten worden vervolgens gecorreleerd met de fMRI signalen. Het resultaat

is een correlatiepatroon dat aangeeft welke gebieden significant actief zijn tijdens epileptische ontladingen.

De technische problemen rondom het simultaan meten van EEG en fMRI zijn grotendeels opgelost; het is onder bepaalde condities en met speciale apparatuur veilig om EEG in een MRI scanner te meten. De analyse en interpretatie van de gegevens is echter niet eenvoudig. Het doel van dit proefschrift was dan ook om de analyse van EEG-fMRI data te optimaliseren en de resultaten te evalueren, zodat EEG-fMRI gebruikt kan worden als een niet-invasieve techniek in de prechirurgische evaluatie van patiënten die kandidaat zijn voor een operatie.

Hoofdstuk 2 en 3 van dit proefschrift richten zich op de analyse van de fMRI data. In hoofdstuk 2 zijn verschillende analytische strategieën met elkaar vergeleken. De resultaten laten zien dat de opbrengst van EEG-fMRI verhoogd wordt wanneer er in het model vooraf geen aannames gemaakt worden over de precieze onderliggende relatie tussen EEG en fMRI, de hemodynamische respons functie (HRF). Omdat de HRF verschillend kan zijn tussen patiënten en tussen hersengebieden, is een flexibel model beter dan een model waarin een vaste vorm voor de HRF wordt aangenomen.

In hoofdstuk 3 wordt aangetoond dat het belangrijk is om de fMRI signalen te corrigeren voor variaties in fysiologische functies. De veranderingen in fysiologische functies zijn te meten met een puls oximeter op een vinger van de proefpersoon. De verschillen in amplitude van dit signaal geven de veranderingen in de ratio van oxy- en deoxyhemoglobine weer als gevolg van ademhaling en hartslag. Deze veranderingen blijken een groot gedeelte van het fMRI signaal te verklaren, zowel bij gezonde proefpersonen als bij patiënten met epilepsie. Daarom is het belangrijk om te corrigeren voor de artefacten die gerelateerd zijn aan fysiologische functies. Deze bevinding is niet alleen van belang voor fMRI-onderzoek bij patiënten met epilepsie, maar ook voor fMRI-onderzoeken tijdens rusttoestand waarbij spontane hersenactiviteit gemeten wordt.

De sensitiviteit van de fMRI analyse is afhankelijk van de epileptische pieken die in het EEG gemarkeerd worden. De detectie van epileptische pieken wordt echter bemoeilijkt door de artefacten die in het EEG ontstaan door de schakelingen van de gradiënten en pulsaties van het hart. Beide typen artefacten kunnen met behulp van algoritmes achteraf gecorrigeerd worden. De meeste algoritmes maken een sjabloon van het artefact door middeling van diverse artefacten; vervolgens wordt dit sjabloon van de ruwe EEG-signalen afgetrokken. Hoofdstuk 4 bevat een voorstel voor uitbreiding van dit algoritme, zodat de correctie van het EEG verbeterd kan worden. De gradiënt-artefacten blijken niet zo stabiel te zijn tijdens lange registraties. Door de artefacten te clusteren en per cluster een sjabloon te maken, wordt de correctie beter. Verder werd aangetoond dat de pulsatie-artefacten van hartslag tot hartslag kunnen overlappen. Door hier rekening mee te houden wordt de correctie van het EEG beter.

In hoofdstuk 5 en 6 ligt de nadruk op de evaluatie van de EEG-fMRI resultaten.

Het correlatiepatroon bestaat bij de meeste patiënten uit meerdere geactiveerde gebieden. De hypothese is dat deze gebieden een interictaal netwerk vormen van gebieden die het begin van de epileptische ontladingen weergeven en gebieden die gerelateerd zijn aan spreiding van deze activiteit over de cortex. Om dit te valideren, zijn de uitkomsten van EEG-fMRI vergeleken met intracraniële registraties. Bij deze registraties worden diepte-elektroden in de hersenen of matjes met elektroden op de hersenschors geplaatst. Deze invasieve registraties worden gezien als de gouden standaard om het begin van de aanvallen te lokaliseren. In hoofdstuk 5 beschrijven we een methode om de invasieve data te analyseren, zodat een systematische vergelijking met EEG-fMRI mogelijk is. Deze methode is geëvalueerd voor vijf patiënten die in het kader van hun operatie geregistreerd zijn met diepte-elektroden. Vervolgens is deze methode in hoofdstuk 6 toegepast op data van zestien patiënten waarbij elektrodenmatjes geplaatst zijn. De sensitiviteit van EEG-fMRI bleek hoog: voor bijna iedere patiënt kwam tenminste één gebied overeen met het gebied waar volgens de invasieve EEG data de interictale epileptische ontladingen beginnen. Daarnaast zijn de EEG-fMRI resultaten ook vergeleken met de klinische meer relevante informatie, zoals de plek waar de aanvallen beginnen en het gebied dat succesvol verwijderd is tijdens de operatie. Het EEG-fMRI correlatiepatroon liet tenminste één geactiveerd gebied zien dat overeenstemming vertoont met de locatie van het begin van de aanvallen in 83 % van de data sets en het geopereerde gebied in 93 % van de data sets. In sommige patiënten was het mogelijk om een epileptisch netwerk te valideren dat zowel het begin als de spreiding van epileptische ontladingen weergeeft.

Een nadeel van EEG-fMRI is dat de analyse sterk afhankelijk is van het aantal epileptische pieken die in het EEG gedetecteerd worden. Het komt regelmatig voor dat er geen pieken optreden tijdens het scannen door de relatief korte tijdspanne van het onderzoek (ongeveer 45 minuten). Ook kan het voorkomen dat de pieken niet door het EEG gezien worden, omdat ze in diep gelegen hersengebieden gegeneerd worden. Wanneer er geen pieken in het EEG te zien zijn, kan de data niet geanalyseerd worden met de hiervoor beschreven modelmatige aanpak. Hoofdstuk 7 beschrijft een exploratieve studie naar het gebruik van een data-gestuurde aanpak die onafhankelijk is van het EEG, namelijk independent component analyse (ICA). Het lijkt erop dat het mogelijk is om met ICA het epileptische netwerk te vinden, óók als er geen pieken in het EEG gevonden worden. Dit biedt perspectief voor het gebruik van fMRI in de prechirurgische evaluatie zonder noodzakelijke co-registratie van het EEG.

Tot slot, met dit onderzoek hebben we laten zien dat de opbrengst van EEG-fMRI onderzoeken hoger wordt met de door ons voorgestelde methodologische verbeteringen. Verder hebben we ook aangetoond dat de resultaten van EEG-fMRI relevant zijn voor de prechirurgische evaluatie. Omdat EEG-fMRI niet veel informatie mist (in bijna alle gevallen vond EEG-fMRI het gebied waar de aanvallen beginnen),

lijkt de methode zeer geschikt om te bepalen waar invasieve elektroden geplaatst moeten worden. Ten opzichte van andere onderzoeken heeft EEG-fMRI vooral een toegevoegde waarde voor de prechirurgische evaluatie van patiënten bij wie diep in de hersenen gelegen bronnen of multifocale epilepsie verwacht wordt. Verder kan met EEG-fMRI het hele netwerk dat betrokken is bij interictale epileptiforme ontladingen in kaart gebracht worden, iets wat veel moeilijker is met EEG alleen.

Dankwoord

Na ruim vier jaar is mijn proefschrift dan bijna af! Dit had ik niet kunnen bereiken zonder de begeleiding, ondersteuning en medewerking van anderen. Ik wil dan ook graag van deze gelegenheid gebruik maken om iedereen te bedanken.

Als eerste wil ik mijn beide copromotoren bedanken, Pauly Ossenblok en Jan C. de Munck. Ik heb ontzettend veel van jullie geleerd, zowel op inhoudelijk als persoonlijk vlak. Jullie waren letterlijk mijn dagelijkse begeleiders, want ik kon altijd bij jullie terecht met mijn vragen. Dat heb ik enorm gewaardeerd! Pauly, jij bent de initiator en kartrekker van het EEG-fMRI project. Bedankt dat je me overgehaald hebt om bij Kempenhaeghe te blijven. Ik heb veel geleerd van je doelgerichte manier van werken. Je hebt me laten inzien hoe ik onze resultaten zo gestructureerd en overzichtelijk mogelijk kon weergeven. Ik twijfel er niet aan dat ik daar in de toekomst nog veel profijt van zal hebben. Jan, bedankt voor je nuchterheid en optimisme van de afgelopen jaren! Jij ziet overal mogelijkheden en zit vol waardevolle ideeën. Het was erg prettig om met jouw zelfgeschreven software te mogen werken. Het was een voordeel dat het meteen aangepast of uitgebreid kon worden naar onze wensen.

Beste Prof. Boon, bedankt voor alle goede adviezen, vertrouwen en ondersteuning van de afgelopen jaren! Vooral uw correcties en klinische blik op de artikelen waren erg waardevol. Beste Prof. Stam, in de eerste jaren was u mijn promotor. Ik heb me er altijd over verbaasd hoe snel u artikelen kon lezen en van nuttig commentaar kon voorzien. Prof. Verdaasdonk, bedankt dat u het stokje van Prof. Stam hebt overgenomen in de laatste fase van mijn promotietraject en voor de mogelijkheid dat ik daarna in het VUmc aangesteld kon worden.

Beste Frans Leijten, ik wil je bedanken voor je enthousiasme en prettige samenwerking van de afgelopen jaren. Ik heb veel van u geleerd over de prechirurgische evaluatie. Vooral in het begin was het erg nuttig om mee te kunnen kijken bij de operaties. Voor hoofdstuk 6 ben ik vaak langs geweest om ECoG data te selecteren en te bespreken. Bedankt dat je ruim de tijd hebt genomen om mij de nodige informatie en uitleg te geven. Geertjan Huiskamp, ook jij bedankt voor de prettige samenwerking en de vlotte manier waarop data uitgewisseld kon worden. Maeike

Zijlmans, voor hoofdstuk 2 heb ik de data gebruikt van een van jouw artikelen. De avonden waarop we samen de resultaten doornamen waren nuttig en efficiënt.

Albert Colon, jij was van de neurologen van Kempenhaeghe het meest betrokken bij het project. Samen met Mike heb je heel wat uurtjes naar de EEGs gekeken om te bepalen of iets wel of geen piek was. Het was erg leerzaam om die discussies te volgen. Jouw enthousiasme voor een interessante casus of leuk resultaat werkt aanstekelijk. Heel veel succes met het afronden van je eigen promotie! Ook de overige artsen verspreid over het hele land wil ik bedanken voor hun betrokkenheid en doorverwijzing van patiënten, met name Louis Wagner, Cyrille Ferrier, Demetrios Velis en Johan Arends.

Beste Roy Krijn, bedankt voor je hulp bij alle matlab, computer en MRI trigger problemen. Ik weet niet wat we wat dat betreft zonder jou hadden moeten doen. Peter Bijkerk, bedankt voor de technische ondersteuning bij het verbeteren en testen van de EEG-fMRI metingen. Bij deze wil ik ook Jan Verwoerd van Philips Healthcare bedanken voor zijn bijdrage aan de ontwikkeling van de scan protocollen.

Zonder de ervaren hulp van de laboranten van de KNF en MRI van Kempenhaeghe hadden we de EEG-fMRI metingen niet kunnen uitvoeren. Met name Mike van der Mierden, Marlies Dolmans en Ine Keulen hebben een grote bijdrage geleverd bij het plaatsen van de EEG cap en de analyse van de EEG gegevens. Mike, je hebt me laten inzien dat het beoordelen van een EEG niet zo zwart-wit is als ik graag zou willen. Remco Berting en Esther Peeters ben ik dankbaar voor hun hulp tijdens het scannen. Harrie Geeris wil ik bedanken voor de selectie van de SEEG data voor hoofdstuk 5. De secretaresses van de KNF, Maike van der Steen en Ivanka van Wanrooy, bedankt voor de ondersteuning bij de selectie van de patiënten.

Bij deze wil ik ook alle patiënten en vrijwilligers bedanken voor hun deelname aan het onderzoek. Zonder hen had ik niet zulke mooie plaatjes in mijn proefschrift gehad.

Alle leden van de lees en/of promotiecommissie, prof.dr. Frederik Barkhof, prof.dr. Jean Gotman, prof.dr. Bart ter Haar Romeny, prof.dr. Jan Heimans, dr. Frans Leijten, prof.dr. Fernando Lopes da Silva, prof.dr.ir. Natasha Maurits, prof.dr. Philip Scheltens en prof.dr. Stefan Sunaert, ben ik dankbaar voor de tijd en energie die zij gestoken hebben in de beoordeling van mijn proefschrift. Jean Gotman, thank you for giving me the opportunity to visit your lab. Although only for a short period, I experienced a very warm welcome.

Mijn collega's van de Boondert, eerst Tamara en Saskia, daarna Sylvie, Evelien, Reina, Geke, Maartje, Debby S, Debby K, Zita, Dominique en Janneke. Veel vrouwen bij elkaar en daarom werd er veel gekletst en gelachen. Het was fijn dat ik voor zowel serieuze als minder-serieuze gesprekken bij jullie terecht kon! De gezellige etentjes en (al dan niet educatieve) activiteiten zal ik missen!

Beste Corry Bos van Creatief Roermond, bedankt voor je creatieve input bij het maken van de kafft!

Lieve familie en vrienden, bedankt voor jullie interesse in mijn onderzoek én voor de nodige afleiding. Ook al gaan we nu verhuizen, ik hoop niet dat dit verandering brengt in alle gezellige thee-drink momenten, haakmiddagen, etentjes, feestjes, uitstapjes, en weekendjes weg, want daar kijk ik altijd naar uit! Als laatste in de rij van promoties van de TU-vriendinnengroep, heb ik al veel goede voorbeelden gehad! Ik kan alleen maar wensen dat ik het er net zo goed vanaf breng. Linda, ik vind het leuk om te zien hoe jij nu ook met wetenschap bezig bent. Veel succes met het afronden van je master. Lieve papa en mama, jullie wil ik in het bijzonder bedanken voor de liefde en mogelijkheden die jullie ons altijd gegeven hebben! Jullie onvoorwaardelijke steun betekent veel voor mij.

Lieve Davy, ik zal je deze keer niet vergeten te noemen! Maar toch denk ik dat er niet genoeg woorden zijn om mijn dank uit te drukken. Je hebt me ontzettend geholpen, zowel op professioneel als op persoonlijk vlak. En ik hoop dat je zonder dit dankwoord ook wel weet hoeveel dat voor me betekent. Hopelijk kan ik je net zoveel kan bijstaan met je eigen promotie!

Liefs,
Petra
Oktober 2012

List of Publications

Papers

van Houdt PJ, de Munck JC, Zijlmans M, Huiskamp G, Leijten FS, Boon PA, and Ossenblok PP (2010a). Comparison of analytical strategies for EEG-correlated fMRI data in patients with epilepsy. *Magn Reson Imaging* 28.8, pp. 1078–86

van Houdt PJ, Ossenblok PP, Boon PA, Leijten FS, Velis DN, Stam CJ, and de Munck JC (2010b). Correction for pulse height variability reduces physiological noise in functional MRI when studying spontaneous brain activity. *Hum Brain Mapp* 31.2, pp. 311–25

van Houdt PJ, Ossenblok PP, van Erp MG, Schreuder KE, Krijn RJ, Boon PA, and Cluitmans PJ (2011). Automatic breath-to-breath analysis of nocturnal polysomnographic recordings. *Med Biol Eng Comput* 49.7, pp. 819–30

van Houdt PJ, Ossenblok PP, Colon AJ, Boon PA, and de Munck JC (2012). A framework to integrate EEG-correlated fMRI and intracerebral recordings. *Neuroimage* 60.4, pp. 2042–53

van Houdt PJ, de Munck JC, Leijten FSS, Huiskamp GJM, Colon AJ, Boon PAJM, and Ossenblok PPW (submitted). Understanding EEG-fMRI correlation patterns in presurgical evaluation of focal epilepsy

van Houdt PJ, Ossenblok PPW, Colon AJ, Leijten FSS, Verdaasdonk R, Boon PAJM, and de Munck JC (in prep). An explorative study to the existence of an epileptic resting-state fMRI network

de Munck JC, van Houdt PJ, Verdaasdonk RM, and Ossenblok PP (2012). A semi-automatic method to determine electrode positions and labels from gel artifacts in EEG/fMRI-studies. *Neuroimage* 59.1, pp. 399–403

de Munck JC, van Houdt PJ, Gonçalves SI, van Wegen E, and Ossenblok PPW (2013). Novel artefact removal algorithms for co-registered EEG/fMRI based on selective averaging and subtraction. *Neuroimage* 64, pp. 407–415

Book Chapters

de Munck JC, Gonçalves SI, van Houdt PJ, Mammoliti R, Ossenblok P, and Lopes da Silva FH (2010). The hemodynamic response of EEG features. In: *EEG-fMRI: Physiological basis, technique, and applications*. Ed. by Mulert C and Lemieux L. Springer-Verlag. Chap. 3.6, pp. 195–208

Curriculum Vitae

Petra van Houdt was born in Hulst on April 11, 1983. After finishing her secondary education at the Reynaertcollege (previously known as St. Jansenius) in 2001, she studied biomedical engineering at the Eindhoven University of Technology. During her studies, she carried out an internship at the department of physiology of the Semmelweis University (Budapest) investigating the adaptation of the intact rat heart to changes in workload. She performed her graduation project at the department of clinical physics of Kempenhaeghe, working on an algorithm for automatic detection of obstructive sleep disordered breathing. After her graduation in 2007, she worked as a junior researcher in Kempenhaeghe to evaluate and complete this algorithm and to publish the results. At the end of 2007, she started her PhD project at Kempenhaeghe and VU University Medical Center with a focus on epilepsy. The results of the research are described in this thesis, entitled "Simultaneous EEG and functional MRI: a noninvasive tool in the presurgical evaluation of focal epilepsy". In this period, she visited the laboratory of Prof.dr. Jean Gotman of the Montreal Neurological Institute. Currently she is working as a researcher on a follow-up study that focuses on the development of more data-driven analytical approaches.