


# PART 3



**Measurement Issues of physical activity and  
sedentary behavior**



# 5

## **Study protocol of physical activity and sedentary behavior measurement among school-children by accelerometry – Cross-sectional survey as part of the ENERGY project**

Mine Yıldırım, Maïté Verloigne, Ilse de Bourdeaudhuij,  
Odysseas Androutsos, Yannis Manios, Regina Felsó,  
Éva Kovács, Alain Doessegger, Bettina Bringolf-Isler,  
Saskia J te Velde, Johannes Brug, Mai JM Chinapaw  
*BMC Public Health 2011;11:182*

## **Abstract**

**Background:** Physical activity and sedentary behavior among children should be measured accurately in order to investigate their relationship with health. Accelerometry provides objective and accurate measurement of body movement, which can be converted to meaningful behavioral outcomes. The aim of this study was to evaluate the best evidence for the decisions on data collection and data processing with accelerometers among children resulting in a standardized protocol for use in the participating countries.

**Methods/Design:** This cross-sectional accelerometer study was conducted as part of the European ENERGY-project that aimed to produce an obesity prevention intervention among schoolchildren. Five countries, namely Belgium, Greece, Hungary, Switzerland and the Netherlands participated in the accelerometer study. We used three different Actigraph models-Actitrainers (triaxial), GT3Xs and GT1Ms. Children wore the device for six consecutive days including two weekend days. We selected an epoch length of 15 seconds. Accelerometers were placed at children's waist at the right side of the body in an elastic belt.

**Results:** In total, 1082 children participated in the study (mean age =  $11.7 \pm 0.75$ y, 51% girls). Non-wearing time was calculated as periods of more than 20 minutes of consecutive zero counts. The minimum daily wearing time was set to 10 hours for weekdays and 8 hours for weekend days. The inclusion criterion for further analysis was having at least three valid weekdays and one valid weekend day. We selected a cut-point (count per minute (cpm)) of <100 cpm for sedentary behavior, <3000 cpm for light, <5200 cpm for moderate, and >5200 cpm for vigorous physical activity. We also created time filters for school-time during data cleaning in order to explore school-time physical activity and sedentary behavior patterns in particular.

**Discussion:** This paper describes the decisions for data collection and processing. Use of standardized protocols would ease future use of accelerometry and the comparability of results between studies.

## Introduction

The accurate measurement of physical activity (PA) and sedentary behavior is essential to evaluate their health impact in terms of amount and intensity (1,2). The complex nature of PA that covers both sports and non-sport activities makes it complicated to perfectly measure all of its aspects. Ideal measurement should be applied in daily life, lasting long enough to represent habitual activity patterns and with low inconvenience to the participant (3). Self-reported measures may not accurately reflect free-living activity patterns due to recall bias and/or its susceptibility to reporting bias by social desirability (4,5). Furthermore, a recent systematic review reported that there is currently no PA questionnaire for youth with both acceptable reliability and validity (6).

Apart from the importance of PA to maintain good health, there is growing evidence that excessive sedentary behavior is linked to a range of health problems in particular overweight and obesity (7-9). Sedentary behaviors such as sitting and screen-based entertainment include activities that do not increase energy expenditure substantially above resting level (10). Since sedentary behavior such as TV viewing has a strong habitual element (a kind of automatic behavior that is done without thinking) it is necessary to measure it over a period of time (11). It has been shown that using single-item self-reports or selecting only television viewing as an indicator of sedentary behavior does not estimate the broad range of sedentary behaviors in which children participate (12,13).

Over the last decades, motion sensors have been widely used as objective measures of PA as well as sedentary behaviors to overcome the problems related to self-report methods (14). The recent technological advances increase the capabilities of motion sensors such as accelerometers, which are small and light devices that are able to collect data for weeks (3). Accelerometers measure accelerations caused by body movements in one to three orthogonal planes (vertical, mediolateral and anteroposterior) and basically generate the activity counts and minutes spent above predefined thresholds (15). Using accelerometers allows researchers to focus on activity patterns rather than energy expenditure (16). In addition, they provide data to further investigate the dose-response relationship between activity patterns and health and providing evidence to be used in public health recommendations (17).

Recent reviews reported that accelerometers provide an accurate and precise measurement of all intensity levels of PA with the entire range from sedentariness to very vigorous (11,15,18). Distinguishing light activity and sedentary behavior is important due to the contribution of total daily light activity to energy expenditure. Accelerometers can make this differentiation that eases to study the relationship between sedentary behavior and health (10). A central issue in using accelerometers is careful planning of data collection, data analysis and interpretation (15). Unfortunately to date there is no consensus on the data analysis and interpretation of accelerometers. In 2004, an international conference was held to close the knowledge gap in the science of accelerometry. As a result, the best practice recommendations were derived on accelerometer use but still there was no clear consensus (19). For this reason, unresolved issues in methodology and data processing should be taken into account in the research planning phase. Therefore, the current paper aims to critically appraise the best evidence

for the decisions on data collection and data processing with accelerometers among children. With this protocol we also aimed to standardize the implementation of the accelerometer study between participating countries.

In the "European Energy balance Research to prevent excessive weight Gain among Youth" (ENERGY)-project, we objectively assessed PA and sedentary time by accelerometers in a subsample of about 1000 children aged 10-12 from five European countries. This paper describes the protocol of the data collection, processing and analysis.

## **Methods**

### **Project overview**

The ENERGY-project is a cross-European project aimed at the systematic development of a cross-European school-based and family involved intervention program to prevent overweight among children. The project is carried out by a multidisciplinary team of researchers from eleven European countries and Australia. The design and conceptual framework of the ENERGY-project have been previously described (20). As part of the development of this obesity prevention program, cross-sectional school-based surveys were conducted among 10-12 year old children and their parents in eight European countries (Belgium, Greece, Norway, Hungary, Slovenia, Spain, Switzerland and the Netherlands). The aim of the survey was to provide up to date information on the prevalence of overweight and obesity - measured by objective measures of height, weight and waist circumference-, on the most important energy balance related behaviors (EBRBs) and their social, cognitive and school environmental determinants. Schools were employed in each country to reach 1000 eligible children. The schools were randomly selected concerning the degree of urbanization of the different provinces and the socioeconomic status (SES) of the different areas within the selected provinces. All participating countries obtained ethical committee approvals. The details on the survey are described elsewhere (21).

### **Accelerometry study**

In five countries -Belgium, Greece, Hungary, Switzerland and the Netherlands- accelerometer data were collected in a subsample of approximately 200 children per country. The selection of schools was balanced across the participating provinces selected for the main survey. The distribution of the number of selected schools for accelerometry was proportional to the number of schools in the selected cities (thus more schools from larger cities). One identical recruitment letter was sent to all randomly selected schools explaining that in some schools children would be asked to wear an accelerometer during six consecutive days. The selection of the schools for accelerometer data collection was based on the logistic planning of the data collection process as well as willingness of the schools. Trained researchers collected accelerometer data according to a standardised protocol.

### **Type of accelerometers**

We used three different models of Actigraph (Pensacola, FL) accelerometers namely Actitrainers (triaxial), GT3Xs and GT1Ms. The GT3X and GT1M (dimensions: 3.8 cm × 3.7 cm × 1.8 cm) and Actitrainer (dimensions: 8.6 cm × 3.3 cm × 1.5 cm) are lightweight devices weighing 27, 27 and 51 grams, respectively (22). A recent study confirmed that vertical axis output of GT1M and GT3X was similar (23). Actigraph accelerometers have shown adequate reproducibility, validity and feasibility in children and adolescents (18).

### **Epoch length and duration of measurement**

The accelerometer collects and sums the activity counts over a user-defined epoch (24). Bailey et al. (25) reported that children tend to have very short bursts of high intensity activities. In their study sample, the median duration of a high intensity activity was 3 seconds with 95% lasting <15 seconds. Therefore, to be able to accurately capture the pattern of high intensity and short duration PA, we selected a time interval/epoch length of 15 seconds.

The duration of the measurement should reflect habitual activity and variability in activity and sedentary behavior patterns from day to day (15). To acquire a valid representation of activity for children and adolescents four to nine days of monitoring are needed. Due to differences between weekday and weekend activity patterns, it is advised to combine weekdays and weekend days in measurement protocols (24). Based on these findings, we collected data during six consecutive days including two weekend days.

### **Initialising the devices**

The ActiLife software (at least version 4.3.0) was used to initialise the accelerometers and upload the collected data (26). Before initialising, the devices were fully charged by connecting the device to a standard USB port. The start date was set at the day when the devices were handed out to the children starting at 05.00 AM. We did not select a stop time. For optional modes dual axis and third axis were selected for GT3X and Actitrainers additional to activity mode.

### **Placement of devices**

Godfrey et al. (27) emphasized the necessity of studying 'whole body' movements presented by placement of a motion sensor as close as possible to the centre of mass of the body, such as the waist. Children were fitted with accelerometers located at the waist at the right side of the body in an elastic belt. Children were asked to wear the device either under or on their clothes during all waking hours, apart from water-based activities. We marked the top of the device with a sticker and instructed the children to be aware that the accelerometer should be placed with the sticker on top.

### **Diaries (log-books)**

We asked children to fill in a diary recording the time of getting up in the morning and going to bed for sleeping. They also noted the time and reason why the device was removed for 5 minutes or more for any activity such as swimming, or showering. We also

asked whether the measurement day was a school day, half-school day or non-school day (See additional file 1: Diary).

### Pre-testing the protocol

The researchers who were responsible for the accelerometer data collection were trained to work according to a standard protocol. The participating countries pre-tested the protocol with at least five 10-12 year old children for feasibility. Based on the pre-test results, some changes were made in the layout of the diary. We emphasized that children need to fill in the diaries themselves and that the devices should also be worn during all activities except water activities.

### Data collection

Researchers distributed the accelerometers face-to-face at schools. Information about accelerometer use was given to the children orally and at the end of the information session accelerometers were handed out. Researchers placed the accelerometer to the children's waist. Additionally, children and parents received a brochure about accelerometer use including the instructions for children (See additional file 2: instructions and information brochure) and the diary. Teachers were also informed about the procedure and asked to remind the children to wear the devices every day. Researchers called and asked the teachers one day before collecting the devices to remind children to bring the device and diary back to school. After the proposed wearing period, children brought the device back to school and handed it out to their teacher. Downloading the data from accelerometers was done as soon as possible on the same computer where it was initialised to prevent disturbances that can be caused by the time offset between computers. The diary data were entered manually in excel files.

Table 1 shows the characteristics of the children who participated in the accelerometer study per country. Sixty-seven percent of children (n = 722) had enough data to be included in further analyses.

**Table 1.** Characteristics of children who participated in the accelerometer study per country.

Country	Initial number	Final number*	Age <sup>a</sup> (mean±SD)	BMI <sup>a</sup> (mean±SD)	Gender <sup>a</sup> (girl%)
Belgium	196	107	11.4 (0.7)	18.0 (2.6)	55.7
Greece	215	160	11.3 (0.6)	20.9 (3.9)	53.8
Hungary	194	147	12.2 (0.7)	19.9 (3.3)	50.0
Switzerland	277	206	11.4 (0.9)	17.8 (2.8)	58.0
the Netherlands	200	102	11.8 (0.6)	18.6 (3.0)	43.4
<b>TOTAL</b>	<b>1082</b>	<b>722</b>	<b>11.7 (0.7)</b>	<b>18.9 (3.4)</b>	<b>53.2</b>

\* The number of children after data reduction, BMI; Body mass index, SD; Standard deviation

<sup>a</sup> Based on the final number of children

### Data processing

All data collected in the study were transferred to VU University Medical Center (VUmc), the Netherlands for data processing. Meterplus (28) was used for data reduction and

analysis. Meterplus is a Windows-based program developed by researchers from San Diego State University in partnership with Actigraph. The Meterplus program was used to prepare and clean the accelerometer data files according to non-wearing time (as described below), invalid data (i.e., days that have not enough wearing time and implausibly high count values) and specific activity bout definitions such as bouts of sedentary time.

### **Data reduction**

It is known that PA and sedentary time results may change substantially depending on how data is processed (29, Fischer et al. unpublished data). Therefore, the quality of the accelerometer data should be checked first using systematic data reduction procedures. The decisions on minimum daily wearing time and number of required days for data analysis are critical data reduction issues. Wearing time calculation is not solely excluding all zero count values from data, since sedentary behavior is part of data. For this reason decisions should be made regarding duration of consecutive zeros to distinguish between non-wearing and sedentary time (30). Non-wearing time is the time period that participants did not wear the device such as during sleeping or water sports/activities. In the current study non-wearing time was calculated as periods of more than 20 minutes of consecutive zero counts (31,32). Wearing time was calculated by subtracting non-wearing time from 24 hours (33). The minimum daily wearing time was set at 10 hours for weekdays and 8 hours for weekend days considering different sleep patterns at weekends. Another critical issue is to include enough days to be able to reflect children's habitual PA pattern. A valid day requires a minimum number of hours of wearing. Children who had at least three valid weekdays and one valid weekend day were included in the further data analysis (34).

### **Cut-points for activity categories**

The end results from accelerometer measurement are count values. Counts have been calibrated against energy expenditure in order to get a biological meaning (15). The most common way to estimate time spent in a specific activity category is to develop a regression equation that defines the relationship between counts and energy expenditure, allowing counts to be converted to units of energy expenditure. To date, a number of calibration equations for youth were developed for the Actigraph accelerometers, though the methods used were rather different (35). Table 2 shows the most common used cut-points that are appropriate for children aged 10-12.

Cut-points by Freedson et al. (36) are completely laboratory-based. Puyau et al. (37) and Treuth et al. (38) used more free living activities, ranging in intensity from sedentary to vigorous. Mattocks et al. (39) also used free living activities but only for moderate and vigorous activity intensities. Treuth et al. (38) measured energy expenditure at rest and during activity with portable indirect calorimetry and used resting energy expenditure to establish MET-levels. This is important since children have higher resting metabolic rates than adults. A recent study comparing the different cut-points for sedentary behavior (100, 300, 800, 1100) with the observation of children's behavior showed that Actigraph

cut-point of  $\leq 100$  cpm is the most appropriate one for quantifying time children spent on sedentary activities (Fischer et al. unpublished data). Another study by Trost et al. (40) evaluated the classification accuracy of cut-points (including those showed in Table 2) using energy expenditure measured by indirect calorimetry. They reported that the cut-points from Freedson et al. (36) and Treuth et al. (38) showed good classification accuracy. They also reported that 100 cpm as a cut-point for sedentary behavior showed good to excellent classification accuracy. Therefore, we selected the cut-points from Treuth et al. (38) for sedentary and activity classifications.

**Table 2.** Actigraph accelerometer cut-points that are appropriate for children aged 10-12.

Study	Sample	Cut-points (counts/minute)			
		Sedentary	Light	Moderate	Vigorous
Freedson et al.* (2005)	6-18 y, 80 children	-	100-2219	2220-4135	$\geq 4136$
Puyau et al. (2002)	6-16 y, 26 children	$\leq 800$	801-3199	3200-8199	$\geq 8200$
Treuth et al. (2004)	13-14 y, 74 girls	$\leq 100$	101-2999	3000-5199	$\geq 5200$
Mattocks et al. (2007)	12 y, 163 children	-	-	3581-6129	$\geq 6130$

\* Age specific equation, the cut-points for 12 years of age.

We also set a cut-point for the upper limit of count values to avoid spurious data based on the recommendations from Esliger et al. (16). It was shown that count values higher than 15.000 per minute are very unusual and implausible. For this reason the count values higher than 15.000 per minute were considered as missing values.

### Bouts of activity

Bouts of activity are sustained periods of elevated counts. Bouts of sedentary time are sustained periods of low counts. MeterPlus provides information on bouts of a specific intensity such as the frequency, when the bouts occurred, and how long they lasted. Analysis of bouts provides information on possible health benefits of the accumulation of certain intensity bouts (16). We selected to analyse sedentary behavior bouts lasting at least ten continuous minutes with two minutes tolerance.

### Outcome measures

We calculated total counts per day (the volume of activity), mean counts per minute (total counts divided by wearing time), and the amount of time spent in sedentary, and in light, moderate and vigorous intensity PA based on the vertical axis counts. We calculated the vector magnitude, a composite movement score for all three directions ( $[x^2 + y^2 + z^2]^{0.5}$ ).

### **Time filters**

We did not create time filters for sleeping time because this (as there are  $\geq 20$  minutes of consecutive zeros) is automatically coded as "non-wearing" time in Meterplus. We did not impute missing time because the review of Esliger et al. (16) concluded that in many cases duration of the non-wearing time obtained by diaries is ambiguous. In addition, imputation conflicts with the objectivity of accelerometer data.

#### *School-time time filters*

Schoolchildren spend 57% of their waking time at school and school time PA opportunities accounts for >70% of moderate and vigorous PA/day for children (41). Therefore, we plan to analyse school-time activity and sedentary behavior patterns apart from the total day. For this reason, we created time filters for school-times during data cleaning in Meterplus. We collected the exact school-times from the school management forms filled out by the manager of each school. We used the same data processing criterion of a recent study from Van Sluis et al. (42) and children who did not reach 500 minutes collected on at least 2 weekdays were excluded from the school-time analysis. We obtained the total valid hours per school day and the total time spent in sedentary behavior, light, moderate and vigorous activity. We calculated the mean percentages of time spent in sedentary, moderate and vigorous activity categories per school day.

### **Discussion**

The cross-European survey in the ENERGY-project aimed to identify the potential determinants of EBRBs and the differences between countries regarding these determinants among schoolchildren and their parents. The accelerometer sub-study provides an objective measurement of PA and sedentary behavior of children. With the systematic development of the accelerometer study protocol, we aimed to use the best evidence for the decisions on data collection and data processing with accelerometers. We also standardised the use of accelerometers between the participating countries including data collection and data analysis.

To examine health impacts of PA and sedentary behavior, researchers transfer accelerometry count values to behavioral outcome such as time spent in a specific activity category. It is important to select appropriate data processing criteria. For example, applying different cut-points to the same data significantly changes the estimation of activity and sedentary time (43,44). This will affect the relationship of activity and sedentary time with a specific health outcome. To choose an evidence based cut-point, we summarised and evaluated existing paediatric calibration studies. When selecting cut-points for the activity categories the age group of study sample needs to be taken into consideration as the same cut-point may not work well across different age groups (10). In the current study we have only one age group, i.e., 10-12 years old.

Furthermore, a recent study that used NHANES 2005-2006 accelerometry data showed that applying stricter compliance criteria during data processing resulted in somewhat higher moderate to vigorous PA levels (29). In the current study we had to exclude nearly 30% of children from further analyses due to insufficient valid days. More studies on

differences between different compliance criteria are needed, for example, the effect of lowering the minimum of three weekdays to two weekdays.

Another critical issue with accelerometer use is compliance to the protocol. Sirard and Slater (45) compared different methods among high school students to increase compliance such as giving incentives, diary and telephone calls. They showed that giving a diary to note down non-wearing time and reasons is an effective, low cost method to increase compliance. We used the diary method resulting in 67% compliance, which is worse to what has been found before by Sirard and Slater (45). It indicates the need for additional compliance strategies for younger children.

One of the strengths of the accelerometer study is a large sample size from different countries in Europe that provides an overview of activity patterns across countries. Researchers should be aware of weaknesses of accelerometry use as well. Some of the limitations of accelerometers are not measuring the movement of arms in terms of carrying, lifting weights, not distinguishing between sitting, lying, and standing still and poor measurement of cycling behavior. Furthermore, water-based activities cannot be measured by accelerometry. For the selection of accelerometer type several issues should be considered such as costs, software support, comparability with other studies and device features (46). In the current study we selected the Actigraph models due to their common use and good psychometric properties against other accelerometer types among children (18).

To our knowledge, there is no validation study of vector magnitude value for children. Since it contains information of body movements of all three planes, it may increase the accuracy of the behavioral outcomes. There is a need for future studies on validation of vector magnitude for children. Even though there is best practice evidence on accelerometry use, there is still no consensus on several issues. For example, the number of days that are needed for school children to reflect their habitual activity patterns is not clear. Another critical issue that needs further study is missing data imputation using diary data. We decided not to impute data due to the ambiguity of diary data and also unknown mechanisms of missingness of the accelerometer data. Catellier et al. (47) indicated that imputation of missing data would bias the result when missingness is not random, such as people less likely to wear the device when they are more inactive. Future studies might explore the usefulness of data imputation and comparison of different techniques of imputation with accelerometer data such as a recent study from Ottevaere et al. (48).

## **Conclusion**

This evidence-based protocol of accelerometry use in children may guide future studies in the field of PA and sedentary behavior measurement. The decisions on data collection and processing should be sample-specific and based on critical evaluation of available evidence. Despite its shortcomings, the accelerometry measurement is an advantageous method in PA and sedentary behavior measurement among children. Further study on the unresolved methodological issues such as validation of vector magnitude and imputation techniques is recommended.

## References

1. Kohl HW, Fulton JE, Caspersen CJ. Assessment of physical activity among children and adolescents: a review and synthesis. *Prev Med* 2000; 31:54-76.
2. Lamonte MJ, Ainsworth BE. Quantifying energy expenditure and physical activity in the context of dose response. *Med Sci Sports Exerc* 2001 ;33(6):370-378.
3. Plasqui G, Westerterp KR. Physical activity assessment with accelerometers: an evaluation against doubly labeled water. *Obesity* 2007 ;15(10):2371-2379.
4. Mattocks C, Tilling K, Ness A, Riddoch C. Improvements in the measurement of physical activity in childhood obesity research; lessons from large studies of accelerometers. *Clin Med: Pediatrics* 2008 ;2:27-36.
5. Bringolf-Isler B, Grize L, Mader U, Ruch N, Sennhauser FH, Braun-Fahrlander C. Assessment of intensity, prevalence and duration of everyday activities in Swiss school children: a cross-sectional analysis of accelerometer and diary data. *Int J Behav Nutr Phys Act* 2009 ;6:50.
6. Chinapaw MJM, Mokkink LB, van Poppel M, van Mechelen W, Terwee CB. Physical activity questionnaires for youth. A systematic review of measurement properties. *Sports Med* 2010; 40(7):539-563.
7. Biddle SJH. Sedentary behavior. *Am J Prev Med* 2007; 33(6):502-503.
8. Tremblay MS, Colley RC, Saunders TJ, Healy GN, Owen N. Physiological and health implications of a sedentary lifestyle. *Appl Physiol Nutr Metab* 2010 ;35(6):725-740.
9. Chinapaw MJM, Proper KI, Brug J, van Mechelen W, Singh AS. The relationship between young peoples' sedentary behavior and biomedical health indicators: A systematic review of prospective studies. *Obes Rev* 2011, 12(7):e621-632.
10. Pate RR, O'Neill JR, Lobelo F. The evolving definition of "sedentary.". *Exerc Sport Sci Rev* 2008; 36:173-178.
11. Biddle SJH, Pearson N, Ross GM, Braithwaite R. Tracking of sedentary behaviors of young people: a systematic review. *Prev Med* 2010; 51:345-351.
12. Mitchell JA, Mattocks C, Ness AR, Leary SD, Pate RR, Dowda M, et al. Sedentary behavior and obesity in a large cohort of children. *Obesity* 2009; 17(8):1596-1602.
13. Marshall SJ, Gorely T, Biddle SJH. A descriptive epidemiology of screen based media use in youth: a review and critique. *J Adolesc* 2006 ;29:333-349.
14. Murphy SL. Review of physical activity measurement using accelerometers in older adults: considerations for research design and conduct. *Prev Med* 2009; 48:108-114.
15. Rowlands AV. Accelerometer assessment of physical activity in children: An update. *Pediatric Exer Sci* 2007; 19:252-266.
16. Esliger DW, Copeland JL, Barnes JD, Tremblay MS. Standardizing and optimizing the use of accelerometer data for free-living physical activity monitoring. *J Phys Act Health* 2005; 3:366-383.
17. Reilly JJ, Penpraze V, Hislop J, Davies G, Grant S, Paton JY. Objective measurement of physical activity and sedentary behavior: review with new data. *Arch Dis Child* 2008; 93:614-619.
18. De Vries SI, Bakker I, Hopman-Rock M, Hirasiga RA, van Mechelen W. Clinimetric review of motion sensors in children and adolescents. *J Clin Epid* 2006; 59(7):670-680.
19. Ward D, Evenson KR, Vaughn A, Rodgers AB, Troiano RP. Accelerometer use in physical activity: best practices and research recommendations. *Med Science Sports Exerc* 2005; 37(11):582-588.
20. Brug J, te Velde SJ, Chinapaw MJM, Bere E, de Bourdeaudhuij I, Moore H, et al. Evidence-based development of school-based and family-involved prevention of overweight across Europe: The ENERGY project's design and conceptual framework. *BMC Public Health* 2010; 10:276.
21. Van Stralen MM, Te Velde SJ, Singh AS, De Bourdeaudhuij I, Martens MK, Van der Sluis M, et al. European Energy balance Research to prevent excessive weight Gain among Youth (ENERGY) project: Design and methodology of the ENERGY cross sectional survey. *BMC Public Health* 2011; 11:65.
22. The Actigraph website. [<http://www.theactigraph.com>].
23. John D, Sasaki JE, Freedson PS. Comparison of Activity Counts from the Actigraph GT3X and GT1M. 2010 [<http://www.theactigraph.com/article/research-database/validation/comparison-of-activity-counts-from-theactigraph-gt3x-and-gt1m>].
24. Trost SG, McIver KL, Pate RR. Conducting accelerometer-based activity assessments in field-based research. *Med Sci Sports Exerc* 2005; 37(11):531-543.

25. Bailey RC, Olson J, Pepper SL, Porszasz J, Barstow TJ, Cooper DM. The level and tempo of children's physical activities: an observational study. *Med Sci Sports Exerc* 1995; 27:1033-1041.
26. The Actilife-5 User Manual. Prepared by ActiGraph R&D and Software Departments. [<http://support.theactigraph.com/downloads/>].
27. Godfrey A, Conway R, Meagher D, ÓLaighin G. Direct measurement of human movement by accelerometry. *Med Engineering & Physics* 2008 ;30:1364-1386.
28. MeterPlus version 4.2 software from Santech, Inc. [<http://www.meterplussoftware.com>].
29. Tucker JM, Welk GJ. Accelerometer Data Processing in NHANES 2005-2006: Evaluation of Physical Activity Compliance Criteria. 2010 [<http://www.theactigraph.com/article/research-database/compliance/accelerometer-data-processing-in-nhanes-2005-2006-evaluation-of-physicalactivity-compliance-criteria/>].
30. Colley R, Gorber SC, Tremblay MS. Quality control and data reduction procedures for accelerometry-derived measures of physical activity. *Health Reports* 2010; 21(1), Statistics Canada, Catalogue no. 82-003-XPE.
31. Treuth MS, Sherwood NE, Butte NF, McClanahan B, Obarzanek E, Zhou A, et al. Validity and reliability of activity measures in African-American girls for GEMS. *Med Sci Sports Exerc* 2003; 35(3):532-539.
32. Stevens J, Murray DM, Baggett CD, Elder JP, Lohman TG, Lytle LA, et al. Objectively assessed associations between physical activity and body composition in middle school girls. The trial of activity for adolescent girls. *Am J Epid* 2007 ;166(11):1298-1305.
33. Troiano RP. Large-scale applications of accelerometers: New frontiers and new questions. *Med Sci Sports Exerc* 2007; 39(9):1501.
34. Stone MR, Rowlands AV, Eston R. Relationships between accelerometer assessed physical activity and health in children: impact of the activity intensity classification method. *J Sports Sci Med* 2009; 8:136-143.
35. Trost SG. State of the art reviews: measurement of physical activity in children and adolescents. *Am J Lifestyle Med* 2007; 1:299-314.
36. Freedson PS, Pober D, Janz KF. Calibration of accelerometer output for children. *Med Sci Sports Exerc* 2005; 37(11):523-530.
37. Puyau MR, Adolph AL, Vohra FA, Butte NF. Validation and calibration of physical activity monitors in children. *Obes Res* 2002 ;10:150-157.
38. Treuth MS, Schmitz K, Catellier DJ, McMurray RG, Murray DM, Almeida MJ, et al. Defining accelerometer thresholds for activity intensities in adolescent girls. *Med Sci Sports Exerc* 2004 ;36:1259-1266.
39. Mattocks C, Leary S, Ness A, Deere K, Saunders J, Tilling K, et al. Calibration of an accelerometer during free-living activities in children. *Int J Ped Obes* 2007; 2:218-226.
40. Trost SG, Loprinzi PD, Moore R, Pfeiffer KA. Comparison of accelerometer cut-points for predicting activity intensity in youth. *Med Sci Sports Exerc* 2010 ;43(7):1360-1368.
41. Guinhouya BC, Lemdani M, Vilhelm C, Hubert H, Apété GK, Durocher A. How school time physical activity is the "big one" for daily activity among schoolchildren: a semi-experimental approach. *J Phys Act Health* 2009; 6(4):510-519.
42. Van Sluijs EM, Jones NR, Jones AP, Sharp SJ, Harrison F, Griffin SJ. School level correlates of physical activity intensity in 10-year-old children. *Int J Pediatr Obes* 2010 ;6(2-2):e574-581.
43. Penpraze V, Reilly JJ, Montgomery C, Kelly LA, Paton JY, Grant S. Effect of accelerometer cut-points on apparent levels of physical activity and inactivity in young children. *Med Sci Sports Med* 2006; 38(5):565.
44. Guinhouya CB, Hubert H, Soubrier S, Vilhelm C, Lemdani M, Durocher A. Moderate-to-vigorous physical activity among children: discrepancies in accelerometry-based cut-off points. *Obesity* 2006; 14:774-777.
45. Sirard JR, Slater ME. Compliance with wearing physical activity accelerometers in high school students. *J Phys Act Health* 2009; 6(Suppl 1):148-155.
46. Cliff DP, Reilly JJ, Okely AD. Methodological considerations in using accelerometers to assess habitual physical activity in children aged 0-5 years. *J Scien Med in Sport* 2009; 12:557-567.
47. Catellier DJ, Hannan PJ, Murray DM, Addy CL, Conway TL, Yang S, et al. Imputation of missing data when measuring physical activity by accelerometry. *Med Sci Sports Exerc* 2005; 37(11):555-562.
48. Ottevaere C, Huybrechts I, De Meester F, De Bourdeaudhuij I, Cuenca-García M, De Henauw S. The use of accelerometry in adolescents and its implementation with non-wear time activity diaries in free-living conditions. *J Sports Sci* 2011; 29(1):103-13.

## SUPPLEMENTARY FILES

### Supplementary file 1. Diary

#### KIDS-METER LOG

At the next table, you see an example of how to fill in the diary for the kids meter. Write down the dates and days on which you wear the kids-meter. Note the times that you put it on and take it off during each day for getting up in the morning and going to bed for sleeping. If you put it off for more than 5 minutes please write this down below and for what reason (e.g., swimming, etc.)

#### Sample Entry

<b>Day</b>	Tuesday	
<b>Date</b>	March 1, 2010	
<b>School Day</b>	Yes (x) No ( ) Half-day ( )	
	Get up	Sleeping
<b>Time</b>	7.30	21.00
<b>Activity</b>	Start time	End time
<b>1.Swimming</b>	10.00	10.40
<b>2.</b>		

#### DON'T FORGET TO FILL THIS TABLE IN DURING THE NEXT 6 DAYS

Day												
Date												
School Day	Yes ( ) No ( ) Half-day ( )		Yes ( ) No ( ) Half-day ( )		Yes ( ) No ( ) Half-day ( )		Yes ( ) No ( ) Half-day ( )		Yes ( ) No ( ) Half-day ( )		Yes ( ) No ( ) Half-day ( )	
	Get up	Sleeping										
Time												
Activity	Start time	End time										
1.												
2.												
3.												

#### RESEARCHER WILL FILL IN THIS PART

Country Code:

School Code:

Class Code:

Date of Birth (Age):

Student Code:

Serial number:

Date of distributing:

Date of collecting back:

**Supplementary file 2. Instructions and information brochure.**



**KIDS-METER INFORMATION LETTER FOR PARENTS**

Dear Parents,

Thank you for participating in the ENERGY project. As part of the ENERGY project your child will wear a kids-meter during 6 days. The purpose of this letter is to inform you how your child should wear this kids-meter.

**What is a Kids-Meter?**

The Kids-Meter measures all the movements of your child. It is a mechanical device without radiation. We are interested in what kind of movements children like to do.

**How to wear the Kids-Meter?**

The Kids-Meter (attached to a belt) must be worn on the right waist; directly on the hip bone. Your child can wear it under or on his/her clothes. Please, make sure that the belt is tight enough (not too loose, and not too tight).



**How long and When to wear the Kids-Meter?**

Please remind your child to wear the Kids-meter for 6 consecutive days. Your child should put it on when she/he wakes up in the morning until she/he goes to bed for sleeping at night. It is not waterproof and should be removed during bathing, showering and swimming. Please remind your child to fill in the diary he/she received from us. In the diary he/she can write the exact time to put it off and on and the activities your child did during nonwear time.

Your child is supposed to wear the kids-meter from ..... till ..... Please remember her/him to bring the kids-meter back to school and give it to the teacher at .....

If you or your child has *any* questions or concerns regarding the study, please do not hesitate to contact us.

Sincerely,

Dr Mai Chin A Paw  
EMGO Institute, Vumc  
Van der Boechorststraat 7  
1081 BT Amsterdam  
Phone: +31 (0)20-4449988  
Email: [k.vanderweiden@vumc.nl](mailto:k.vanderweiden@vumc.nl)



## INSTRUCTIONS FOR KIDS-METER USE

- **WEAR** the monitor everyday ALL DAY for the next 6 days.
- **REMOVE** the monitor just before going to bed. Leave it on a table or dresser where you will be sure to see it and put it on first thing the next morning.
- **PUT** your monitor on each morning when you get out of bed.
- **WEAR** the kids-meter with the **STICKER ON TOP**.



- **BE SURE** the monitor is on the right side of your waist. **NOT** at the front or back of your waist.
  - **BE SURE** the monitor fits tightly around your waist. You can wear it under or above your clothes.
  - **DON'T** drop the monitor.
  - **DON'T** let the monitor get wet. Remove the device during showering, swimming and bathing or any sports in which it can get wet.
- **BE SURE** to put it back on when you are out of the water.
  - If you forget to put it on for any part of the day put it on as soon as you remember.
  - **Please NOTE** at what time you put the accelerometer on and off and why in your diary.
  - Please **DON'T** forget to bring it back to school after 6 days.



