

## PART II

---

# An exercise trial in patients with early-onset dementia



## CHAPTER 5

---

# Exercise and early-onset Alzheimer's disease; theoretical considerations

---

ASTRID M. HOOGHMSTRA<sup>1,2</sup>, LAURA H.P. EGGERMONT<sup>1</sup>,  
PHILIP SCHELTENS<sup>2</sup>, WIESJE M. VAN DER FLIER<sup>2,3</sup>, ERIK J.A. SCHERDER<sup>1</sup>

---

### **Affiliations**

<sup>1</sup> Department of Clinical Neuropsychology, VU University, Amsterdam, the Netherlands

<sup>2</sup> Alzheimer Center & Department of Neurology, VU University Medical Center, Neuroscience Campus Amsterdam, Amsterdam, the Netherlands

<sup>3</sup> Department of Epidemiology and Biostatistics, VU University Medical Center, Amsterdam, the Netherlands

*Published in:*

Dementia and Geriatric Cognitive Disorders Extra, 2012; 2:132-145.

## Abstract

---

**Background/Aims:** Although studies show a negative relationship between physical activity and the risk for cognitive impairment and late-onset Alzheimer's disease, studies concerning early-onset Alzheimer's disease (EOAD) are lacking. This review aims to justify the value of exercise interventions in EOAD by providing theoretical considerations that include neurobiological processes.

**Methods:** A literature search on key words related to early-onset dementia, exercise, imaging, neurobiological mechanisms and cognitive reserve was performed.

**Results/conclusion:** Brain regions and neurobiological processes contributing to the positive effects of exercise are affected in EOAD and ,thus, provide theoretical support for exercise interventions in EOAD. Finally, we present the design of a randomized controlled trial currently being conducted in early-onset dementia patients.

## Introduction

---

Epidemiological studies demonstrate a positive relationship between physical activity and cognitive functioning.<sup>1,2</sup> A decrease in the level of physical activity by a disturbance in gait and, consequently, in walking, coincides with a decline in cognitive functioning.<sup>1</sup> A decrease in the level of physical activity might even predict dementia.<sup>3</sup> In contrast, maintaining a physically active lifestyle may protect against dementia.<sup>4,5</sup>

A causal relationship, i.e. does physical activity improve cognitive functioning, can only be demonstrated by randomized controlled trials (RCTs). In one study, a daily physical activity intervention in healthy sedentary older persons improved executive functions, in particular working memory.<sup>6</sup> Intervention studies that examine the effects of physical activity on cognition in demented patients are limited and show mixed results. Positive effects on executive functions after aerobic exercise were found in small groups of elderly patients suffering from mild cognitive impairment.<sup>7</sup> Another study also observed positive effects of exercise on cognitive functioning in older persons at risk for Alzheimer's disease (AD).<sup>8</sup> In contrast, in older persons with moderate dementia, no positive effects on cognition were found, potentially explained by the fact that most of the patients were suffering from concomitant cardiovascular disease<sup>9</sup> (for a review, see<sup>10</sup>). In summary, positive effects of physical activity have been found in healthy older individuals, patients with mild cognitive impairment, and persons at risk for AD. In persons with moderate dementia, these positive effects were not found.

One of the brain regions that play a crucial role in executive functions is the prefrontal cortex.<sup>11</sup> The functions of the prefrontal cortex react positively to increased physical activity.<sup>12</sup> The functioning of other cortical areas, such as the parietal lobe, also show a positive relationship with physical activity.<sup>13</sup> The prefrontal and parietal lobe are particularly vulnerable in early-onset Alzheimer's disease (EOAD).<sup>14,15</sup> It is therefore remarkable that studies examining the effects of physical activity in this population are lacking.

Studies on the effects of physical activity interventions on cognition in this younger population may be worthwhile for a number of reasons. Firstly, few specific treatments are available for patients with EOAD. However, EOAD is increasingly recognised as a problem.<sup>16</sup> It places a large psychological and economic burden on patients and caregivers because of the patient's prominent role in society at time of disease onset.<sup>17</sup> A treatment, such as a physical activity program, might bring positive effects for both the patient and the caregiver. Secondly, EOAD patients suffer less from physical inconveniences than patients with late-onset Alzheimer's disease (LOAD)<sup>18</sup> and may therefore participate in a more intensive program; intensity of an exercise program is important for its effect on cognition.<sup>19</sup> Finally, positive effects of exercise on cognition in normal ageing and (very) early dementia can be explained by their beneficial effect on several neurobiological processes, such as neurogenesis, synaptogenesis, and angiogenesis.<sup>20</sup> Improvement of these processes may also benefit patients with EOAD.

The goal of the present review is to provide theoretical considerations that justify exercise interventions in EOAD. Within this scope, we will address the following topics. First, the physical functioning of early-onset Alzheimer's disease patients will be addressed. Subsequently, brain regions that respond positively to exercise and that are affected in EOAD will be discussed. Next, neurobiological mechanisms such as neurogenesis, synaptogenesis, angiogenesis, and neurotrophins that may underlie the effects of exercise on cognition in EOAD will be highlighted. Subsequently, the cognitive reserve hypothesis will be reviewed. Finally, we present the design of an RCT currently being conducted in early-onset dementia (EOD) patients in our center.

## Methods

---

Search databases were PubMed/MEDLINE and Web of Science. The search terms used were combinations of the keywords: early-onset and presenile in combination with: dementia, Alzheimer's disease, vascular dementia, and

frontotemporal dementia. During the literature search, it appeared that studies on EOD involve almost exclusively studies on EOAD. Therefore, this review targets exclusively this type of EOD.

With respect to studies on physical functioning the following search terms were used: physical functioning, gait (disturbance\*), balance, falling, and walking (speed). The search for imaging studies included the following search terms: image\*, damage\*, atrophy\* degenerate\*, single-photon emission computed tomography (SPECT), (functional) magnetic resonance imaging ((f)MRI), and positron emission tomography (PET), in combination with the terms brain, cortical, cerebral, lobe, cerebrum, encephalon, and grey matter. In relation to exercise, only studies concerning structural MRI were included. Studies on neurogenesis were searched through: neuro\*, brain, cell, dendri\*, in combination with, genesis, growth, branch\* survival, prolifera\*, plastic\*, death, damage, atrophy\*, and degenerat\*. The search for studies on synaptogenesis included the terms: synap\*, in combination with, genesis, formation, elimination, pruning, synaptophysin, and synaptotagmin. With respect to studies focusing on angiogenesis, the following search terms were used: angio\*, arterio, vessel\*, vascul\*, in combination with, genesis, growth, branch\*, prolifera\*, death, deterioration, elimination, and SPECT. Neurotrophin studies were gathered by the search terms: neurotrophin\*, neurotrophic factor, brain derived neurotrophic factor (BDNF), neuro-trophin-4 (NT-4), nerve growth factor (NGF), neuro-trophin-3 (NT-3), insuline-like growth factor (IGF), and new-neuro-trophin-1 (NNT-1). Studies regarding the cognitive reserve hypothesis included the search terms: cognitive reserve, brain reserve, and neural reserve. Studies were first selected based on the title. Subsequently, the residual studies were selected using the abstract and the content of the article. The final search was performed in June 2011.

## **Physical functioning of patients with EOAD in relation to LOAD**

---

We found one study on gait disturbances in EOAD compared to LOAD, showing that EOAD patients experience less gait disturbances (16% of the patients) in contrast to late-onset Alzheimer's disease patients (45%).<sup>18</sup> In LOAD, gait disturbances have recently been studied (for reviews, see <sup>21-23</sup>). Gait disturbances are caused by neuropathology in subcortical brain regions, e.g. basal ganglia, and in cortical areas, e.g. frontal lobe, and can hence be divided in parkinsonian and pseudoparkinsonian gait disturbances, respectively.<sup>24</sup> Gait disturbances can be experienced even in mild stages of AD (i.e. cautious gait),<sup>25,26</sup> and include decreased gait velocity, step length, static and dynamic balance, and a widened base.<sup>27</sup> In more advanced stages of AD 'frontal gait' can be observed: patients show a shuffling walking style and start and turn difficulties.<sup>25</sup>

Taken together, although studies are scarce, we assume that the physical condition of EOAD patients would permit participation in a more intensive exercise program than LOAD patients.

## **Brain regions that respond positively to exercise and are affected in EOAD**

---

*Exercise.* In Individuals with higher levels of cardiorespiratory fitness, the loss of grey matter, characteristic for aging, is reduced in prefrontal, superior and anterior parietal, medial temporal (specifically in the hippocampus), and occipital regions.<sup>12,28-31</sup> It is known that exercise has a beneficial influence on cardiorespiratory fitness.<sup>32</sup> Indeed, after aerobic exercise, increases in grey matter density were observed in prefrontal and temporal cortices; concerning the latter, these increases are particularly seen in the hippocampus.<sup>33,34</sup> For more details see Table 1 (p. 100).

*EOAD*. Neocortical atrophy is a neuropathological hallmark of *EOAD*.<sup>35-37</sup> MRI, SPECT and PET imaging techniques show that the above-mentioned brain regions that respond positively to exercise are affected in *EOAD*.<sup>14,15,35,36,38-42</sup> For more details on the aforementioned areas and on areas affected in *EOAD* but not related to exercise, see Table 2 (p. 101). Of note is that the primary sensory and motor areas are relatively preserved in *EOAD*,<sup>35</sup> implying that motor activity is still possible in these patients.

## **Neurobiological mechanisms, exercise, cognition, and *EOAD***

---

Animal and human experimental studies show that 3 major neurobiological mechanisms underlie the positive effects of exercise on brain structures and subsequently on cognitive function: neurogenesis, synaptogenesis, and angiogenesis.<sup>20,43,44</sup>

### **NEUROGENESIS**

*Exercise*. In a landmark paper, cell proliferation and neurogenesis is shown in the hippocampal area of mice having access to a running wheel.<sup>45</sup> Other studies report similar findings, suggesting that prolonged physical activity (voluntary wheel running) enhances neurogenesis in the hippocampus, both in adult mice and rats<sup>46-48</sup> as well as in aged mice and rats.<sup>49</sup> More specifically, wheel running in mice stimulates survival of newly generated neurons and their development into functional hippocampal neurons.<sup>50</sup> Furthermore, continued physical activity reduces the adult-dependent decrease in adult neurogenesis.<sup>51</sup> Increased neurogenesis after exercise is mainly coupled with positive effects on BDNF levels, as has been shown in an early study<sup>52</sup> and supported by several studies (for reviews, see<sup>20,53</sup>). Cerebral blood volume, coupling neuro- and angiogenesis, proves to be

elevated in the hippocampus in healthy subjects after 3 months of aerobic exercise.<sup>54</sup>

*EOAD.* Mutations in the presenilin 1 and 2 (PS1 and PS2) genes are linked to most autosomal dominantly inherited forms of EOAD. Several studies suggest an association between PS1<sup>55-58</sup> and PS2 mutations<sup>59,60</sup> and cell apoptosis, due to withdrawal of neurotrophins, and amyloid beta disposition.<sup>61,62</sup> PS1 mutations further impair enrichment-induced neurogenesis of hippocampal neural progenitor cells.<sup>63</sup> Although some studies reported increased hippocampal proliferation in EOAD,<sup>64,65</sup> this hippocampal proliferation does probably not reflect neurogenesis but rather glial proliferation and vascular changes.<sup>64</sup>

## SYNAPTOGENESIS

*Exercise.* Animal models show that aerobic training increases synaptic development and synaptic plasticity.<sup>66,67</sup> Voluntary exercise increases dendritic complexity in the dentate gyrus<sup>47,68</sup> and small heat shock proteins and pre- and post-synaptic proteins in the hippocampus in rats.<sup>69</sup> Moderate physical activity changes the level of synaptic proteins in the motor areas of the brain, and hence, may trigger brain plasticity in these areas.<sup>70</sup> Apart from exercise, synaptogenesis has mainly been studied in relation to environmental enrichment. Motor learning (rotarod training) increases synapse formation in the cerebellar cortex of female rats,<sup>71-73</sup> and acrobatic training enhances synaptogenesis in the motor cortex of male rats.<sup>74</sup> These synaptic changes rely on motor learning and not on repetitive use of synapses during physical activity only.<sup>71</sup> Cortical levels of synaptophysin are increased after stimulation (living in cages with toys, tunnels and a running wheel in comparison to regular cages) for 20 weeks.<sup>75</sup>

*EOAD.* Research on synapses is often performed using specific synaptic vesicle proteins, such as synaptophysin and synaptotagmin.<sup>76</sup> The level of synaptophysin is lower in EOAD than in LOAD, indicating a higher synapse

loss in EOAD.<sup>77</sup> In a preliminary study, synaptotagmin also seemed to be reduced in both cerebral spinal fluid and brain tissue in EOAD compared to age-matched healthy individuals.<sup>78,79</sup> Greater metabolic dysfunction in the hippocampi and the basal frontal cortex, reflecting greater synapse loss, is found in EOAD patients carrying the apolipoprotein ε4 (ApoE4) allele, compared to EOAD patients not carrying the ApoE4 allele and LOAD patients.<sup>14</sup>

## ANGIOGENESIS

*Exercise.* In rats, prolonged exercise (30 days of wheel running) induces angiogenesis and increased blood flow in the cerebellum, motor cortex and hippocampus.<sup>71,80-82</sup> Angiogenesis occurs in the motor cortex within 30 days from the onset of the exercise program and these effects seem to last over time.<sup>82</sup> Three weeks of exercise reduces neurologic deficits and infarct volume after an induced stroke in rats – a finding that is attributed to angiogenesis.<sup>83,84</sup> Elevated microvessel density is revealed in the striatum after exercise.<sup>83,84</sup> Growth factors that stimulate angiogenesis are already increased after 1–3 weeks of exercise in rats, and the levels of these factors are further elevated after 3 weeks of exercise.<sup>83</sup> Older adults who perform regular exercise show more constant levels of cerebral blood flow in comparison to an inactive control group.<sup>85</sup>

*EOAD.* Angiogenesis is hypothesized to be reduced in AD,<sup>86</sup> but it has not been specifically studied in EOAD.

## NEUROTROPHINS

Neurotrophins are proteins that support neural networks by stimulating the development of synapses, synaptic efficiency, and survival of neurons.<sup>87</sup> There are several different neurotrophins, e.g. NGF, BDNF, NT-3, NT-4, and IGF-1. Neurotrophins act in brain areas with a high degree of plasticity, such as the cerebral cortex and the hippocampus.<sup>88</sup>

*Exercise.* Several animal experimental studies reveal increased BDNF levels in rats after voluntary exercise<sup>48,53,88-92</sup> in Ammon's horn areas (CA1 and CA4) of the hippocampus, in layers II and III of the caudal cortex, and in retrosplenial cortices.<sup>93</sup> Also, NGF levels are increased in the dentate gyrus, in CA4 of the hippocampus, and in areas II and III of the caudal cortex after exercise.<sup>93</sup> Both BDNF and NGF levels increased in the motor cortex (layer V; neuron) and in the striatum (glia) after 3 weeks of wheel running.<sup>84</sup> Additionally, IGF-1 levels are increased after exercise;<sup>94,95</sup> which is thought to be neuroprotective.<sup>96</sup>

*EOAD.* Mutations in the PS1 and PS2 genes are thought to contribute to apoptotic cell death by means of trophic factor withdrawal.<sup>55,62,97</sup> PS1 mutations may alter cellular signalling systems associated with trophic factor-induced differentiation in PC12 cells.<sup>98</sup> This altered responsiveness to neurotrophic factors could play a role in the pathogenesis of neuritic degeneration and cell death in human PS1 mutation carriers.<sup>98</sup> It is also known that IGF-1 has anti-apoptotic effects. In EOAD, PS1 mutation-related apoptotic neuronal cell death may be caused by disruptions of IGF-1 signalling.<sup>99</sup>

## **Cognitive reserve**

---

In connection with neurogenesis and exercise, the 'cognitive reserve hypothesis' has been mentioned. Cognitive reserve is thought to operate as a buffer against cognitive decline in both healthy ageing<sup>100</sup> and neurodegenerative processes.<sup>101</sup> A broad set of determinants contribute to a greater cognitive brain reserve,<sup>102</sup> including exercise.<sup>6,103,104</sup> There is some inconsistency in the use of the term cognitive reserve. The literature shows a classification of cognitive reserve into more passive<sup>105</sup> and more active models.<sup>102</sup> Passive reserve is defined by quantitative neurobiological measures, such as brain volume and the number of neurons and synapses. If more neurons and synaptic connections are present, the brain is able to function longer

at a normal level after neuropathological damage resulting from a neurodegenerative process has been inflicted.<sup>106</sup> On the other hand, active reserve implies that the brain actively attempts to cope with neuropathology. Active reserve is determined by how efficient neural networks operate in a healthy brain (neural reserve) and by the ability to compensate via cognitive strategies and the deployment of different neural networks when the pre-existing networks are damaged (neural compensation).<sup>102</sup> To date, only a few studies have addressed the specific role of cognitive reserve in EOAD. Studies on the relation between *passive* reserve and EOAD conclude that passive reserve is *lower* in EOAD than in LOAD since EOAD patients show clinical symptoms at an earlier age and therefore have a lower premorbid count of neurobiological measures.<sup>14,77</sup> In contrast, studies investigating the association between *active* reserve and EOAD state that the degree of neuropathology at the moment of symptom onset is greater in EOAD than in LOAD. Active reserve is consequently presumed to be *larger* in younger than in older patients, since the younger patients are able to cope with a more severe state of neuropathology.<sup>107,108</sup>

## Conclusions

---

- The notion emerging from this review is that brain regions responding positively to exercise, such as frontal and parietal regions, are particularly affected in EOAD.<sup>109</sup> Damage in these areas results in a variety of EOAD-related clinical features, such as loss of planning skills, loss of initiative, and personality changes,<sup>110</sup> which are detrimental to an individual's autonomy.<sup>111</sup> From this result, it follows that exercise may help fight the symptoms associated with EOAD.
- Exercise leads to neurogenesis in the hippocampal formation, to synaptogenesis (particularly) in the cerebellum, to angiogenesis in the motor cortex, and to increased levels of neurotrophins. In EOAD, synapse loss and loss of neurons are neuropathological hallmarks. This suggests that exercise

may partly reverse the pathological mechanisms in EOAD.

- With respect to the cognitive reserve hypothesis, passive reserve is thought to be lower, and active reserve is considered to be higher in EOAD than in LOAD. Of note is that, to date, only a few studies have addressed this topic directly.
- There might be a difference between EOAD and LOAD in the way these disorders respond to a treatment. This difference may be due to a variation in disease progression. Patients with EOAD show a more rapid cognitive decline than patients with LOAD.<sup>40,112</sup> In a meta-analysis, AD patients with a more rapid disease progression showed greater cognitive benefit from rivastigmine treatment, than slowly progressing patients.<sup>113</sup> The question arises whether the same may account for non-pharmacological treatments, such as exercise interventions.

This review provides theoretical support for exercise interventions in EOAD. We now present the design of an RCT currently being conducted studying the effects of exercise on the course of dementia in EOD patients.

## **Design of an RCT in EOD: the EXERCISE-ON study**

---

The EXERcise and Cognition In Sedentary adults with Early-ONset dementia (EXERCISE-ON) study is a multicenter RCT in patients with EOD (AD, Vascular dementia, frontotemporal dementia, or other types of dementia). The aim of this study is to assess whether exercise slows down the progressive course of symptoms of EOD. Participants are randomly assigned to 1 of 2 exercise programs: the *aerobic exercise program* (using a bicycle ergometer) and the *flexibility and relaxation program* (flexibility and relaxation exercises). Both programs last 3 months, with a frequency of 3 times a week and are situated in a rehabilitation center. Measurements take place at baseline, after 3

months (end of the exercise program) and after 6 months. Primary outcomes are cognitive functioning (in particular executive functioning), (instrumental) activities of daily living, and quality of life. Secondary measures include physical, neuropsychological, and rest-activity rhythm measures. Outcome measures will be controlled for comorbid medical conditions (medical chart), depressive symptoms, and ApoE genotype in view of possible moderating effects on treatment outcome.

This study is the first to assess the effect of exercise on cognition in EOD patients.

# References

- 1.** Rosano C, Simonsick EM, Harris TB, et al. Association between physical and cognitive function in healthy elderly: The health, aging and body composition study. *Neuroepidemiology*. 2005;24(1-2):8-14.
- 2.** Verghese J, Wang CL, Lipton RB, Holtzer R, Xue XN. Quantitative gait dysfunction and risk of cognitive decline and dementia. *J Neurol Neurosurg Psychiatry*. 2007;78(9):929-935.
- 3.** Waite LM, Grayson DA, Piguot O, et al. Gait slowing as a predictor of incident dementia: 6-year longitudinal data from the Sydney Older Persons Study. *J Neurol Sci*. 2005;229-230:89-93.
- 4.** Fratiglioni L, Paillard-Borg S, Winblad B. An active and socially integrated lifestyle in late life might protect against dementia. *Lancet Neurol*. 2004;3(6):343-353.
- 5.** Rovio S, Kareholt I, Helkala EL, et al. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet Neurol*. 2005;4(11):705-711.
- 6.** Kramer AF, Hahn S, Cohen NJ, et al. Ageing, fitness and neurocognitive function. *Nature*. 1999;400(6743):418-419.
- 7.** Scherder EJA, Van Paasschen J, Deijen J-B, et al. Physical activity and executive functions in the elderly with mild cognitive impairment. *Aging Ment Health*. 2005;9(3):272-280.
- 8.** Lautenschlager NT, Cox KL, Flicker L, et al. Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *JAMA*. 2008;300(9):1027-1037.
- 9.** Eggermont LHP, Swaab DF, Hol EM, Scherder EJA. Walking the line: a randomised trial on the effects of a short term walking programme on cognition in dementia. *J Neurol Neurosurg Psychiatry*. 2009;80(7):802-804.
- 10.** Eggermont L, Swaab D, Luiten P, Scherder E. Exercise, cognition and Alzheimer's disease: More is not necessarily better. *Neurosci Biobehav Rev*. 2006;30(4):562-575.
- 11.** Stuss DT, Alexander MP. Executive functions and the frontal lobes: a conceptual view. *Psychol Res Forsch*. 2000;63(3-4):289-298.
- 12.** Colcombe SJ, Erickson KI, Raz N, et al. Aerobic fitness reduces brain tissue loss in aging humans. *J Gerontol A-Biol*. 2003;58(2):176-180.
- 13.** Rosano C, Venkatraman VK, Guralnik J, et al. Psychomotor Speed and Functional Brain MRI 2 Years After Completing a Physical Activity Treatment. *J Gerontol A-Biol*. 2010;65(6):639-647.
- 14.** Mosconi L, Herholz K, Prohovnik I, et al. Metabolic interaction between ApoE genotype and onset age in Alzheimer's disease: implications for brain reserve. *J Neurol Neurosurg Psychiatry*. 2005;76(1):15-23.
- 15.** Seo SW, Im K, Lee JM, et al. Effects of demographic factors on cortical thickness in Alzheimer's disease. *Neurobiol Aging*. 2011;32(2):200-209.

**16.**

Sampson EL, Warren JD, Rossor MN. Young onset dementia. *Postgrad Med J*. 2004;80(941):125-139.

**17.**

Shinagawa S, Ikeda M, Toyota Y, et al. Frequency and clinical characteristics of early-onset dementia in consecutive patients in a memory clinic. *Dement Geriatr Cogn Disord*. 2007;24(1):42-47.

**18.**

Seltzer B, Sherwin I. A comparison of clinical features in early-onset and late-onset primary degenerative dementia - one entity or 2. *Arch Neurol*. 1983;40(3):143-146.

**19.**

Rolland Y, Abellan van Kan G, Vellas B. Physical activity and Alzheimer's disease: from prevention to therapeutic perspectives. *J Am Med Dir Assoc*. 2008;9(6):390-405.

**20.**

Churchill JD, Galvez R, Colcombe S, Swain RA, Kramer AF, Greenough WT. Exercise, experience and the aging brain. *Neurobiol Aging*. 2002;23(5):941-955.

**21.**

Scherder E, Eggermont L, Sergeant J, Boersma F. Physical Activity and Cognition in Alzheimer's Disease: Relationship to Vascular Risk Factors, Executive Functions and Gait. *Rev Neurosci*. 2007;18:149-158.

**22.**

Scherder E, Eggermont L, Visscher C, Scheltens P, Swaab D. Understanding higher level gait disturbances in mild dementia in order to improve rehabilitation: "Last in-first out." *Neurosci Biobehav Rev*. 2011;35(3):699-714.

**23.**

Van Iersel MB, Hoefsloot W, Munneke M, Bloem BR, Rikkert MGMO. Systematic review of quantitative clinical gait analysis

in patients with dementia. *Z Gerontol Geriatr*. 2004;37(1):27-32.

**24.**

Kurlan RE, Richard IH, Papka M, Marshall F. Movement disorders in Alzheimer's disease: More rigidity of definitions is needed. *Mov Disord*. 2000;15(1):24-29.

**25.**

O'keefe ST, Kazeem H, Philpott RM, Playfer JR, Gosney M, Lye M. Gait disturbance in Alzheimer's disease: A clinical study. *Age Ageing*. 1996;25(4):313-316.

**26.**

Prehogan A, Cohen CI. Motor dysfunction in dementias - Differential diagnosis of dementia requires acute understanding of motor symptoms. *Geriatrics*. 2004;59(11):53-60.

**27.**

Petterson AF, Engardt M, Wahlund LO. Activity level and balance in subjects with mild Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2002;13(4):213-216.

**28.**

Erickson KI, Prakash RS, Voss MW, et al. Aerobic Fitness is Associated With Hippocampal Volume in Elderly Humans. *Hippocampus*. 2009;19(10):1030-1039.

**29.**

Erickson KI, Raji CA, Lopez OL, et al. Physical activity predicts gray matter volume in late adulthood The Cardiovascular Health Study. *Neurology*. 2010;75(16):1415-1422.

**30.**

Gordon BA, Rykhlevskaia EI, Brumback CR, et al. Neuroanatomical correlates of aging, cardiopulmonary fitness level, and education. *Psychophysiology*. 2008;45(5):825-838.

**31.**

Rovio S, Spulber G, Nieminen LJ, et al. The effect of midlife physical activity on struc-

tural brain changes in the elderly. *Neurobiol Aging*. 2010;31(11):1927–1936.

**32.**

Pollock M, Gaesser G, Butcher J, et al. American College of Sports Medicine Position Stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sport Exerc*. 1998;30(6):975–991.

**33.**

Colcombe SJ, Erickson KI, Scalf PE, et al. Aerobic exercise training increases brain volume in aging humans. *J Gerontol A-Biol*. 2006;61(11):1166–1170.

**34.**

Erickson KI, Voss MW, Prakash RS, et al. Exercise training increases size of hippocampus and improves memory. *P Natl Acad Sci USA*. 2011;108(7):3017–3022.

**35.**

Frisoni GB, Pievani M, Testa C, et al. The topography of grey matter involvement in early and late onset Alzheimer's disease. *Brain*. 2007;130:720–730.

**36.**

Ishii K, Kawachi T, Sasaki H, et al. Voxel-based morphometric comparison between early- and late-onset mild Alzheimer's disease and assessment of diagnostic performance of Z score images. *Am J Neuroradiol*. 2005;26(2):333–340.

**37.**

Mizuno Y, Ikeda K, Tsuchiya K, Ishihara R, Shibayama H. Two distinct subgroups of senile dementia of Alzheimer type: Quantitative study of neurofibrillary tangles. *Neuropathology*. 2003;23(4):282–289.

**38.**

Johnson KA, Lopera F, Jones K, et al. Presenilin-1-associated abnormalities in regional cerebral perfusion. *Neurology*. 2001;56(11):1545–1551.

**39.**

Kemp PM, Holmes C, Hoffmann SMA, et al. Alzheimer's disease: differences in technetium-99m HMPAO SPECT scan findings between early onset and late onset dementia. *J Neurol Neurosurg Psychiatry*. 2003;74(6):715–719.

**40.**

Rabinovici GD, Furst AJ, Alkalay A, et al. Increased metabolic vulnerability in early-onset Alzheimer's disease is not related to amyloid burden. *Brain*. 2010;133:512–528.

**41.**

Shiino A, Watanabe T, Maeda K, Kotani E, Akiguchi I, Matsuda M. Four subgroups of Alzheimer's disease based on patterns of atrophy using VBM and a unique pattern for early onset disease. *Neuroimage*. 2006;33(1):17–26.

**42.**

Shiino A, Watanabe T, Kitagawa T, et al. Different atrophic patterns in early- and late-onset Alzheimer's disease and evaluation of clinical utility of a method of regional z-score analysis using voxel-based morphometry. *Dement Geriatr Cogn Disord*. 2008;26(2):175–186.

**43.**

Kramer AF, Erickson KI, Colcombe SJ. Exercise, cognition, and the aging brain. *J Appl Physiol*. 2006;101(4):1237–1242.

**44.**

Mcauley E, Kramer AF, Colcombe SJ. Cardiovascular fitness and neurocognitive function in older Adults: a brief review. *Brain Behav Immun*. 2004;18(3):214–220.

**45.**

Van Praag H, Christie BR, Sejnowski TJ, Gage FH. Running enhances neurogenesis, learning, and long-term potentiation in mice. *P Natl Acad Sci USA*. 1999;96(23):13427–13431.

**46.**

Browne L, Sweeney BJ, Farrell MA. Late-onset neuroaxonal leucoencephalopathy with spheroids and vascular amyloid. *Eur Neurol.* 2003;50(2):85–90.

**47.**

Redila VA, Christie BR. Exercise-induced changes in dendritic structure and complexity in the adult hippocampal dentate gyrus. *Neuroscience.* 2006;137(4):1299–1307.

**48.**

Farmer J, Zhao X, van Praag H, Wodtke K, Gage FH, Christie BR. Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience.* 2004;124(1):71–79.

**49.**

Van Praag H, Shubert T, Zhao CM, Gage FH. Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci.* 2005;25(38):8680–8685.

**50.**

Bednarczyk MR, Aumont A, Decary S, Bergeron R, Fernandes KJL. Prolonged Voluntary Wheel-Running Stimulates Neural Precursors in the Hippocampus and Forebrain of Adult CD1 Mice. *Hippocampus.* 2009;19(10):913–927.

**51.**

Kronenberg G, Bick-Sander A, Bunk E, Wolf C, Ehninger D, Kempermann G. Physical exercise prevents age-related decline in precursor cell activity in the mouse dentate gyrus. *Neurobiol Aging.* 2006;27(10):1505–1513.

**52.**

Neeper SA, Gomezpinilla F, Choi J, Cotman C. Exercise and Brain Neurotrophins. *Nature.* 1995;373(6510):109.

**53.**

Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain

health and plasticity. *Trends Neurosci.* 2002;25(6):295–301.

**54.**

Pereira AC, Huddlestone DE, Brickman AM, et al. An in vivo correlate of exercise-induced neurogenesis in the adult dentate gyrus. *P Natl Acad Sci USA.* 2007;104(13):5638–5643.

**55.**

Guo Q, Sopher BL, Furukawa K, et al. Alzheimer's presenilin mutation sensitizes neural cells to apoptosis induced by trophic factor withdrawal and amyloid beta-peptide: Involvement of calcium and oxyradicals. *J Neurosci.* 1997;17(11):4212–4222.

**56.**

Guo Q, Furukawa K, Sopher BL, et al. Alzheimer's PS-1 mutation perturbs calcium homeostasis and sensitizes PC12 cells to death induced by amyloid beta-peptide. *Neuroreport.* 1996;8(1):379–383.

**57.**

Morgan GA, Guo Q, Chan SL, Gary DS, Osborne BA, Mattson MP. Defects of immune regulation in the presenilin-1 mutant knockin mouse. *Neuromolecular Med.* 2007;9(1):35–45.

**58.**

Gupta S, Singh R, Datta P, et al. The C-terminal tail of presenilin regulates Omi/HtrA2 protease activity. *J Biol Chem.* 2004;279(44):45844–45854.

**59.**

Da Costa CA, Paitel E, Mattson MP, et al. Wild-type and mutated presenilins 2 trigger p53-dependent apoptosis and down-regulate presenilin 1 expression in HEK293 human cells and in murine neurons. *P Natl Acad Sci USA.* 2002;99(6):4043–4048.

**60.**

Janicki S, Monteiro MJ. Increased apoptosis arising from increased expression of the Alzheimer's disease-associated

presenilin-2 mutation (N141I). *J Cell Biol.* 1997;139(2):485–495.

**61.**

Mattson MP, Robinson N, Guo Q. Estrogens stabilize mitochondrial function and protect neural cells against the pro-apoptotic action of mutant presenilin-1. *Neuroreport.* 1997;8(17):3817–3821.

**62.**

Mattson MP, Guo Q, Furukawa K, Pedersen WA. Presenilins, the endoplasmic reticulum, and neuronal apoptosis in Alzheimer's disease. *J Neurochem.* 1998;70(1):1–14.

**63.**

Choi SH, Veeraraghavalu K, Lazarov O, et al. Non-cell-autonomous effects of presenilin 1 variants on enrichment-mediated hippocampal progenitor cell proliferation and differentiation. *Neuron.* 2008;59(4):568–580.

**64.**

Boekhoorn K, Joels M, Lucassen PJ. Increased proliferation reflects glial and vascular-associated changes, but not neurogenesis in the presenile Alzheimer hippocampus. *Neurobiol Dis.* 2006;24(1):1–14.

**65.**

Ramirez-Rodriguez G, Itez-King G, Kempermann G. The new neuron formation in the adult hippocampus: Neurogenesis. *Salud Ment.* 2007;30(3):12–19.

**66.**

Lu B, Chow A. Neurotrophins and hippocampal synaptic transmission and plasticity. *J Neurosci Res.* 1999;58(1):76–87.

**67.**

Vaynman S, Ying Z, Gomez-Pinilla F. Interplay between brain-derived neurotrophic factor and signal transduction modulators in the regulation of the effects of exercise on synaptic-plasticity. *Neuroscience.* 2003;122(3):647–657.

**68.**

Eadie BD, Redila VA, Christie BR. Voluntary exercise alters the cytoarchitecture of the adult dentate gyrus by increasing cellular proliferation, dendritic complexity, and spine density. *J Comp Neurol.* 2005;486(1):39–47.

**69.**

Hu SX, Ying Z, Gomez-Pinilla F, Frautschy SA. Exercise can increase small heat shock proteins (sHSP) and pre- and post-synaptic proteins in the hippocampus. *Brain Res.* 2009;1249:191–201.

**70.**

Ferreira AFB, Real CC, Rodrigues AC, Alves AS, Britto LRG. Moderate exercise changes synaptic and cytoskeletal proteins in motor regions of the rat brain. *Brain Res.* 2010;1361:31–42.

**71.**

Black JE, Isaacs KR, Anderson BJ, Alcantara AA, Greenough WT. Learning Causes Synaptogenesis, Whereas Motor-Activity Causes Angiogenesis, in Cerebellar Cortex of Adult-Rats. *P Natl Acad Sci USA.* 1990;87(14):5568–5572.

**72.**

Kleim JA, Vij K, Ballard DH, Greenough WT. Learning-dependent synaptic modifications in the cerebellar cortex of the adult rat persist for at least four weeks. *J Neurosci.* 1997;17(2):717–721.

**73.**

Kleim JA, Swain RA, Armstrong KA, Napper RMA, Jones TA, Greenough WT. Selective synaptic plasticity within the cerebellar cortex following complex motor skill learning. *Neurobiol Learn Mem.* 1998;69(3):274–289.

**74.**

Jones TA, Chu CJ, Grande LA, Gregory AD. Motor skills training enhances lesion-induced structural plasticity in the motor cortex of adult rats. *J Neurosci.* 1999;19(22):10153–10163.

**75.**

Levi O, Jongen-Relo AL, Feldon J, Roses AD, Michaelson DM. ApoE4 impairs hippocampal plasticity isoform-specifically and blocks the environmental stimulation of synaptogenesis and memory. *Neurobiol Dis.* 2003;13(3):273-282.

**76.**

Thome J, Pesold B, Baader M, et al. Stress differentially regulates synaptophysin and synaptotagmin expression in hippocampus. *Biol Psychiatry.* 2001;50(10):809-812.

**77.**

Bigio EH, Hynan LS, Sontag E, Satumtira S, White CL. Synapse loss is greater in presenile than senile onset Alzheimer disease: implications for the cognitive reserve hypothesis. *Neuropathol Appl Neurobiol.* 2002;28(3):218-227.

**78.**

Davidsson P, Jahn R, Bergquist J, Ekman R, Blennow K. Synaptotagmin, a synaptic vesicle protein, is present in human cerebrospinal fluid - A new biochemical marker for synaptic pathology in Alzheimer disease? *Mol Chem Neurobiol.* 1996;27(2):195-210.

**79.**

Gottfries CG, Lehmann W, Regland B. Early diagnosis of cognitive impairment in the elderly with the focus on Alzheimer's disease. *J Neural Transm.* 1998;105(8-9):773-786.

**80.**

Isaacs KR, Anderson BJ, Alcantara AA, Black JE, Greenough WT. Exercise and the Brain - Angiogenesis in the Adult-Rat Cerebellum After Vigorous Physical-Activity and Motor Skill Learning. *J Cereb Blood Flow Metab.* 1992;12(1):110-119.

**81.**

Kleim JA, Cooper NR, Van den Berg PA. Exercise induces angiogenesis but does not alter movement representations within rat motor cortex. *Brain Res.* 2002;934(1):1-6.

**82.**

Swain RA, Harris AB, Wiener EC, et al. Prolonged exercise induces angiogenesis and increases cerebral blood volume in primary motor cortex of the rat. *Neuroscience.* 2003;117(4):1037-1046.

**83.**

Ding YH, Luan XD, Li J, et al. Exercise-induced overexpression of angiogenic factors and reduction of ischemia/reperfusion injury in stroke. *Curr Neurovasc Res.* 2004;1(5):411-420.

**84.**

Ding Y, Li J, Luan X, et al. Exercise pre-conditioning reduces brain damage in ischemic rats that may be associated with regional angiogenesis and cellular overexpression of neurotrophin. *Neuroscience.* 2004;124(3):583-591.

**85.**

Rogers RL, Meyer JS, Mortel KF. After Reaching Retirement Age Physical-Activity Sustains Cerebral Perfusion and Cognition. *J Am Geriatr Soc.* 1990;38(2):123-128.

**86.**

Vagnucci AH, Li WW. Alzheimer's disease and angiogenesis. *Lancet.* 2003;361(9357):605-608.

**87.**

Meisner F, Scheller C, Kneitz S, et al. Memantine upregulates BDNF and prevents dopamine deficits in SIV-infected macaques: A novel pharmacological action of memantine. *Neuropsychopharmacology.* 2008;33(9):2228-2236.

**88.**

Tapia-Arancibia L, Aliaga E, Silhol M, Arancibia S. New insights into brain BDNF function in normal aging and Alzheimer disease. *Brain Res Rev.* 2008;59(1):201-220.

**89.**

Gomez-Pinilla F, Ying Z, Roy RR, Molteni R, Edgerton VR. Voluntary exercise induces a BDNF-mediated mechanism that

promotes neuroplasticity. *J Neurophysiol*. 2002;88(5):2187-2195.

**90.**

Oliff HS, Berchtold NC, Isackson P, Cotman CW. Exercise-induced regulation of brain-derived neurotrophic factor (BDNF) transcripts in the rat hippocampus. *Mol Brain Res*. 1998;61(1-2):147-153.

**91.**

Seifert T, Brassard P, Wissenberg M, et al. Endurance training enhances BDNF release from the human brain. *Am J Physiol Regul Integr Comp Physiol*. 2010;298(2):R372-R377.

**92.**

Vaynman S, Ying Z, Gomez-Pinilla F. Exercise induces BDNF and synapsin I to specific hippocampal subfields. *J Neurosci Res*. 2004;76(3):356-362.

**93.**

Neeper SA, Gomezpinilla F, Choi J, Cotman CW. Physical activity increases mRNA for brain-derived neurotrophic factor and nerve growth factor in rat brain. *Brain Res*. 1996;726(1-2):49-56.

**94.**

LLorens-Martin M, Torres-Aleman I, Trejo JL. Mechanisms Mediating Brain Plasticity: IGF1 and Adult Hippocampal Neurogenesis. *Neuroscientist*. 2009;15(2):134-148.

**95.**

Trejo JL, Carro E, Torres-Aleman I. Circulating insulin-like growth factor I mediates exercise-induced increases in the number of new neurons in the adult hippocampus. *J Neurosci*. 2001;21(5):1628-1634.

**96.**

Carro E, Trejo JL, Busiguina S, Torres-Aleman I. Circulating insulin-like growth factor I mediates the protective effects of physical exercise against brain insults of different etiology and anatomy. *J Neurosci*. 2001;21(15):5678-5684.

**97.**

Mattson MP, Guo Q. Cell and molecular neurobiology of presenilins: A role for the endoplasmic reticulum in the pathogenesis of Alzheimer's disease? *J Neurosci Res*. 1997;50(4):505-513.

**98.**

Furukawa KS, Guo Q, Schellenberg GD, Mattson MP. Presenilin-1 mutation alters NGF-induced neurite outgrowth, calcium homeostasis, and transcription factor (AP-1) activation in PC12 cells. *J Neurosci Res*. 1998;52(5):618-624.

**99.**

Tanii H, Ankarcrona M, Flood F, et al. Alzheimer's disease presenilin-1 exon 9 deletion and L250S mutations sensitize SH-SY5Y neuroblastoma cells to hyperosmotic stress-induced apoptosis. *Neuroscience*. 2000;95(2):593-601.

**100.**

Corral M, Rodriguez M, Amenedo E, Sanchez JL, Diaz F. Cognitive reserve, age, and neuropsychological performance in healthy participants. *Dev Neuropsychol*. 2006;29:479-491.

**101.**

Valenzuela MJ, Sachdev P. Brain reserve and dementia: a systematic review. *Psychol Med*. 2006;36(4):441-454.

**102.**

Stern Y. Cognitive reserve. *Neuropsychologia*. 2009;47(10):2015-28.

**103.**

Lam LC, Tam CW, Lui VW, et al. Modality of physical exercise and cognitive function in Hong Kong older Chinese community. *Int J Geriatr Psychiatry*. 2009;24:48-53.

**104.**

Richards M, Hardy R, Wadsworth MEJ. Does active leisure protect cognition? Evidence from a national birth cohort. *Soc Sci Med*. 2003;56(4):785-792.

**105.**

Katzman R. Education and the prevalence of dementia and Alzheimer's disease. *Neurology*. 1993;43:13-20.

**106.**

Mori E, Hirono N, Yamashita H, et al. Premorbid brain size as a determinant of reserve capacity against intellectual decline in Alzheimer's disease. *Am J Psychiatr*. 1997;154(1):18-24.

**107.**

Kim EJ, Cho SS, Jeong Y, et al. Glucose metabolism in early onset versus late onset Alzheimer's disease: an SPM analysis of 120 patients. *Brain*. 2005;128:1790-1801.

**108.**

Marshall GA, Fairbanks LA, Tekin S, Vinters H V, Cummings JL. Early-onset Alzheimer's disease is associated with greater pathologic burden. *J Geriatr Psychiatry Neurol*. 2007;20(1):29-33.

**109.**

McMurtray A, Clark DG, Christine D, Mendez MF. Early-onset dementia: frequency and causes compared to late-onset dementia. *Dement Geriatr Cogn Disord*. 2006;21(2):59-64.

**110.**

Kelley BJ, Boeve BF, Josephs K a. Cognitive and noncognitive neurological features of young-onset dementia. *Dement Geriatr Cogn Disord*. 2009;27(6):564-571.

**111.**

Cahn-Weiner DA, Farias ST, Julian L, et al. Cognitive and neuroimaging predictors of instrumental activities of daily living. *J Int Neuropsychol Soc*. 2007;13(5):747-757.

**112.**

Van der Vlies AE, Koedam ELGE, Pijnenburg YAL, Twisk JWR, Scheltens P, van der Flier WM. Most rapid cognitive decline in APOE epsilon 4 negative Alzheimer's disease with early onset. *Psychol Med*. 2009;39(11):1907-1911.

**113.**

Farlow MR, Small GW, Quarg P, Krause A. Efficacy of rivastigmine in Alzheimer's disease patients with rapid disease progression: Results of a meta-analysis. *Dement Geriatr Cogn Disord*. 2005;20(2-3):192-197.

**114.**

Frisoni GB, Testa C, Sabattoli F, Beltramello A, Soininen H, Laakso MP. Structural correlates of early and late onset Alzheimer's disease: voxel based morphometric study. *J Neurol Neurosurg Psychiatry*. 2005;76(1):112-114.

**115.**

Karas G, Scheltens P, Rombouts S, et al. Precuneus atrophy in early-onset Alzheimer's disease: A morphometric structural MRI study. *Neuroradiology*. 2007;49(12):967-976.

**Table 1. Brain regions that respond positively to exercise – structural MRI**

Reference	Design	Population (n)	Age (years) <sup>a</sup>	Exercise measurement	Brain region
Colcombe et al., 2003 <sup>12</sup>	Relational	Healthy older adults (55)	66.5 (5.3)	VO <sup>2</sup> max estimate	PFC; superior parietal
Colcombe et al., 2006 <sup>33</sup>	RCT 6 month (aerobic training vs. stretching toning)	<ul style="list-style-type: none"> <li>Healthy older adults (59)</li> <li>Young controls MRI (20)</li> </ul>	<ul style="list-style-type: none"> <li>EG 65.5</li> <li>CG 66.9</li> </ul>	VO <sup>2</sup> max estimate	PFC; temporal cortex
Erickson et al., 2011 <sup>34</sup>	RCT 1 year (aerobic exercise vs. stretching)	Healthy older adults (120)	<ul style="list-style-type: none"> <li>EG 67.6 (5.81)</li> <li>CG 65.5 (5.44)</li> </ul>	VO <sup>2</sup> max	Hippocampus
Erickson et al., 2010 <sup>29</sup>	Relational	Healthy older adults (299)	78	Number of blocks walked in one week.	PFC; occipital; entorhinal; hippocampus
Erickson et al., 2009 <sup>28</sup>	Relational	Healthy older adults (165)	66.5 (5.6)	VO <sup>2</sup> peak	Hippocampus
Gordon et al., 2008 <sup>30</sup>	Relational	<ul style="list-style-type: none"> <li>Healthy older adults (40)</li> <li>Young controls (20)</li> </ul>	71.5 (4.7)	VO <sup>2</sup> max	Medial temporal; anterior parietal; inferior frontal
Rovio et al., 2010 <sup>31</sup>	Follow-up (20.9 (4.9) years) after survey	<ul style="list-style-type: none"> <li>Healthy adults (31),</li> <li>MCI (23),</li> <li>Dementia (21)</li> </ul>	<ul style="list-style-type: none"> <li>Active (n = 32): 73.0 (3.5)</li> <li>Sedentary (n = 43): 72.1 (4.4)</li> </ul>	At follow-up: indication of weekly participation in physical activity	Frontal

**Note:**

<sup>a</sup>Mean and standard deviation unless otherwise specified

Abbreviations: CG = control group; EG = exercise group; MCI = mild cognitive impairment; MRI = magnetic resonance imaging; PFC = prefrontal cortex; RCT = randomized controlled trial; VO<sup>2</sup> max = maximum oxygen volume; VO<sup>2</sup> peak = peak oxygen volume. www

**Table 2. Brain regions that are affected in early-onset Alzheimer's disease**

Reference	Design	Population (n)	Age (years) <sup>a</sup>	Imaging technique	Brain region
Frisoni et al., 2007 <sup>35</sup>	EOAD vs. LOAD	<ul style="list-style-type: none"> <li>• EOAD (15)</li> <li>• LOAD (15)</li> <li>• Younger healthy adults (15)</li> <li>• Older healthy adults (15)</li> </ul>	<ul style="list-style-type: none"> <li>• 62.5 (5.4)</li> <li>• 78.5 (6.2)</li> <li>• 62.5 (5.4)</li> <li>• 76.8 (3.4)</li> </ul>	MRI	Occipital; frontal
Frisoni et al., 2005 <sup>114</sup>	EOAD vs. younger healthy adults	<ul style="list-style-type: none"> <li>• EOAD (9)</li> <li>• LOAD (9)</li> <li>• Younger healthy adults (9)</li> <li>• Older healthy adults (17)</li> </ul>	<ul style="list-style-type: none"> <li>• 62 (7)</li> <li>• 78 (4)</li> <li>• 61 (4)</li> <li>• 74 (6)</li> </ul>	MRI	Temporoparietal junction
Ishii et al., 2005 <sup>36</sup>	<ul style="list-style-type: none"> <li>• EOAD vs. LOAD</li> <li>• EOAD vs. younger healthy adults</li> </ul>	<ul style="list-style-type: none"> <li>• EOAD 1<sup>st</sup> group (30) 2<sup>nd</sup> group (20)</li> <li>• LOAD (30) (20)</li> <li>• Younger healthy adults (30) (20)</li> <li>• Older healthy adults (30) (20)</li> </ul>	<ul style="list-style-type: none"> <li>• 60.2 (5.2) 60.8 (4.6)</li> <li>• 71.5 (2.6) 72.2 (3.2)</li> <li>• 59.6 (3.8) 59.1 (2.7)</li> <li>• 71.4 (3.5) 70.3 (4.2)</li> </ul>	MRI	<ul style="list-style-type: none"> <li>• EOAD vs. LOAD: Precuneus; parietal; middle temporal; fusiform gyrus.</li> <li>• EOAD vs. younger healthy adults: Medial temporal, inferior parietal; precuneus; perisylvian; basal forebrain; inferior frontal areas</li> </ul>
Johnson et al., 2001 <sup>38</sup>	<ul style="list-style-type: none"> <li>• AD vs. PS1- healthy</li> <li>• AD vs. PS1+ asymptomatic</li> </ul>	<ul style="list-style-type: none"> <li>• PS1- healthy (23)</li> <li>• PS1+ asymptomatic (18)</li> <li>• PS1+ diag. AD (16)</li> </ul>	<ul style="list-style-type: none"> <li>• 42.7 (7.9)</li> <li>• 38.1 (7.2)</li> <li>• 51.0 (6.4)</li> </ul>	SPECT	<ul style="list-style-type: none"> <li>• AD vs. PS1- healthy: Posterior parietal; superior frontal</li> <li>• AD vs. PS1+ asymptomatic: Temporoparietal</li> </ul>
Karas et al., 2007 <sup>115</sup>	<ul style="list-style-type: none"> <li>• EOAD vs. LOAD</li> <li>• Correlational (MRI - age)</li> </ul>	AD (51)	69 (8.5)	MRI	Precuneus
Kemp et al., 2003 <sup>39</sup>	EOAD vs. LOAD retrospective	<ul style="list-style-type: none"> <li>• EOAD (20)</li> <li>• LOAD (44)</li> </ul>	<ul style="list-style-type: none"> <li>• 57.8 (4.1)</li> <li>• 76.4 (4.5)</li> </ul>	SPECT	Posterior association areas

Table 2 (continued)

Reference	Design	Population (n)	Age (years) <sup>a</sup>	Imaging technique	Brain region
Kim et al., 2005 <sup>107</sup>	EOAD vs. LOAD	<ul style="list-style-type: none"> <li>• EOAD (74)</li> <li>• LOAD (46)</li> <li>• Younger healthy adults (20)</li> <li>• Older healthy adults (13)</li> </ul>	Onset <ul style="list-style-type: none"> <li>• 55.7 (5.4)</li> <li>• 69.6 (3.1)</li> </ul> Exam. <ul style="list-style-type: none"> <li>• 59.1 (5.7)</li> <li>• 72.8 (3.6)</li> <li>• 56.4 (4.9)</li> <li>• 71.5 (2.0)</li> </ul>	PET	Superior temporal; inferior parietal; middle occipital; precuneus
Mosconi et al., 2005 <sup>14</sup>	EOAD vs. LOAD	<ul style="list-style-type: none"> <li>• EOAD ApoE4- (15)</li> <li>• EOAD ApoE4+ (12)</li> <li>• LOAD ApoE4- (34)</li> <li>• LOAD ApoE4+ (31)</li> <li>• Healthy adults (35)</li> </ul>	<ul style="list-style-type: none"> <li>• 60 (8)</li> <li>• 65 (5)</li> <li>• 77 (4)</li> <li>• 77 (4)</li> <li>• 69.3 (5.6)</li> </ul>	PET	Orbitofrontal; inferior parietal; inferior temporal
Rabinovici et al., 2010 <sup>40</sup>	EOAD vs. LOAD	<ul style="list-style-type: none"> <li>• EOAD (21)</li> <li>• LOAD (18)</li> <li>• Healthy adults (30)</li> </ul>	Onset <ul style="list-style-type: none"> <li>• 55.2 (5.9)</li> <li>• 72.0 (4.7)</li> </ul> Exam. <ul style="list-style-type: none"> <li>• 60.2 (6.2)</li> <li>• 77.8 (4.9)</li> <li>• 73.7 (6.4)</li> </ul>	PET	Temporo-parietal; middle temporal; precuneus; posterior cingulate; occipital
Seo et al., 2011 <sup>15</sup>	Correlational (age at onset – MRI)	<ul style="list-style-type: none"> <li>• AD (193)</li> <li>• Healthy adults (142)</li> </ul>	<ul style="list-style-type: none"> <li>• 73.5 (7.3)</li> <li>• 66.0 (7.9)</li> </ul>	MRI	Parietal
Shiino et al., 2008 <sup>42</sup>	Correlational (age – MRI)	<ul style="list-style-type: none"> <li>• AD (50)</li> <li>• Healthy adults (83)</li> </ul>	<ul style="list-style-type: none"> <li>• 73.1 (8.7)</li> <li>• 70.6 (6.4)</li> </ul>	MRI	Temporal; posterior cingulate
Shiino et al., 2006 <sup>41</sup>	Comparison of 4 subgroups of atrophy	<ul style="list-style-type: none"> <li>• AD (40)</li> <li>• MCI (20)</li> <li>• Younger healthy adults (40)</li> <li>• Older healthy adults (88)</li> </ul>	<ul style="list-style-type: none"> <li>• 71.1 (9.7)</li> <li>• 67.7 (9.0)</li> <li>• 24.5 (2.1)</li> <li>• 68.7 (8.7)</li> </ul>	MRI	Posterior cingulate; posterior cortices

**Note:** a Mean and standard deviation unless otherwise specified.

Abbreviations: AD = Alzheimer's disease; ApoE4 +/- = Apolipoprotein allele 4 present/absent; EOAD = early-onset Alzheimer's disease; exam. = examination; LOAD = late onset Alzheimer's disease; MRI = magnetic resonance imaging; PET = positron emission tomography; PS1+/- = presenilin 1 mutation present/absent; SPECT = single-photon emission computed tomography.