

Table of contents

	List of abbreviations.	9
Chapter 1	General introduction. <i>Adapted from Perit Dial Int. 2009 Nov-Dec;29(6):605-17.</i>	11
Chapter 2	Long-term intervention with heparins in a rat model of peritoneal dialysis. <i>Perit Dial Int. 2009 Jan-Feb;29(1):26-35.</i>	33
Chapter 3	BMP-7 blocks mesenchymal conversion of mesothelial cells and prevents peritoneal damage induced by dialysis fluid exposure. <i>Nephrol Dial Transplant. 2010 Apr;25(4):1098-1108.</i>	51
Chapter 4	Celecoxib treatment reduces peritoneal fibrosis and angiogenesis and prevents ultrafiltration failure in experimental peritoneal dialysis. <i>Nephrol Dial Transplant. 2009 Dec;24(12):3669-76.</i>	75
Chapter 5	Sunitinib treatment prevents angiogenesis in a rat model for peritoneal dialysis. <i>Submitted.</i>	93
Chapter 6	Vitamin D receptor activation reduces peritoneal remodelling in experimental peritoneal dialysis. <i>Submitted.</i>	103
Chapter 7	Peritoneal dialysis fluid incompatibility and new vessel formation promote leukocyte-endothelium interactions in a chronic rat model for peritoneal dialysis. <i>Microcirculation, 2010 May;17(4):271-280.</i>	123
Chapter 8	Loss of mesothelial Neutrophil-Gelatinase-Associated-Lipocalin (NGAL) reflects epithelial to mesenchymal transition in experimental peritoneal dialysis. <i>Submitted.</i>	141
Chapter 9	Protein identification in effluents of patients treated with different peritoneal dialysis fluids.	161
Chapter 10	Summary and conclusions	179
Chapter 11	Nederlandse samenvatting	190
	Dankwoord	196
	Curriculum Vitae	198
	List of Publications	199