

CONTENTS

• <i>Chapter 1: General introduction</i>	9
• <i>Chapter 2: Relation between objective and subjective measures of bladder dysfunction in multiple sclerosis</i>	23
Part 1: Physicians' perspective	
• <i>Chapter 3: How similar are commonly combined criteria for EDSS progression in multiple sclerosis?</i>	33
• <i>Chapter 4: Clinical impact of 20 % worsening on Timed 25-foot Walk and 9-hole Peg Test in multiple sclerosis</i>	45
• <i>Chapter 5: Responsiveness and predictive value of EDSS and MSFC in primary progressive MS</i>	55
Part 2: Patients' perspective	
• <i>Chapter 6: Psychometric evaluation of the Multiple Sclerosis Impact Scale (MSIS-29) for proxy use</i>	75
• <i>Chapter 7: Proxy measurements in multiple sclerosis: agreement between patients and their partners on the impact of multiple sclerosis in daily life</i>	89
• <i>Chapter 8: Responsiveness of two self-report outcome measures: the Multiple Sclerosis Impact Scale (MSIS-29) and the Multiple Sclerosis Walking Scale (MSWS-12)</i>	109
Part 3: Physicians' vs. patients' perspectives	
• <i>Chapter 9: Assessing disease progression in multiple sclerosis: combining physicians' and patients' perspectives?</i>	127
• <i>Chapter 10: General discussion</i>	145

Part 4: Appendices

• <i>Clinical measures</i>	163
• <i>Summary</i>	197
• <i>Nederlandse samenvatting</i>	203
• <i>Dankwoord</i>	211
• <i>Curriculum vitae</i>	219
• <i>List of Publications</i>	223