

CHAPTER 1

General introduction

The aging population

Life expectancy has risen remarkably over the last century in most developed countries. An almost linear lengthening of life span with a steady pace of almost three months per year in women and a little over two months in men has been reported since 1840 (1). The trend started off by saving the young until the 1950's and continued by an increased life expectancy at age 65 and later on at age 80. Over the last decades, the increase in life expectancy at birth is, apart from the increase of the proportion of people reaching old age, mostly caused by the fact that old age is prolonged in developed countries (2). Among persons aged 65 years and older, the proportion of the oldest old is growing. In many populations, the absolute number of persons aged 65 years and over has thus increased rapidly. In the Netherlands for example, more than 2.5 million persons were aged over 65 years in 2010, of whom about 650,000 persons were aged over 80 years. These numbers have increased with approximately 400,000 and 150,000 over the last ten years (3).

Obesity in the aging population

A worldwide increase of the prevalence of overweight and obesity has taken place over the last decades (4, 5). Overweight and obesity are mostly assessed by the Body Mass Index (BMI), the body weight (kg) divided by body height (m) squared. A BMI of ≥ 25 -30 kg/m² is considered overweight and a BMI of ≥ 30 kg/m² is considered obese. The prevalence of overweight and obesity has increased in all age groups, including older persons. Currently, the prevalence of overweight and obesity is highest among persons aged around 60-70 years in most countries (6). In the Netherlands for example, the prevalence of overweight and obesity in persons aged over 65 years has risen from 54.9% to 56.7% over the last ten years based on self-reported data on weight and height. In persons aged 75 years and over, the percentages are somewhat lower, but the increase is steeper, from 48.3% to 51.6%. Relatively more overweight older persons are obese as compared to ten years earlier, so also within the overweight population the severity of overweight has increased (3). In participants of the Longitudinal Aging Study Amsterdam (LASA) aged 55-65 years, the prevalence of obesity based on measured weight and height in women rose from 20.5% to 27.5% between 1992

and 2002. In men, there was an even more striking increase from 9.5% to 18.4% (7). The high and still increasing prevalence of obesity in a vastly growing older population causes an enormous raise in the absolute numbers of overweight and obese older adults.

The prevalence of overweight in younger populations is rising as well. Recent generations develop overweight at younger ages and most people tend to gain weight over their life span (8-10). Children and adolescents who are obese, have a higher risk of becoming an obese adult (11, 12). As a result, the prevalence of overweight and obesity in old age is expected to keep rising the upcoming decades. Moreover, future generations will have experienced greater cumulative exposure to excess weight when they reach old age (10).

Definition of obesity using anthropometry

Obesity is a disease in which an excess of body fat causes an increased risk for adverse health effects (4). A positive energy balance, caused by dietary intake of energy exceeding the energy expenditure, results in storage of the excess of energy in the body, mostly as fat mass. For practical reasons, obesity is usually assessed by indirect anthropometric measures. By use of anthropometric indicators individuals at risk are identified, selection of individuals for/in need of interventions takes place, and the effect of changing conditions or interventions is evaluated.

The most commonly used anthropometric measures are body mass index (BMI) and waist circumference. Adolphe Quetelet was the first to describe a ratio of weight adjusted by height squared in 1832. The term BMI was introduced by Keys et al. (13) in 1972. They concluded that the ratio best indicated adiposity because of its high correlation to weight in combination with its low correlation with height and its sensitivity to differences in weight. The cut-off values of BMI are the same for men and women and were based on the shape of the association between BMI and mortality. The cut-off values for under weight normal weight, overweight and obesity, which are displayed in Table 1, are adopted in the World Health Organization (WHO) guidelines since 1995 (14). Waist circumference is a measure of abdominal obesity as is the waist-hip ratio, which denotes the circumference of the waist relative to the circumference of the hip. Cut-off values

for high risk waist circumference were proposed by Lean et al. (15) in order to correctly classify those with a high BMI and/or waist-hip ratio in adult populations. Two different action levels were proposed for men and women separately (Table 1). Action level I was formulated to be a warning to avoid further weight gain while persons in action level II should be urged to lose weight. The WHO included both the cut-off values of BMI and waist circumference in their current guidelines and states that they are applicable in adults aged ≤ 70 years (16).

Table 1. Categories of BMI and waist circumference according to the WHO guidelines.

Risk categories	BMI	
	Men and women	
Underweight	$\leq 18.5 \text{ kg/m}^2$	
Normal weight	$\geq 18.5 - < 25 \text{ kg/m}^2$	
Overweight	$\geq 25 - < 30 \text{ kg/m}^2$	
Obesity	$\geq 30 \text{ kg/m}^2$	


	Waist circumference	
	Men	Women
Healthy waist	$< 94 \text{ cm}$	$< 80 \text{ cm}$
Action level I	$\geq 94 - < 102 \text{ cm}$	$\geq 80 - < 88 \text{ cm}$
Action level II	$\geq 102 \text{ cm}$	$\geq 88 \text{ cm}$

Limitations of current definition of overweight in older adults

With aging, the body composition changes. A (relative) loss of lean mass occurs and the distribution of fat over the body changes. At a given BMI an older adult has a higher body fat percentage as compared to younger adults (17, 18). Fat mass is increasingly stored around the organs and inside organs (visceral and

ectopic fat) and inside muscles (ectopic fat). As body fat redistributes to a more central fat distribution, subcutaneous fat on the extremities redistributes to visceral and ectopic fat (Figure 1) (17, 19, 20), causing a higher waist circumference relative to BMI in older as compared to younger adults (21). Moreover, an age-related height loss takes place which causes the BMI to increase with aging when weight remains stable (22-24). Because of these processes, anthropometric measures should be interpreted with caution in older adults. The measures might relate differently to health in older adults as compared to younger adult populations and cut-off values of anthropometric measures for the identification of overweight might shift.

Figure 1. Age-related redistribution of fat.



Consequences of overweight and obesity in old age

Research into the health consequences of obesity has been mostly performed in (young) adult populations. In addition, it focused mainly on mortality, cardiovascular diseases and cardiovascular risk factors like diabetes and hypertension. Overweight and obesity and its consequences do not develop over night; it takes quite some years before they emerge in clinical practice. Older persons have a lifetime of influences behind them and it might make quite a difference whether they developed overweight recently or already during childhood or adolescence.

The strength of the association of overweight and obesity with mortality declines in older adults as compared to (younger) adults and most findings suggest that mortality risks start to rise at relatively higher levels of overweight (25-30). As compared to the association with BMI, the association between waist circumference and mortality seems somewhat stronger in old age. Several reasons for the declining strength of the association between obesity and mortality with age have been suggested (26, 31). First, a survival effect could cause persons susceptible for the adverse effects of obesity to die prior to old age. Furthermore, the relative importance of a healthy BMI and waist circumference decreases because older adults have a higher (competing) risk to die from other causes. Also, obesity-related mortality risk increases over the years after the onset of obesity and the limited life span left in old age in which obesity-related health consequences can develop might play a role. Finally, a relative increasing importance of body mass (both lean and fat) as a reserve in periods of illness and confounding influences of smoking and weight loss are mentioned.

A trend of decreasing obesity-related mortality has been reported, while an increase in the association between obesity and disability is seen (32). Persons with obesity live longer with disease and disability, which is likely to increase health care costs (33, 34) and decrease quality of life (35, 36). Besides mortality and cardiovascular abnormalities, research on the consequences of obesity in old age has focused on functional measures and health outcomes affecting quality of life. Obesity is associated with an acceleration of age-related declines in health and function. Because of age-related losses in muscle mass and muscle strength, older adults are particularly susceptible for the negative consequences of overweight and obesity on physical function (37-39).

Several metabolic abnormalities, like glucose intolerance and hypertension, are independently associated both with aging and with (abdominal) obesity (40, 41). Arthritis is a highly prevalent condition in old age, which is also strongly associated with overweight and obesity (42). Especially, but not exclusively, the association with knee osteoarthritis is strong, probably because of the mechanical strain on the weight-bearing joints (43). Furthermore, the prevalence of incontinence increases with age, predominantly in women. The independent association of obesity with this age-related condition has been reported repeatedly (44, 45). These obesity-related health problems have all been related to a decreased quality of life in older adults (46). Apart from the indirect detrimental effect of overweight and obesity on quality of life through the increased risk for chronic disease and functional limitations, a direct effect has been described. A strong negative association between BMI and quality of life has been described in older adults (47, 48), which was particularly strong in the physical domains. The association is partly mediated by chronic diseases and age-related losses in function, but remains existent after adjustment.

Objectives

The general aim of this thesis is to add to the limited knowledge on the assessment and consequences of obesity in old age. By carefully describing its association with important health outcomes in large samples of older adults we aim to gain the insight in the consequences of obesity in old age. Furthermore, we describe the execution of advanced analyses to establish criteria for high risk waist circumference in old age which are currently lacking.

Epidemiology

The results described in this thesis were obtained using data from large, prospective cohort studies on aging. By use of these observational data, we describe obesity measures and their health consequences. In this way, the natural course of the (development of) consequences of obesity can be described. Using population based data, the results of the study have a high potential to be generalizable to the general older population. In all chapters, data from the Longitudinal Aging Study Amsterdam (LASA) were used (49, 50, 51). LASA is an

ongoing multidisciplinary study in which a representative sample of Dutch older adults is included. In Chapter 6, data from three other prospective cohort studies on aging were used; the Invecchiare in Chianti (InCHIANTI) study, the Age Gene/Environment Susceptibility—Reykjavik Study (AGES-Reykjavik) and the Health, Aging and Body Composition Study (Health ABC). In Table 2, an outline of the data used in each chapter of this thesis can be found as well as some of the studies' characteristics.

Table 2. Data of cohort studies used in each chapter.

Chapter	Study	Years of assessment	Number of participants	Inclusion criteria	Outcomes
2	LASA	2001-2002 2005-2006	899 770	Participants aged 55-65 at baseline Complete data on overweight history and functional limitations	Functional limitations
3	LASA	1992-1993 1995-1996 1998-1999	2,000 1,478 1,271	Presence of data on pain, BMI and waist circumference	Pain
4	LASA	1995-1996	1,049	Age \geq 70 years Complete data on self-reported mobility limitations, BMI and waist circumference	Mobility limitations
5	LASA	1992-1993 1995-1996 1998-1999 2001-2002 2005-2006	1,219 1,062 1,034 868 813	Age \geq 70 years Complete data on waist circumference	Pain Mobility limitations Urinary incontinence Knee osteoarthritis Quality of life Cardiovascular disease Diabetes
6	Health ABC InCHIANTI AGES-Reykjavik LASA	2002-2003 1998-2000 2002-2006 1992-1993	1,310 673 5,117 1,204	Age \geq 70 years Complete data on waist circumference Waist circumference >70 cm (women) or >80 cm (men)	Pain Mobility limitations Urinary incontinence Knee osteoarthritis Cardiovascular disease Diabetes

Outline of this thesis

In Chapter 2, we describe the possible cumulative impact of obesity over the life course on mobility limitations in late middle-age. In Chapter 3, the associations of both BMI and waist circumference with (the development of) pain are described. In Chapter 4, we use several methodological approaches in order to find the optimal method to identify an appropriate cut-off value of waist circumference for older adults aged ≥ 70 years. This optimal method is applied in the study described in Chapter 5, in which we develop potential cut-off values for waist circumference using a broad range of important health outcomes related to obesity. These new cut-off values are cross-validated using data of four different cohort studies from four different countries in Chapter 6. In Chapter 7, the results of the individual chapters are put into context; we discuss the over-all findings of this thesis and elaborate on their implications.

References

1. Oeppen J, Vaupel JW. Demography. Broken limits to life expectancy. *Science*. 2002;296(5570):1029-1031.
2. Christensen K, Doblhammer G, Rau R, et al. Ageing populations: the challenges ahead. *Lancet*. 2009;374(9696):1196-1208.
3. CBS StatLine, Statistics Netherlands (The Hague), 2010. (<http://statline.cbs.nl>).
4. James W, Jackson-Leach R, Mhurchu CN, et al. Overweight and obesity (high body mass index) In: Ezzati M, Lopez A, Rodgers A, et al., eds. Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors Geneva: WHO, 2004.
5. Finucane MM, Stevens G, Cowan MJ, et al. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *The Lancet*. 2011;377(9765):557-567.
6. Flegal KM, Carroll MD, Ogden CL, et al. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA*. 2010;303(3):235-241.
7. Visser M, Pluijm SM, van der Horst MH, et al. [Lifestyle of Dutch people aged 55-64 years less healthy in 2002/'03 than in 1992/'93]. *Ned Tijdschr Geneeskd*. 2005;149(53):2973-2978.
8. Nooyens AC, Visscher TL, Verschuren WM, et al. Age, period and cohort effects on body weight and body mass index in adults: The Doetinchem Cohort Study. *Public Health Nutr*. 2009;12(6):862-870.
9. Barone BB, Clark JM, Wang NY, et al. Lifetime weight patterns in male physicians: the effects of cohort and selective survival. *Obesity (Silver Spring)*. 2006;14(5):902-908.
10. Lee JM, Pilli S, Gebremariam A, et al. Getting heavier, younger: trajectories of obesity over the life course. *Int J Obes (Lond)*. 2010;34(4):614-623.
11. Serdula MK, Ivery D, Coates RJ, et al. Do obese children become obese adults? A review of the literature. *Prev Med*. 1993;22(2):167-177.
12. The NS, Suchindran C, North KE, et al. Association of adolescent obesity with risk of severe obesity in adulthood. *JAMA*. 2010;304(18):2042-2047.
13. Keys A, Fidanza F, Karvonen MJ, et al. Indices of relative weight and obesity. *J Chronic Dis*. 1972;25(6):329-343.
14. World Health Organisation (WHO) Expert Committee. *Physical status: the use and interpretation of anthropometry*. Technical Report Series. Geneva: WHO, 1995.

15. Lean ME, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ*. 1995;311(6998):158-161.
16. Clinical Guidelines on the Identification, Evaluation, and treatment of overweight and obesity in adults--The evidence report. National Institutes of Health. *Obes Res*. 1998;6 Suppl 2:51S-209S.
17. Hughes VA, Roubenoff R, Wood M, et al. Anthropometric assessment of 10-y changes in body composition in the elderly. *Am J Clin Nutr*. 2004;80(2):475-482.
18. Gallagher D, Visser M, Sepulveda D, et al. How useful is body mass index for comparison of body fatness across age, sex, and ethnic groups? *Am J Epidemiol*. 1996;143(3):228-239.
19. Cartwright MJ, Tchkonja T, Kirkland JL. Aging in adipocytes: potential impact of inherent, depot-specific mechanisms. *Exp Gerontol*. 2007;42(6):463-471.
20. Carmelli D, McElroy MR, Rosenman RH. Longitudinal changes in fat distribution in the Western Collaborative Group Study: a 23-year follow-up. *Int J Obes*. 1991;15(1):67-74.
21. Stevens J, Katz EG, Huxley RR. Associations between gender, age and waist circumference. *Eur J Clin Nutr*. 2010;64(1):6-15.
22. Sorkin JD, Muller DC, Andres R. Longitudinal change in height of men and women: implications for interpretation of the Body Mass Index: The Baltimore Longitudinal Study of Aging. *Am J Epidemiol*. 1999;150(9):969-977.
23. Visser M, Deeg DJ. The effect of age-related height loss on the BMI classification of older men and women. *Int J Body Compos Res*. 2007;5(1):35-40.
24. van Leer EM, van Noord PA, Seidell JC. Components of adult height and height loss. Secular trend and effects of aging in women in the DOM project. *Ann Epidemiol*. 1992;2(5):611-615.
25. Stevens J, Cai J, Pamuk ER, et al. The effect of age on the association between body-mass index and mortality. *N Engl J Med*. 1998;338(1):1-7.
26. Janssen I, Mark AE. Elevated body mass index and mortality risk in the elderly. *Obes Rev*. 2007;8(1):41-59.
27. Flicker L, McCaul KA, Hankey GJ, et al. Body mass index and survival in men and women aged 70 to 75. *J Am Geriatr Soc*. 58(2):234-241.
28. Baik I, Ascherio A, Rimm EB, et al. Adiposity and mortality in men. *Am J Epidemiol*. 2000;152(3):264-271.
29. Dolan CM, Kraemer H, Browner W, et al. Associations between body composition, anthropometry, and mortality in women aged 65 years and older. *Am J Public Health*. 2007;97(5):913-918.

30. Kuk JL, Ardern CI. Influence of age on the association between various measures of obesity and all-cause mortality. *J Am Geriatr Soc.* 2009;57(11):2077-2084.
31. Zamboni M, Mazzali G, Zoico E, et al. Health consequences of obesity in the elderly: a review of four unresolved questions. *Int J Obes (Lond).* 2005;29(9):1011-1029.
32. Alley DE, Chang VW. The changing relationship of obesity and disability, 1988-2004. *JAMA.* 2007;298(17):2020-2027.
33. Zizza CA, Herring A, Stevens J, et al. Obesity affects nursing-care facility admission among whites but not blacks. *Obes Res.* 2002;10(8):816-823.
34. Yang Z, Hall AG. The financial burden of overweight and obesity among elderly Americans: the dynamics of weight, longevity, and health care cost. *Health Serv Res.* 2008;43(3):849-868.
35. Fine JT, Colditz GA, Coakley EH, et al. A prospective study of weight change and health-related quality of life in women. *JAMA.* 1999;282(22):2136-2142.
36. Orfila F, Ferrer M, Lamarca R, et al. Gender differences in health-related quality of life among the elderly: the role of objective functional capacity and chronic conditions. *Soc Sci Med.* 2006;63(9):2367-2380.
37. Visser M, Goodpaster BH, Kritchevsky SB, et al. Muscle mass, muscle strength, and muscle fat infiltration as predictors of incident mobility limitations in well-functioning older persons. *J Gerontol A Biol Sci Med Sci.* 2005;60(3):324-333.
38. Visser M, Langlois J, Guralnik JM, et al. High body fatness, but not low fat-free mass, predicts disability in older men and women: the Cardiovascular Health Study. *Am J Clin Nutr.* 1998;68(3):584-590.
39. Launer LJ, Harris T, Rumpel C, et al. Body mass index, weight change, and risk of mobility disability in middle-aged and older women. The epidemiologic follow-up study of NHANES I. *JAMA.* 1994;271(14):1093-1098.
40. Goodpaster BH, Krishnaswami S, Harris TB, et al. Obesity, regional body fat distribution, and the metabolic syndrome in older men and women. *Arch Intern Med.* 2005;165(7):777-783.
41. Cefalu WT, Wang ZQ, Werbel S, et al. Contribution of visceral fat mass to the insulin resistance of aging. *Metabolism.* 1995;44(7):954-959.
42. Blagojevic M, Jinks C, Jeffery A, et al. Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage.* 2010;18(1):24-33.

43. Lohmander LS, Gerhardsson de Verdier M, Rollof J, et al. Incidence of severe knee and hip osteoarthritis in relation to different measures of body mass: a population-based prospective cohort study. *Ann Rheum Dis.* 2009;68(4):490-496.
44. Townsend MK, Curhan GC, Resnick NM, et al. BMI, Waist circumference, and incident urinary incontinence in older women. *Obesity (Silver Spring).* 2008;16(4):881-886.
45. Krause MP, Albert SM, Elsangedy HM, et al. Urinary incontinence and waist circumference in older women. *Age Ageing.* 2010;39(1):69-73.
46. Hawkins K, Pernarelli J, Ozminkowski RJ, et al. The prevalence of urinary incontinence and its burden on the quality of life among older adults with medicare supplement insurance. *Qual Life Res.* 2011;20(5):723-732.
47. Yan LL, Daviglius ML, Liu K, et al. BMI and health-related quality of life in adults 65 years and older. *Obes Res.* 2004;12(1):69-76.
48. Heo M, Allison DB, Faith MS, et al. Obesity and quality of life: mediating effects of pain and comorbidities. *Obesity Res.* 2003;11(2):209-216.
49. Deeg DJ, van Tilburg T, Smit JH, et al. Attrition in the Longitudinal Aging Study Amsterdam. The effect of differential inclusion in side studies. *J Clin Epidemiol.* 2002;55(4):319-328.
50. Deeg DJH, Westendorp-de Serière M. Autonomy and well-being in the aging population; Report from the longitudinal Aging Study Amsterdam 1992-1993. Amsterdam: VU University Press, 1994.
51. Huisman M, Poppelaars J, van der Horst M, et al. Cohort profile: The Longitudinal Aging Study Amsterdam. *Int J Epidemiol.* 2011;;40(4):868-876.