

# chapter **FIVE**

## HYPEROSMOLARITY AND HYPOXIA INDUCE CHONDROGENESIS OF ADIPOSE DERIVED STEM CELLS IN A COLLAGEN TYPE 2 HYDROGEL

Wouter JFM. Jurgens, Zufu Lu, Behrouz Zandieh-Doulabi,  
Dirk J. Kuik, Marco JPF. Ritt, Marco N. Helder

## ABSTRACT

Apart from soluble growth factors, various other biophysicochemical cues are known to promote chondrogenesis. Under physiological conditions cartilage in the joint comprises a hyperosmotic and hypoxic environment. Therefore, in this study, we examined the inductive effects of hyperosmotic and/or hypoxic conditions on cultured adipose stem cells (ASCs) and compared them with conventional TGF $\beta$ 1-induction. After encapsulation in collagen type II hydrogels and specific induction, ASCs were assessed for viability, proliferation, morphology and chondrogenic differentiation potential. Viability was similar under all conditions with low proliferative activity. After 4 days, hypoxia and/or hyperosmolarity did not affect round cell morphology in collagen type II hydrogels, while cells were mainly stretched in the TGF $\beta$ 1-induced group. At 21 days, the TGF $\beta$ 1-treated group had aggregated into a cell nodule. Hyperosmolarity mimicked this aggregation to a lesser extent, whereas cells under hypoxia stretched out after 21 days, with a combined effect in the hypoxic/hyperosmotic group. Both individual and combined hyperosmotic and/or hypoxic conditions significantly upregulated SOX5, SOX9, COMP and Link-p gene expression compared with the non-induced group, and to similar levels as the TGF $\beta$ 1-induced group. GAG synthesis in both hydrogel and medium was increased under hypoxic conditions, whereas hyperosmolarity decreased GAG formation in the hydrogels, but increased GAG formation in the medium. We conclude that in a joint mimicking the three-dimensional (3D) micro-environment, a combination of hyperosmolarity and hypoxia is able to induce chondrogenesis to the same extent as TGF $\beta$ 1. This might lead to an interesting alternative when considering short-term triggering in a one-step surgical procedure for the treatment of cartilaginous defects.

## INTRODUCTION

Cartilage tissue is well known for its avascular and aneural nature. As a consequence, cartilage shows only limited self-regenerative potential. Conventional treatments aimed at restoring or repairing the damaged cartilage tissue result in suboptimal functional repair<sup>1,2</sup>. Recently, regenerative therapies combining cells, scaffolds and biological factors aimed at biological repair and restoration of function are emerging. Mesenchymal stem cells (MSC) may be ideal candidates for these regenerative cell therapies, due to their high proliferation rate, multidifferentiation capacity<sup>3</sup> and superior cartilage quality<sup>4,5</sup>. Besides bone marrow, MSCs have been isolated from other sources like muscle, periosteum, cord blood and adipose tissue<sup>6</sup>. Adipose tissue is an attractive source since it is abundant, easy accessible through liposuction or resection, and contains a large number of stem cells, referred to as adipose derived stem cells (ASC)<sup>7</sup>. These characteristics make them a suitable candidate to apply in the one-step procedure for the treatment of (osteo)chondral defects<sup>8,9</sup>.

In general, induction of these cells into the chondrogenic lineage is accomplished by the addition of biochemical cues like (combinations of) growth factors, in particular members of the tissue growth factor  $\beta$  superfamily (TGF $\beta$ 1 and BMPs), insulin growth factor, or fibroblast growth factor<sup>10</sup>. In addition, biophysical cues like hypoxia<sup>11,12</sup>, hydrostatic pressure<sup>13,14</sup>, and ultrasound<sup>15</sup> have been found to induce chondrogenesis in MSCs.

Next to the induction by biophysicochemical cues, the behaviour and differentiation of MSC can be influenced by the complex biophysicochemical environment in which chondrocytes of articular cartilage reside<sup>16</sup>. This environment is avascular and the residing chondrocytes exist at a decreased oxygen tension of (1 to 7%)<sup>17,18</sup>. In chondrocyte culture systems increased synthesis of extracellular matrix (ECM) by chondrocytes under hypoxia has been shown<sup>19</sup>, and this has been extended to stem cells from bone marrow<sup>20</sup> and liposuction-derived adipose tissue<sup>17</sup> undergoing chondrogenesis. Next to a hypoxic environment, chondrocytes experience a unique ionic/osmotic environment, which is attributable to the polyanionic proteoglycans. They carry a high density of fixed negative charges that attract cations into the interstitial fluid<sup>21</sup>. Extracellular osmolality, which arises principally from the free extracellular ions, is in the range of 350–485 mOsmol, depending on cartilage type and zone, with cation concentrations, and hence interstitial osmolarity, following local proteoglycan gradients<sup>22</sup>. In both chondrocytes and intervertebral disc cells improvement in chondrogenic gene- and phenotype was found in an hyperosmotic environment<sup>23,24</sup> and similar results were found in stem cells from bone marrow upon chondrogenic induction<sup>25</sup>. As stated above, this environment cannot only negatively influence resident cells leading to the degeneration of joints, but moreover, may be critical for ASCs upon implantation in the joints for regenerative therapies, e.g. in a one-step surgical procedure for cartilage regeneration<sup>26</sup>. So far, the behaviour of ASC to the microenvironmental conditions of the joint is quite unknown or only partially investigated<sup>11,12,17,27</sup>.

In this study, ASC were encapsulated in a collagen type 2 gel to induce chondrogenesis. The effect of two biophysicochemical microenvironmental conditions of the joint were

explored, i.e. reduced oxygen concentration and high extracellular osmolarity. The influence of these two biophysical cues on the chondrogenesis of ASCs were compared with conventional biochemical induction with TGF $\beta$ 1. The parameters examined were viability, cellular morphology (well-known to influence stem cell lineage commitment<sup>32-34</sup>, proliferation, and the chondrogenic differentiation capacity of cultured ASC encapsulated in a collagen type II hydrogel.

## MATERIAL AND METHODS

### Adipose tissue-derived stem cell (ASCs) isolation and culture

Retrieval of human specimens was approved by the ethical committee of the VU University medical center, and informed consent was obtained.

ASCs were isolated from subcutaneous adipose tissue from patients ( $n = 9$ ) undergoing elective surgical procedures, as described previously<sup>31</sup>. Briefly, harvested tissue was enzymatically dissociated with 0.01 M Collagenase A (Roche diagnostics), decontaminated from erythrocytes by Ficoll density centrifugation, and the remaining stromal cells were plated in 25 cm<sup>2</sup> tissue culture flasks. Plating and expansion medium consisted of Dulbecco's modified Eagle's medium (DMEM; Invitrogen) supplemented with 10% fetal bovine serum (FBS; Hyclone), 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin and 2 mM L-glutamine (all Invitrogen). Cultures were grown in a humidified incubator at 37°C in an atmosphere of 5% CO<sub>2</sub>. On reaching 80-90% confluency, cells were detached with 0.5 mM EDTA/0.05% trypsin (Invitrogen), for 5 min at 37°C, and replated. In this way, a homogeneous population of ASCs was obtained, which was subsequently used for differentiation experiments at passage 3–4.

### Embedment of ASC in collagen II hydrogels and in vitro chondrogenic induction.

Collagen type II hydrogels were prepared by mixing six parts of collagen II (chicken sternal cartilage; Sigma), one part of 10x DMEM, one part of reconstitution buffer (2.2 g NaHCO<sub>3</sub> in 100 ml of 0.05 N NaOH and 200 mM Hepes) and two parts of cell suspension in culture medium. The final concentration of collagen type II was 2.4 mg/ml and the cell density was  $1.5 \times 10^6$  cells/ml hydrogel. All the components were mixed on ice; 100  $\mu$ l mixtures were seeded in 6-well plates and collagen hydrogel lattices were formed by placing the plates in a humidified incubator at 37°C in an atmosphere of 5% CO<sub>2</sub> for 4 h. Thereafter constructs were divided in 5 different experimental groups, in which different media under various culture conditions was carefully overlaid:

1. *The non-induced group* in plain medium [DMEM, containing ITS+™ Premix (BD Biosciences), 1% FCS (Hyclone), 25 $\mu$ M ascorbate-2-phosphate (Sigma), 100 U/ml penicillin, 100 $\mu$ g/ml streptomycin and 2 mM L-glutamine], under normoxic and iso-osmotic culture condition.
2. *The positive control group* in chondrogenic, iso-osmotic medium [plain medium supplemented with 10 ng/ml transforming growth factor- $\beta_1$  (TGF $\beta_1$ , ITK-diagnostics)] under normoxic culture condition.
3. *The hyperosmotic group* in plain, hyperosmotic medium (500 mOsm) under normoxic culture conditions.

3. *The hyperosmotic group* in plain, hyperosmotic medium (500 mOsm) under normoxic culture conditions.
4. *The hypoxic group* in plain, iso-osmolar medium under hypoxic (1% oxygen) culture conditions.
5. *The combined group* in plain, hyperosmotic medium under hypoxic culture conditions.

Osmolarity was adjusted by adding 5.4 g/L NaCl to the plain DMEM medium (320 mOsm), and measured using an osmometer (KNAUER, Germany). The hypoxic and combined groups were exposed to hypoxic conditions in a 5% CO<sub>2</sub> / 1% oxygen custom-designed hypoxia workstation (T.C.P.S. Rotselaar, Belgium) at 37°C in a humidified atmosphere; values for hyperosmolarity and hypoxia were based on previous experiments and experiments performed by others<sup>9,24,25</sup>. Hypoxic and combined hypoxic and hyperosmotic media were preincubated for 2h to reach the designated oxygen level prior to application. Medium (2 ml per well) was refreshed every 3–4 days until day 21.

### Cell viability and proliferation

Cell viability ( $n = 4$ ) in all groups was tested using a Toxilight® kit (Lonza) after 4, 7, 11, 14, 17 and 21 days according to the manufacturer's protocol. This assay is based on the release of the enzyme Adenylate Kinase (AK) from damaged cells and allows the accurate and sensitive analysis of cytotoxicity and cytolysis. ADP is added as substrate for AK, in which presence it is converted to ATP releasing a signal for assay by bioluminescence. Luminescence was measured using a luminometer (Victor microplate reader). To assess cellular toxicity, 100  $\mu$ L spent medium was used per condition and time-point, and measured in duplicate. Blank medium and medium from lysed cells served as negative and positive controls, respectively. The results are depicted as luminometer light output (RLU). Proliferation ( $n = 4$ ) of ASC in all groups was determined using CyQUANT® (Molecular Probes) analysis, as described previously<sup>9</sup>.

### Cytoskeletal organisation

The effects of different culture conditions on the cytoskeletal organization of ASCs was assessed by labeling the actin cytoskeleton, as previously described ( $n = 5$ )<sup>32</sup>. ASCs were cultured under specific conditions, as described above for 4 and 21 days, and fixed using 4% formaldehyde. F-actin fibers within the ASC were stained with 2.5 Units Alexa fluor 488 Phalloidin (Invitrogen) in PBS in the dark for 1 hour, followed by direct acquisition with confocal laser scanning microscope (Leica TCS SP5) using a 20x Carl-Zeiss lens. In this way, 2D images of the 3D environment of the hydrogel were obtained.

### Quantitative Real-time-Polymerase chain reaction (qRT-PCR)

Total RNA was extracted from the tissue using Trizol (Invitrogen) according to the manufacturer's instructions ( $n = 9$ ). RNA isolation and Reverse transcription were performed as previously described<sup>30</sup>. Relative target gene expression of aggrecan (ACAN), collagen type II (COL2A), collagen type X (COL10 $\alpha$ 1), cartilage oligomeric matrix protein (COMP), link protein (LINK-p), and the early transcription factors SOX-5/6/9 were determined in a

LightCycler 480 (Roche Diagnostics). Target genes were normalized to 18s housekeeping gene-expression to obtain the relative gene expression. Primers (Invitrogen) used for real-time PCR were previously described<sup>9,30</sup>.

Using Light Cycler software (version 4), the crossing points were assessed and plotted versus the serial dilution of known concentrations of the standards derived from each gene using Fit Points method. PCR efficiency was calculated by Light Cycler software and the data were used only if the calculated PCR efficiency was between 1.85-2.0.

### Histochemistry

Histochemistry ( $n = 4$ ) was performed as described previously<sup>31</sup>. In brief, medium was removed from the construct, rinsed with PBS twice and fixed using 4% paraformaldehyde. Then collagen type II hydrogels from each culture condition were stained directly with Alcian blue (Sigma) at acidic pH (1.0) for visualization of proteoglycans after 4 and 21 days.

### Extracellular Matrix Biosynthesis

Extracellular sulfated glycosaminoglycans (GAG) deposition in the collagen type II hydrogels was quantified after harvesting at 14 and 21 days. Overlying media were collected after 7, 14 and 21 days ( $n = 4$ ). The tissue-containing hydrogels were digested in 3% papain solution buffer overnight at 60°C. GAG were measured using a Blyscan kit (Biocolor LTD, Northern Ireland). Results were depicted as the total glycosaminoglycan production per gel, corrected for the amount of cells (DNA) in the gels using CyQUANT® analysis according to the manufacturer's protocol. Furthermore GAG release in the medium was depicted per ml medium as a treatment over control ratio (non-induced value set at 1).


### Statistics

Kolmogorov-Smirnov tests were used to determine the normalcy of measurements and, if appropriate, their logarithmics. Repeated measures ANOVA was used to determine significant differences when increasing time-points in one donor within one variable were compared, followed by Bonferroni post-hoc analysis. If levels of gene expression were below the detection limit (0.05), values were set at  $10^{-2}$  (or log level at -2). All statistical tests used a significance level of  $\alpha=0.05$ .

## RESULTS

### Cultured ASC in collagen type II hydrogel show similar viability and proliferative activity in all groups

Toxicity was 5-fold higher during the first 4 days in all groups (Figure 1). Compared with the positive control (known amounts of lysed ASC) and values of the CyQUANT assay, this corresponded with  $\pm 75\ 000$  cells. Thereafter, toxicity dropped to low levels, and cell number and viability remained constant, indicating a low proliferative activity (Figure 1). No significant differences in proliferative activity could be detected between the various groups at any time point (data not shown).




**Figure 1.** Cellular toxicity under different biophysicochemical conditions of ASCs cultured in collagen type II gel. Cellular toxicity was initially high due to harsh biophysicochemical induction on day 4 in all groups, thereafter decreasing to low values. Results are depicted as mean  $\pm$  SEM (n=4); \*\*\*p<0.001

### Hypoxia and hyperosmolarity cause round morphology of ASCs in collagen type II hydrogel during early induction, whereas hypoxia stimulates stretching during later induction

Actin filaments were stained to visualize cell shape or cytoskeletal organisation of the cells (Figure 2). This figure represents 2D images from the 3D situation in the hydrogels. This implicates that not all cells are in the same plane of focus in the gel. The morphological descriptions below therefore may not be captured in the images provided, but may be deduced from full 3D analysis of the gels.

After 4 days, most of the ASC in all groups -except the TGFβ1-induced group- showed round cell morphology in a collagen type II hydrogel with non-stressed or granule-like actin



**Figure 2.** Morphology of ASCs in collagen type II gel after culture for 4 days (A–E) and after 21 days (F–J) under various biophysicochemical conditions. Green fluorescent staining represents actin filaments, blue staining represents cell nuclei. ASCs in collagen type II gel mainly showed round morphology after 4 days under various biophysicochemical conditions, with relaxed actin filaments, except for the TGFβ1-group (upper row). After 21 days of culture under various conditions, the non-induced cells remained round, whereas the TGFβ1-group contracted into a nodule (see insert in G). Cells cultured under hypoxic conditions clearly stretched out after prolonged culture, whereas hyperosmolarity induced small nodules residing in the gel on top of stretched cells with stressed actin filaments. Bars=20μm

filaments (Figure 2A-E). Only ASC in the collagen type II hydrogels that touched the bottom of the wells stretched out, showing stressed actin filaments. Combined hyperosmotic/hypoxic conditions seemed to reduce cell surface slightly (Figure 2E). In contrast, TGF $\beta$ 1-stimulated ASC were mostly elongated with stressed actin filaments (Figure 2B). After 21 days, round cell morphology with membrane ruffles was maintained in the non-induced ASCs (Figure 2F), whereas TGF $\beta$ 1 stimulation resulted in chondrogenic nodule formation with round cells and dispersed actin cytoskeleton (Figure 2G). Hyperosmolarity maintained round cell shape after 21 days, with low aggregation level resulting in only small round cell masses with relaxed actin filaments and slightly more elongated cells with stressed actin filaments at the hydrogel bottom (Figure 2H). Hypoxic conditions caused elongated cells with stressed actin filaments after 21 days (Figure 2I). A combined pattern could be observed in the hypoxic and hyperosmotic group (further referred to as combined group; Figure 2J).


### **Hypoxia and/or hyperosmolarity stimulate chondrogenic gene expression comparable to TGF $\beta$ 1-induction**

Chondrospecific gene expression analysis showed significant upregulation of the early transcription factor SOX-5 at both time points in the TGF $\beta$ 1-induced, hyperosmotic and combined groups (Figure 3A), whereas the hypoxic group showed SOX-5 downregulation after both 4 and 14 days when compared to the non-induced group. For SOX-6, only the combined group showed a significant difference (in this case downregulation) compared to the TGF $\beta$ 1-stimulated group after 14 days of induction (Figure 3B). SOX-9 was significantly increased (compared to non-induced control) in the hyperosmotic and combined group after 4 days and at both time points in the TGF $\beta$ 1-induced group. ACAN gene expression was increased in the hyperosmotic group after 4 days and at both 4 and 14 days in the combined group compared with the TGF $\beta$ 1-induced control (Figure 3D). COL2a expression could only be detected, but not quantified, in 5 f 9 donors in all groups (data not shown). The late marker COMP was down-regulated in the hypoxic group, whereas the combined group showed significantly up-regulation of COMP to similar levels as TGF $\beta$ 1-induced controls (Figure 3E). A trend was only visible between the non-induced group and both the TGF $\beta$ 1-induced and combined groups, on both time-points ( $p=0.08$  for both). LINK-p gene expression was upregulated at both time points in the TGF $\beta$ 1-induced and combined groups compared to the non-induced group (Figure 3F). A significant difference in the LINK-p gene expression was observed after 14 days between the positive control and the hyperosmotic, the hypoxic and the non-induced group, but not in the combined group. Lastly, the hypertrophic gene COL10 was clearly expressed significantly more in the TGF $\beta$ 1-induced versus all other groups at both time-points (Figure 3G).


### **Hypoxic condition stimulates extracellular matrix formation**

Production of proteoglycans was assessed qualitatively using Alcian Blue staining after 4 and 21 days (Figure 4A-J), and quantitatively by biochemical measurement of sulphated glycosaminoglycan (GAG) deposition using a Blyscan kit after 14 and 21 days (Figure 5).

After 4 days only the non-induced cells and the positive control started to produce proteoglycans as visualized by Alcian Blue staining, whereas in the hyperosmotic,



**Figure 3.** Gene expression profile of ASCs cultured in collagen type II gel under various biophysico-chemical conditions compared to the non-induced group (treatment/control). The addition of hypoxia and/or hyperosmolarity clearly induced chondrogenic differentiation to similar levels to the TGFβ1-induced group, as indicated by significant differences in various early and late chondrospecific genes. The hypertrophic marker collagen type X was upregulated only in the TGFβ1-induced group, but not in the other groups. The results are depicted on a logarithmic scale as mean ± SEM (n=9). Significant differences to the non-induced group at a specific time-point are marked with an asterisk only, whereas differences to the TGFβ1-induced group are marked by connected lines. Overall differences between groups are marked in the legend: \*p<0.05; \*\*p<0.01; \*\*\*p<0.001




**Figure 4.** Alcian blue staining of ASCs in collagen type II gel after 4 days (A–E) and 21 days (F–J) of induction under various biophysicochemical conditions. After 4 days the non-induced and TGFβ1-induced group already showed some Alcian blue staining, whereas not much staining could be seen in the hyperosmotic, hypoxic and combined groups. After 21 days abundant and disseminated staining could be observed in the hypoxic group, whereas more patchy staining was seen in the combined group and lighter staining in the hyperosmotic group. Very bright staining could be seen in the TGFβ1-induced group; however, due to contraction only a very small nodule was left. The non-induced group stained only very lightly and in a patchy way. Bars=1 cm

hypoxic and combined group less staining could be observed (Figure 4A–E). However, after 21 days abundant proteoglycan formation could be seen [hypoxic > combined group > hyperosmotic group (Figure 4 H–J)]. The TGFβ1-induced positive control also stained intensively; however, due to contraction/aggregation, only a very small hydrogel was remaining (Figure 4G), resulting in a localised (non-uniform) staining. This same aggregation was observed in the non-induced control, staining only faintly and in a patchy way (Figure 4F), and, to a lesser extent, a similar patchy staining could be observed in the combined group (Figure 4J).

Quantification of ECM formation was performed by measuring the sulphated GAG deposition in the hydrogels and excretion of GAG into the medium. Very little GAG was formed in the extracellular environment of all groups after 14 and 21 days, whereas abundant GAG was released in the medium overlying the hydrogels (Figure 5A vs. 5B). After 14 days a trend towards increased GAG deposition could be observed in the hypoxic group compared with the other groups, however this difference did not reach significance, and decreased after 21 days. In contrast, in the hyperosmotic group and the combined group clearly less GAG was formed in the hydrogels (Figure 5A).

The overall GAG release in the medium in the non-induced group increased from  $2.3 \pm 0.9 \mu\text{g/ml}$  at 7 days to  $3.9 \pm 1.1 \mu\text{g/ml}$  after 21 days (data not shown). At the 7-day time-point, GAG release was significantly increased in all groups compared with the non-induced group (Figure 5B). The significance reached  $p < 0.01$  for the TGFβ1-induced and hypoxic groups, whereas the hyperosmotic and the combined hyperosmotic and hypoxic group reached  $p < 0.001$ . After 14 days only a significant increase could be detected due to the addition of hyperosmolarity or hypoxia (both  $p < 0.05$ ) and after 21 days no further significant differences in GAG excretion were detected.



**Figure 5.** Glycosaminoglycan (GAG) deposition of ASCs cultured in collagen type II gel under various biophysicochemical conditions. (A) GAG deposition in the gels after 14 and 21 days of induction, corrected for the number of cells. No significant differences in GAG deposition could be observed, although a trend towards increased GAG deposition could be observed in the hypoxic group after 14 days. (B) GAG excreted in the medium after 7, 14 and 21 days of induction corrected to the non-induced group. Since little GAG is retained in the extracellular matrix, most of it is released into the medium. Over time, excretion of GAG into the medium is diminished as the formed GAGs are becoming sequestered in the extracellular matrix. The results are depicted as mean  $\pm$  SD ( $n=4$ ). Significant differences to the non-induced group at a specific time-point are marked with an asterisk only, whereas differences to the TGFβ1-induced group are marked by connected lines; \* $p<0.05$ ; \*\* $p<0.01$ ; \*\*\* $p<0.001$

## DISCUSSION

This study is the first to investigate the effect of an hyperosmotic and combined hypoxic and hyperosmotic microenvironment on the viability, proliferation, morphology and chondrogenic differentiation potential of cultured ASC when encapsulated in a collagen type II hydrogel. The effect of hyperosmolarity was so far only determined in chondrocytes, where it was found to have an inductive effect on chondrogenic gene expressions and proteoglycan biosynthesis<sup>24,33</sup>. Although we realize ourselves that this experimental study represents only a simplified *in vitro* model of the complex *in vivo* joint environment, we are convinced that by varying the biophysicochemical conditions under controlled *in vitro* conditions, the relative influence of each factor on the viability, proliferation and chondrogenic differentiation capacity of the ASC can be determined.

We found that a) viability and proliferation of cultured ASC did not differ significantly between different culture conditions; b) cultured ASC in the collagen type II hydrogel showed mostly round morphology under all conditions except on TGFβ1-induction, which induced strong aggregation of cells into a big cell nodule; c) the 3D environment of the collagen type II hydrogel induced chondrogenic differentiation; d) a combination of hyperosmolarity and hypoxia had similar effects on the induction of chondrogenesis compared with TGFβ1 induction at both the genetic and protein level, as analyzed with qRT-PCR, histology and biochemical assays.

We hypothesized that hyperosmolarity (500 mOsm) in combination with hypoxia could lead to massive cell death due to harsh changes in biophysicochemical environment<sup>25,34</sup>.

However, contrary to our expectations, all groups initially experienced high toxicity, whereas from 4 days on toxicity was diminished and cell number remained stable after increasing culturing period (Figure 1). This early toxicity might be rather due to harsh experimental conditions embedding the ASC in the collagen type II hydrogel (e.g. it took 4 hours for the collagen type II hydrogel to solidify, before medium could be added), than to specific biophysicochemical cues. Another explanation can be found in the ontogenic process of chondrogenesis. This process is divided into an early phase of recruited mesenchymal cells aggregation/condensation to form chondroblasts, underlining the importance of cell-cell contact<sup>35-37</sup>. However, in a later phase a shifting occurs from cell-cell to cell-matrix interaction due to the production of abundant *ECM*<sup>38,39</sup> and apoptosis of cells. The toxicity can thus also be (partly) attributed to this developmental process of cartilage formation.

It is known that chondrogenic differentiation requires a 3D environment. Several methods exist to induce chondrogenic differentiation, like the commonly used pellet method, alginate beads or the use of scaffold materials or hydrogels<sup>40,41</sup>. In this study a collagen type II hydrogel was used which, in concordance with earlier studies, on itself already induced chondrogenic differentiation<sup>42,43</sup>. As proposed earlier by our group and also found in this study, the micro-environment plays a pivotal role in this induction, by influencing cell shape and the aggregation of cells<sup>43,44</sup>. This mechanism could explain why other studies using *ECM* proteins present in articular cartilage found a similar induction<sup>34,45</sup> and underlines the importance of matrix elasticity on differentiation of cells<sup>28,29</sup>.

While hypoxia and hyperosmolarity resulted in low proliferative activity of ASC, both single and combined induction enhanced differentiation of the ASCs into the chondrogenic lineage, both on the genetic level, by early up-regulation of SOX-genes and COMP and Link-p gene expression, as well as on the protein level, by enhanced proteoglycan production. The effect of hypoxia on ASC was already demonstrated by others<sup>12,27</sup>; however, contradictory results on the influence of hypoxia on the chondrogenic differentiation of ASC were presented. This might be attributed to differences in the culture and/or induction methods used, including the percentage of serum, the use of dexamethasone, the type of growth factor used, glucose and glutamine concentrations and/or the oxygen tensions applied.

Remarkably, in this study, a high gene up-regulation of collagen type X could be observed in the TGFβ1-induced group, whilst the other groups did not show any upregulation. Although sometimes disputed, this gene might be a marker for hypertrophic chondrogenic differentiation<sup>46,47</sup> and concomitant (too) aggressive chondrogenic induction by the addition of TGFβ1. Thus, the combined chondrogenic induction with hypoxia and hyperosmolarity, resulting in a more gradual chondrogenic differentiation, might be preferred.

ECM deposition was compared by visualizing proteoglycan formation with Alcian Blue staining, and quantifying the sulphated GAG deposition in the ECM of the hydrogels and GAG excretion in the medium. The results of the Alcian blue staining seemed to correlate well with the GAG formation in the hydrogels, the hypoxia group staining most intensely and showing the highest GAG formation of all groups at both time-points, followed by the TGF β1-induced and non-induced group. This is in concordance with other studies, showing increased matrix biosynthesis (GAG production) in both chondrocytes

and mesenchymal stem cells when cultured under hypoxic conditions<sup>12,48</sup>. Interestingly, the GAG excretion patterns were clearly different between groups, i.e. whereas GAG excretion was increased in the non-induced, TGF $\beta$ 1-induced and hypoxic group over time, excretion showed a clear decrease in the hyperosmotic and combined groups between 7 and 14 days. This might be attributed to a less developed extracellular collagen type II deposition in these latter groups in the early period, thus hampering sequestration of the formed GAG in the ECM as also reported for osteoarthritic cartilage<sup>49,50</sup>, whereas after a week the ongoing ECM production allowed more efficient GAG encapsulation and concomitant lowered GAG excretion. However, our experimental conditions did not allow this hypothesis to be tested, since cells were cultured in collagen type II hydrogels. This made it impossible to distinguish between native collagen type II and collagen type II from the hydrogel with (immuno)histochemical visualisation or western blot analysis.

This study demonstrated an increase in chondrogenic genes and proteoglycan biosynthesis in response to combined hypoxia and hyperosmolarity, mimicking normal joint environment. However, osteoarthritic joints are characterized by proteoglycan loss, decreased collagen stiffness, and increased cartilage hydration (resulting in a hypo-osmotic environment). Consequently, the resulting increased cell volume of the residing chondrocytes makes them more sensitive to mechanical loading and cell death<sup>51,52</sup>. Although we did not test hypo-osmotic conditions in this study, other studies have shown that GAG production was low under hypo-osmolarity but high when varying the osmolarity between physiological levels<sup>53,54</sup>. This implies that in cellular therapies aimed at curing the osteoarthritic joint, restoration of a GAG sequestering network and re-establishment of a hyperosmotic microenvironment are both important parameters to be taken into account.

From this study it can be concluded that ASCs in a collagen type II hydrogel can differentiate towards the chondrogenic lineage, as investigated by qRT-PCR and histological and biochemical analysis of proteoglycans. Moreover, the combined application of both hyperosmolarity and hypoxia, two biophysicochemical cues present under physiological conditions in the joint environment, is able to induce chondrogenesis to the same level as induction TGF $\beta$ 1. Therefore, these data provide valuable insights for the development of stem cell based cartilage tissue engineering therapies. This might lead to an interesting alternative when considering short-term triggering in a one-step surgical procedure for the treatment of cartilaginous defects.

## REFERENCES

1. Hunziker EB, 2002. Articular cartilage repair: basic science and clinical progress. A review of the current status and prospects, *Osteoarthritis. Cartilage.*, **10**: 432-463
2. Brittberg M, 2010. Cell Carriers as the Next Generation of Cell Therapy for Cartilage Repair: A Review of the Matrix-Induced Autologous Chondrocyte Implantation Procedure, *Am.J.Sports Med.*, **38**: 1259-1271
3. Pittenger MF, Mosca JD, and McIntosh KR, 2000. Human mesenchymal stem cells: progenitor cells for cartilage, bone, fat and stroma, *Curr.Top.Microbiol.Immunol.*, **251**: 3-11
4. Lee HY, Kopesky PW, Plaas A, Sandy J, Kisiday J, Frisbie D, Grodzinsky AJ, and Ortiz C, 2010. Adult bone marrow stromal cell-based tissue-engineered aggrecan exhibits ultrastructure and nanomechanical

- properties superior to native cartilage, *Osteoarthritis Cartilage*, **18**: 1477-1486
5. Kopesky PW, Lee HY, Vanderploeg EJ, Kisiday JD, Frisbie DD, Plaas AH, Ortiz C, and Grodzinsky AJ, 2010. Adult equine bone marrow stromal cells produce a cartilage-like ECM mechanically superior to animal-matched adult chondrocytes, *Matrix Biol.*, **29**: 427-438
  6. Barry FP and Murphy JM, 2004. Mesenchymal stem cells: clinical applications and biological characterization, *Int.J.Biochem. Cell Biol.*, **36**: 568-584
  7. Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, Benhaim P, Lorenz HP, and Hedrick MH, 2001. Multilineage cells from human adipose tissue: implications for cell-based therapies, *Tissue Eng*, **7**: 211-228
  8. Helder MN, Knippenberg M, Klein-Nulend J, and Wuisman PI, 2007. Stem Cells from Adipose Tissue Allow Challenging New Concepts for Regenerative Medicine, *Tissue Eng*, **13**: 1799-1808
  9. Jurgens WJ, van DA, Doulabi BZ, Niessen FB, Ritt MJ, van Milligen FJ, and Helder MN, 2009. Freshly isolated stromal cells from the infrapatellar fat pad are suitable for a one-step surgical procedure to regenerate cartilage tissue, *Cytotherapy*, **11**: 1052-1064
  10. Puetzer JL, Petite JN, and Loba EG, 2010. Comparative Review of Growth Factors for Induction of Three-Dimensional In Vitro Chondrogenesis in Human Mesenchymal Stem Cells Isolated from Bone Marrow and Adipose Tissue, *Tissue Eng Part B Rev.*, **16**: 435-444
  11. Follmar KE, Decroos FC, Prichard HL, Wang HT, Erdmann D, and Olbrich KC, 2006. Effects of glutamine, glucose, and oxygen concentration on the metabolism and proliferation of rabbit adipose-derived stem cells, *Tissue Eng*, **12**: 3525-3533
  12. Merceron C, Vinatier C, Portron S, Masson M, Amiaud J, Guigand L, Chel Y, Weiss P, and Guicheux J, 2009. The differential effects of hypoxia on the osteochondrogenic potential of human adipose-derived stem cells, *Am.J.Physiol Cell Physiol*,
  13. Schumann D, Kujat R, Nerlich M, and Angele P, 2006. Mechanobiological conditioning of stem cells for cartilage tissue engineering, *Biomed.Mater.Eng*, **16**: S37-S52
  14. Elder BD and Athanasiou KA, 2009. Hydrostatic pressure in articular cartilage tissue engineering: from chondrocytes to tissue regeneration, *Tissue Eng Part B Rev.*, **15**: 43-53
  15. Cui JH, Park SR, Park K, Choi BH, and Min BH, 2007. Preconditioning of mesenchymal stem cells with low-intensity ultrasound for cartilage formation in vivo, *Tissue Eng*, **13**: 351-360
  16. Moore KA and Lemischka IR, 2006. Stem cells and their niches, *Science*, **311**: 1880-1885
  17. Wang DW, Fermor B, Gimble JM, Awad HA, and Guilak F, 2005. Influence of oxygen on the proliferation and metabolism of adipose derived adult stem cells, *J.Cell Physiol*, **204**: 184-191
  18. Khan WS, Adesida AB, and Hardingham TE, 2007. Hypoxic conditions increase hypoxia-inducible transcription factor 2alpha and enhance chondrogenesis in stem cells from the infrapatellar fat pad of osteoarthritis patients, *Arthritis Res.Ther.*, **9**: R55
  19. Murphy CL and Sambanis A, 2001. Effect of oxygen tension and alginate encapsulation on restoration of the differentiated phenotype of passaged chondrocytes, *Tissue Eng*, **7**: 791-803
  20. Scherer K, Schunke M, Sellckau R, Hassenpflug J, and Kurz B, 2004. The influence of oxygen and hydrostatic pressure on articular chondrocytes and adherent bone marrow cells in vitro, *Biorheology*, **41**: 323-333
  21. Urban JP, Maroudas A, Bayliss MT, and Dillon J, 1979. Swelling pressures of proteoglycans at the concentrations found in cartilaginous tissues, *Biorheology*, **16**: 447-464
  22. Urban JP, Hall AC, and Gehl KA, 1993. Regulation of matrix synthesis rates by the ionic and osmotic environment of articular chondrocytes, *J.Cell Physiol*, **154**: 262-270
  23. van der Windt AE, Haak E, Das RH, Kops N, Welting TJ, Caron MM, van Til NP, Verhaar JA, Weinans H, and Jahr H, 2010. Physiological tonicity improves human chondrogenic marker expression through nuclear factor of activated T-cells 5 in vitro, *Arthritis Res.Ther.*, **12**: R100
  24. Wuertz K, Urban JP, Klase J, Ignatius A, Wilke HJ, Claes L, and Neidlinger-Wilke C, 2007. Influence of extracellular osmolarity and mechanical stimulation on gene expression of intervertebral disc cells, *J.Orthop.Res.*, **25**: 1513-1522
  25. Wuertz K, Godburn K, Neidlinger-Wilke C, Urban J, and Iatridis JC, 2008. Behavior of mesenchymal stem cells in the chemical microenvironment of the intervertebral disc, *Spine*, **33**: 1843-1849

26. Hoogendoorn RJ, Lu ZF, Kroeze RJ, Bank RA, Wuisman PI, and Helder MN, 2008. Adipose stem cells for intervertebral disc regeneration: current status and concepts for the future, *J.Cell Mol.Med.*, **12**: 2205-2216
27. Malladi P, Xu Y, Chiou M, Giaccia AJ, and Longaker MT, 2006. Effect of reduced oxygen tension on chondrogenesis and osteogenesis in adipose-derived mesenchymal cells, *Am.J.Physiol Cell Physiol*, **290**: C1139-C1146
28. Discher DE, Janmey P, and Wang YL, 2005. Tissue cells feel and respond to the stiffness of their substrate, *Science*, **310**: 1139-1143
29. Engler AJ, Sen S, Sweeney HL, and Discher DE, 2006. Matrix elasticity directs stem cell lineage specification, *Cell*, **126**: 677-689
30. Lu ZF, Zandieh DB, Huang CL, Bank RA, and Helder MN, 2008. Beta1 integrins regulate chondrogenesis and rock signaling in adipose stem cells, *Biochem.Biophys.Res. Commun.*, **372**: 547-552
31. Oedayrajsingh-Varma MJ, van Ham SM, Knippenberg M, Helder MN, Klein-Nulend J, Schouten TE, Ritt MJ, and van Milligen FJ, 2006. Adipose tissue-derived mesenchymal stem cell yield and growth characteristics are affected by the tissue-harvesting procedure, *Cytotherapy*, **8**: 166-177
32. Lu Z, Doulabi BZ, Huang C, Bank RA, and Helder MN, 2010. Collagen type II enhances chondrogenesis in adipose tissue-derived stem cells by affecting cell shape, *Tissue Eng Part A*, **16**: 81-90
33. Hopewell B and Urban JP, 2003. Adaptation of articular chondrocytes to changes in osmolality, *Biorheology*, **40**: 73-77
34. Solchaga LA, Dennis JE, Goldberg VM, and Caplan AI, 1999. Hyaluronic acid-based polymers as cell carriers for tissue-engineered repair of bone and cartilage, *J.Orthop.Res.*, **17**: 205-213
35. Cancedda R, Castagnola P, Cancedda FD, Dozin B, and Quarto R, 2000. Developmental control of chondrogenesis and osteogenesis, *Int.J.Dev.Biol.*, **44**: 707-714
36. Wezeman FH, 1998. Morphological foundations of precartilage development in mesenchyme, *Microsc.Res.Tech.*, **43**: 91-101
37. Hall BK and Miyake T, 1992. The membranous skeleton: the role of cell condensations in vertebrate skeletogenesis, *Anat.Embryol.(Berl)*, **186**: 107-124
38. DeLise AM, Fischer L, and Tuan RS, 2000. Cellular interactions and signaling in cartilage development, *Osteoarthritis. Cartilage*, **8**: 309-334
39. Goldring MB, Tsuchimochi K, and Ijiri K, 2006. The control of chondrogenesis, *J.Cell Biochem.*, **97**: 33-44
40. Randolph MA, Anseth K, and Yaremchuk MJ, 2003. Tissue engineering of cartilage, *Clin.Plast.Surg.*, **30**: 519-537
41. Woodfield TB, Bezemer JM, Pieper JS, Van Blitterswijk CA, and Riesle J, 2002. Scaffolds for tissue engineering of cartilage, *Crit Rev. Eukaryot. Gene Expr.*, **12**: 209-236
42. Bosnakovski D, Mizuno M, Kim G, Takagi S, Okumura M, and Fujinaga T, 2006. Chondrogenic differentiation of bovine bone marrow mesenchymal stem cells (MSCs) in different hydrogels: influence of collagen type II extracellular matrix on MSC chondrogenesis, *Biotechnol.Bioeng.*, **93**: 1152-1163
43. Lu ZF, Doulabi BZ, Wuisman PI, Bank RA, and Helder MN, 2008. Influence of collagen type II and nucleus pulposus cells on aggregation and differentiation of adipose tissue-derived stem cells, *J.Cell Mol.Med.*, **12**: 2812-2822
44. Lu ZF, Zandieh DB, Wuisman PI, Bank RA, and Helder MN, 2007. Differentiation of adipose stem cells by nucleus pulposus cells: configuration effect, *Biochem.Biophys.Res. Commun.*, **359**: 991-996
45. Cheng NC, Estes BT, Awad HA, and Guilak F, 2009. Chondrogenic differentiation of adipose-derived adult stem cells by a porous scaffold derived from native articular cartilage extracellular matrix, *Tissue Eng Part A*, **15**: 231-241
46. Hellingman CA, Koevoet W, Kops N, Farrell E, Jahr H, Liu W, de Jong RJ, Frenz DA, and van Osch GJ, 2010. Fibroblast growth factor receptors in in vitro and in vivo chondrogenesis: relating tissue engineering using adult mesenchymal stem cells to embryonic development, *Tissue Eng Part A*, **16**: 545-556
47. Stevens DA, Hasserjian RP, Robson H, Siebler T, Shalet SM, and Williams GR, 2000. Thyroid hormones regulate hypertrophic chondrocyte differentiation and expression of parathyroid hormone-related peptide and its receptor during endochondral bone formation, *J.Bone Miner.Res.*, **15**: 2431-2442
48. Fermor B, Christensen SE, Youn I, Cernanec JM, Davies CM, and Weinberg JB, 2007.

- Oxygen, nitric oxide and articular cartilage, *Eur.Cell Mater.*, **13**: 56-65
49. Huebner JL, Williams JM, Deberg M, Henrotin Y, and Kraus VB, 2009. Collagen fibril disruption occurs early in primary guinea pig knee osteoarthritis, *Osteoarthritis.Cartilage.*,
  50. Kaplan LD, Lu Y, Snitzer J, Nemke B, Hao Z, Biro S, Albiero W, Stampfli HF, Markel M, Popkin C, and Baum SZ, 2009. The effect of early hyaluronic acid delivery on the development of an acute articular cartilage lesion in a sheep model, *Am.J.Sports Med.*, **37**: 2323-2327
  51. Bush PG, Hodkinson PD, Hamilton GL, and Hall AC, 2005. Viability and volume of in situ bovine articular chondrocytes-changes following a single impact and effects of medium osmolarity, *Osteoarthritis. Cartilage.*, **13**: 54-65
  52. Negoro K, Kobayashi S, Takeno K, Uchida K, and Baba H, 2008. Effect of osmolarity on glycosaminoglycan production and cell metabolism of articular chondrocyte under three-dimensional culture system, *Clin.Exp. Rheumatol.*, **26**: 534-541
  53. Chao PH, West AC, and Hung CT, 2006. Chondrocyte intracellular calcium, cytoskeletal organization, and gene expression responses to dynamic osmotic loading, *Am.J.Physiol Cell Physiol*, **291**: C718-C725
  54. Heywood HK, Bader DL, and Lee DA, 2006. Rate of oxygen consumption by isolated articular chondrocytes is sensitive to medium glucose concentration, *J.Cell Physiol*, **206**: 402-410